

ON CHARACTERS OF $L_{\mathfrak{sl}_n}(-\Lambda_0)$ -MODULES

KATHRIN BRINGMANN, KARL MAHLBURG, ANTUN MILAS

ABSTRACT. We use recent results of Rolin, Zwegers, and the first author to study characters of irreducible (highest weight) modules for the vertex operator algebra $L_{\mathfrak{sl}_\ell}(-\Lambda_0)$. We establish asymptotic behaviors of characters for the (ordinary) irreducible $L_{\mathfrak{sl}_\ell}(-\Lambda_0)$ -modules. As a consequence we prove that their quantum dimensions are one, as predicted by representation theory. We also establish a full asymptotic expansion of irreducible characters for \mathfrak{sl}_3 . Finally, we determine a decomposition formula for the full characters in terms of unary theta and false theta functions which allows us to study their modular properties.

1. INTRODUCTION AND STATEMENT OF RESULTS

Vertex operator algebras have had a profound influence on mathematics and modern theoretical physics. Arguably, the most important examples of vertex algebras are those associated to representations of the affine Kac-Moody Lie algebras at positive integral levels. Every such vertex algebra is rational [12], and its characters are modular functions on the same congruence subgroup. Moreover, characters of representations of affine vertex algebras at certain rational levels, called admissible, are also modular [14].

1.1. Affine vertex algebras at negative levels. More recently, a notable effort has been made to understand modules of affine vertex algebras at a negative integer level [1, 2, 3, 4, 16], etc. Although there is no general Weyl-Kac type formula at these levels, in a few cases their characters (or q -characters) are known explicitly. Here a prominent role is played by the so-called Deligne series of representations at the level $-\frac{h^\vee}{6} - 1$ for the Lie algebras of type D_4, E_6, E_7 , and E_8 [3, 16]. These vertex algebras are irrational and quasi-lisse, and thus their characters are quasi-modular and are also solutions of certain modular linear differential equations [3]. The search for new examples of quasi-lisse (and lisse) vertex algebras is a very active area of current research with important applications to $N = 2$ four dimensional CFT [5] and logarithmic CFT [10].

Apart from Deligne's series, specialized characters of $L_{\mathfrak{sl}_n}(-\Lambda_0)$ -modules have been studied in several recent works [6, 16]. This vertex algebra is no longer quasi-lisse but their characters still enjoy interesting "quantum" modular properties. More precisely, it was proven in [6, 8] that the irreducible characters of level -1 are mixed quantum modular forms by realizing them as Fourier coefficients of meromorphic Jacobi forms of negative index.

1.2. Quantum dimensions and asymptotics. Somewhat independent of these developments, in [10, 11] the authors initiated a study of asymptotic expansions of characters of modules for irrational vertex algebras. Motivated by the success of the theory of rational vertex algebras [13], it is expected that asymptotic expansions of characters can be used to formulate a Verlinde-type formula for the fusion rules. For the singlet vertex algebra and generalizations this was recently established by using the so-called regularized quantum dimension [10]. Despite the progress made it is still unclear how to formulate a Verlinde-type formula for more general vertex algebras. The main issue is that vertex algebras can have representations with very different properties, which makes it difficult to define quantum dimensions, S -matrices, and other standard invariants. But for vertex algebras whose irreducible modules are C_1 -cofinite we expect that asymptotically (as $t \rightarrow 0^+$)

$$\frac{\text{ch}_M(it)}{\text{ch}_V(it)} \sim a_0 + a_1 t + \dots \quad (1.1)$$

for any irreducible V -module M . In particular, (1.1) would imply $\lim_{t \rightarrow 0^+} \frac{\text{ch}_M(it)}{\text{ch}_V(it)} = a_0$, so we have a well-defined notion of quantum dimension (the existence of the limit does not necessarily require (1.1) though). In order to gain a better insight into irrational vertex algebras it is desirable to study asymptotic expansion of characters, and hence of (1.1), for various families of affine vertex algebras (for related W -algebra computations see [10]).

1.3. Main results. In this paper we study asymptotic and modular properties of characters of the vertex algebra $L_{\mathfrak{sl}_n}(-\Lambda_0)$ -modules, $n \geq 3$. This vertex algebra is a good representative of an irrational affine vertex algebra admitting both atypical and typical modules. We only consider atypical representations, leaving Verlinde-type formula for future considerations.

Let us outline the main results. For brevity, we let $\mathfrak{g} = \mathfrak{sl}_\ell$, $\ell \geq 3$, the Lie algebra of trace zero $\ell \times \ell$ matrices. As usual, from \mathfrak{g} we construct the affine Kac-Moody Lie algebra $\widehat{\mathfrak{g}}$. Denote by ω_j , $j \in \{1, \dots, \ell - 1\}$, the fundamental weights of \mathfrak{g} and by Λ_j , $j \in \{0, \dots, \ell - 1\}$, the fundamental weights for $\widehat{\mathfrak{g}}$. For each weight Λ , let $L_{\mathfrak{g}}(\Lambda)$ be the corresponding irreducible highest weight $\widehat{\mathfrak{g}}$ -module.

We equip $L_{\mathfrak{g}}(-\Lambda_0)$ with a conformal vertex algebra structure via Sugawara's construction. It is known by Adamovic and Perse [1] that $L_{\mathfrak{g}}(-\Lambda_0)$ admits countably many inequivalent ordinary modules: $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_1)$ with $s \in \mathbb{N}_0$ and $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_{\ell-1})$ with $s \in \mathbb{N}$, and infinitely many generic modules [2]. We denote the degree operator by $L(0)$ and the central charge by $c := \frac{\dim(\mathfrak{g})(-1)}{(-1)+h^\vee} = -(\ell+1)$, where $h^\vee = \ell$ is the dual Coxeter number of \mathfrak{sl}_ℓ . We use this data to define the relevant characters (here $\Lambda \in \{\Lambda_1, \Lambda_{\ell-1}\}$ and $q := e^{2\pi i\tau}$):

$$\text{ch}[L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda)](\tau) := \text{tr}_{L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda)} q^{L(0) - \frac{c}{24}}.$$

Let

$$h_s := \frac{(s\omega_1 + 2\rho, s\omega_1)}{2((-1) + h^\vee)} = \frac{(s\omega_{\ell-1} + 2\rho, s\omega_{\ell-1})}{2((-1) + h^\vee)} = \frac{s^2}{2\ell} + \frac{s}{2}$$

denote the lowest conformal weight of $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_1)$ and of $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_{\ell-1})$. Note that $L_{\mathfrak{g}}(-\Lambda_0(s+1) + s\Lambda_1)$ and $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_{\ell-1})$ are dual to each other so they share the same character.

By using results of Kac and Wakimoto on specialized characters of the affine Lie algebras $\widehat{\mathfrak{sl}}_\ell$ at level -1 [16, Section 1], we obtain, for $s \in \mathbb{N}_0$ and $\Lambda \in \{\Lambda_1, \Lambda_{\ell-1}\}$,

$$\text{ch}[L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda)](\tau) = q^{h_s - \frac{c}{24}} \frac{F_{\ell,s}(q)}{(q)_\infty^{\ell^2-1}}, \quad (1.2)$$

where

$$F_{\ell,s}(q) := (q)_\infty^{\ell^2-1} \text{coeff}_{\zeta^s} \frac{(q)_\infty}{(\zeta)_\infty^\ell (\zeta^{-1}q)_\infty^\ell}.$$

Here $(a)_\infty = (a; q)_\infty := \prod_{j=0}^{\infty} (1 - aq^j)$. Throughout, we are interested in the range $0 < y < v$, where we write $z = x + iy$, $\tau = u + iv$. We note that other ranges can be treated in a similar way.

Theorem 1.1. *We have, as $t \rightarrow 0^+$,*

$$F_{\ell,s}(e^{-t}) \sim \mathcal{C}_\ell \left(\frac{t}{2\pi} \right)^{1 - \frac{\ell^2}{2}} e^{-\frac{\pi^2(\ell^2-2\ell)}{6t}},$$

where

$$\mathcal{C}_\ell := \frac{2^{1-2\ell}(\ell-1)!}{\Gamma\left(\frac{\ell+1}{2}\right)^2}.$$

The following corollary follows directly (the same results apply for their dual modules, $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_{\ell-1})$ as their q -characters are equal). Let V be a vertex algebra. For a V -module M we define its *quantum* or *asymptotic dimension*:

$$\text{qdim}(M) := \lim_{t \rightarrow 0^+} \frac{\text{ch}[M](it)}{\text{ch}[V](it)}.$$

Corollary 1.2. *Let $\mathfrak{g} = \mathfrak{sl}_\ell$, $\ell \geq 3$ and $s \in \mathbb{N}_0$. Let V_s denote $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_1)$ or $L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_{\ell-1})$. Then we have, as $t \rightarrow 0^+$,*

$$\text{ch}[V_s](e^{-t}) \sim \mathcal{C}_\ell \sqrt{\frac{t}{2\pi}} e^{\frac{\pi^2(2\ell-1)}{6t}}. \quad (1.3)$$

In particular $\text{qdim}(V_s) = 1$ for every ℓ and s .

In fact, the method used to prove Theorem 1.1 should in principle lead to the full asymptotic expansion for any such character. In Proposition 6.1 below, we carry out the details for the case $\ell = 3$.

Now we recall Kac and Wakimoto's work in [16, Section 1], which gives the full character as the coefficients of certain series. Let $\zeta_j := e^{\alpha_j}$ and define

$$F_\ell(\zeta_1, \dots, \zeta_\ell) := (q)_\infty \prod_{j=1}^{\ell} \prod_{k=1}^{\infty} \frac{1}{(1 - \zeta_j^{-1} \dots \zeta_\ell^{-1} q^k) (1 - \zeta_j \dots \zeta_\ell q^{k-1})}.$$

Using the geometric series on every term, we expand this product as a formal series, to obtain a decomposition of the shape

$$F_\ell(\zeta_1, \dots, \zeta_\ell) =: \sum_{s \in \mathbb{Z}} F_{\ell,s}(\zeta_1, \dots, \zeta_{\ell-1}) \zeta_\ell^s.$$

In other words, $F_{\ell,s}(\zeta_1, \dots, \zeta_{\ell-1})$ is defined to be the s -th Fourier coefficient of F_ℓ with respect to ζ_ℓ . In [16], Kac and Wakimoto showed that the $F_{\ell,s}$ are essentially the irreducible characters of $L_{\mathfrak{g}}(-\Lambda_0)$ -modules. More precisely ($\zeta_j := e^{2\pi i z_j}$)

$$\begin{aligned} F_\ell(\zeta_1, \dots, \zeta_\ell) &= \sum_{s=1}^{\infty} (\zeta_1 \dots \zeta_\ell)^s q^{\frac{3s}{2}} \tilde{\text{ch}}[-(1+s)\Lambda_0 + s\Lambda_1](z; \tau) \\ &\quad + \sum_{s=0}^{\infty} \zeta_\ell^{-s} q^{-\frac{s}{2}} \tilde{\text{ch}}[-(1+s)\Lambda_0 + s\Lambda_{\ell-1}](z; \tau), \end{aligned} \quad (1.4)$$

where $\tilde{\text{ch}}[M](z; \tau)$ is the full character of M and $\zeta = (\zeta_1, \dots, \zeta_{\ell-1})$. Note that our definition differs from [16] due to the shift $\zeta_\ell \mapsto \zeta_\ell q^{\frac{1}{2}}$, which affects the q -powers in (1.4).

In Section 7 we prove the following decomposition result for $F_{\ell,s}(\zeta_1, \dots, \zeta_{\ell-1})$.

Theorem 1.3. *Assume that $w_j := -\sum_{k=j}^{\ell-1} z_k$ for $1 \leq j \leq \ell - 1$, and $w_\ell = 0$ are all distinct. We have, for $|q| < |\zeta_j|^\ell < 1$, $1 \leq j \leq \ell - 1$,*

$$F_{\ell,s}(\zeta_1, \dots, \zeta_{\ell-1}) = -i^{\ell+1} q^{-h_s + \frac{c}{24}} \eta(\tau)^{\ell-2} \prod_{j=1}^{\ell-1} \zeta_j^{\frac{js}{\ell}} \sum_{\nu=1}^{\ell} \frac{\vartheta_{s-\frac{\ell}{2}, \ell, \frac{\ell}{2}}^+(w_\nu - \frac{1}{\ell} \sum_{j=1}^{\ell} w_j)}{\prod_{\substack{j=1 \\ j \neq \nu}}^{\ell} \vartheta(w_\nu - w_j)}.$$

where ϑ and $\vartheta_{s-\frac{\ell}{2}, \ell, \frac{\ell}{2}}^+$ are certain theta and partial theta functions defined in (2.6) and (4.2), respectively.

Kac and Wakimoto [16] recently obtained a Weyl-Kac type formula for the character $\text{ch}[-(1+s)\Lambda_0 + s\Lambda](z; \tau)$. Unlike our Theorem 1.3, where the full character splits into rank one pieces, their formula involves a summation over the half-space of the root lattice of \mathfrak{sl}_ℓ (see also [6]). Although very elegant, we found their result more difficult to use for the purpose of computing asymptotic expansions of full characters.

1.4. Organization of the paper. The paper is organized as follows. In Section 2 we recall basic facts about Bernoulli numbers and their generalizations, theta functions, and the Euler-Maclaurin summation formula. We then study in Section 3 the asymptotic behaviour of certain auxiliary functions. Section 4 decomposes $F_{\ell,s}$ in terms of partial theta functions and quasimodular forms. In Section 5, we then finish the proof of Theorem 1.1 and Corollary 1.2. Section 6 treats the case $\ell = 3$ in which we find a full asymptotic expansion for the corresponding characters. We prove a decomposition formula of $F_{\ell,s}(\zeta_1, \dots, \zeta_{\ell-1})$ in Section 7. We finish the paper in Section 8 by computing modular transformation properties of the full character $F_{\ell,s}$ from modular properties of unary partial and false theta functions, which we also derive there.

ACKNOWLEDGMENTS

We thank Victor Kac for raising the question of computing asymptotic expansion of characters and Walter Bridges for his helpful comments. We moreover thank the referee for helpful comments.

The research of the first author is supported by the Alfred Krupp Prize for Young University Teachers of the Krupp foundation and the research leading to these results receives funding from the European Research Council under the European Union's Seventh Framework Programme (FP/2007-2013) / ERC Grant agreement n. 335220 - AQSER. The research of the third author was supported by a Simons Foundation Collaboration Grant (# 317908) and NSF-DMS Grant 1601070.

2. PRELIMINARIES

2.1. Bernoulli number generating functions. Recall that the *Bernoulli numbers* B_k are defined via their generating function

$$\frac{w}{e^w - 1} =: \sum_{k=0}^{\infty} B_k \frac{w^k}{k!}. \quad (|w| < 2\pi) \quad (2.1)$$

We also require the following modified generating function, again assuming $|w| < 2\pi$,

$$S(w) := \sum_{k=1}^{\infty} \frac{B_{2k} w^{2k}}{2k(2k)!}. \quad (2.2)$$

Differentiating (2.2) and using (2.1) gives that

$$S'(w) = \frac{e^{-w}}{1 - e^{-w}} - \frac{1}{w} + \frac{1}{2}$$

and hence, by taking the limit on both sides,

$$S(w) = \text{Log} \left(\frac{e^w - 1}{w} \right) - \frac{w}{2}. \quad (2.3)$$

We also need generalizations of Bernoulli numbers. For this, denote by $B_n^{(r)}(x)$ the *Bernoulli polynomials of order* $r \in \mathbb{N}$, defined via their generating function

$$\left(\frac{w}{e^w - 1}\right)^r e^{xw} =: \sum_{n=0}^{\infty} B_n^{(r)}(x) \frac{w^n}{n!}.$$

For $r = 1$ this recovers the ordinary Bernoulli polynomials $B_n(x)$.

We also require the *Euler polynomials* defined through the generating function

$$\frac{2e^{xw}}{e^w + 1} =: \sum_{n=0}^{\infty} \frac{E_n(x)}{n!} w^n.$$

The *Euler numbers* are in particular given by

$$E_n = 2^n E_n\left(\frac{1}{2}\right). \quad (2.4)$$

The following identity relates Euler and Bernoulli polynomials ($n \in \mathbb{N}, m \in 2\mathbb{N}$)

$$E_n(mx) = -\frac{2}{n+1} m^n \sum_{k=0}^{m-1} (-1)^k B_{n+1}\left(x + \frac{k}{m}\right). \quad (2.5)$$

2.2. Theta functions. Let

$$\vartheta(z) = \vartheta(z; \tau) := \sum_{n \in \frac{1}{2} + \mathbb{Z}} q^{\frac{n^2}{2}} e^{2\pi i n(z + \frac{1}{2})}, \quad \eta(\tau) := q^{\frac{1}{24}}(q)_{\infty}. \quad (2.6)$$

By the Jacobi triple product identity, we have the product expansion

$$\vartheta(z; \tau) = -iq^{\frac{1}{8}} \zeta^{-\frac{1}{2}}(q)_{\infty}(\zeta)_{\infty}(\zeta^{-1}q)_{\infty}.$$

Recall the transformations ($\lambda, \mu \in \mathbb{Z}, \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$)

$$\begin{aligned} \vartheta(z + \lambda\tau + \mu) &= (-1)^{\lambda + \mu} q^{-\frac{\lambda^2}{2}} \zeta^{-\lambda} \vartheta(z), \\ \vartheta\left(\frac{z}{c\tau + d}; \frac{a\tau + b}{c\tau + d}\right) &= \chi \begin{pmatrix} a & b \\ c & d \end{pmatrix} (c\tau + d)^{\frac{1}{2}} e^{\frac{\pi i c z^2}{c\tau + d}} \vartheta(z; \tau), \end{aligned} \quad (2.7)$$

where χ is the multiplier of η^3 . In particular

$$\vartheta\left(\frac{z}{\tau}; -\frac{1}{\tau}\right) = -i\sqrt{-i\tau} e^{\frac{\pi i z^2}{\tau}} \vartheta(z; \tau), \quad \eta\left(-\frac{1}{\tau}\right) = \sqrt{-i\tau} \eta(\tau). \quad (2.8)$$

Next recall that

$$\vartheta(z; \tau) = -2\pi z \cdot \eta(\tau)^3 \cdot \exp\left(-\sum_{k=1}^{\infty} \frac{G_{2k}(\tau)}{2k} z^{2k}\right), \quad (2.9)$$

where for $k \in \mathbb{N}$

$$G_{2k}(\tau) := -\frac{(2\pi i)^{2k} B_{2k}}{(2k)!} + \frac{2(2\pi i)^{2k}}{(2k-1)!} \sum_{n=1}^{\infty} \sigma_{2k-1}(n) q^n. \quad (2.10)$$

Here $\sigma_\ell(n) := \sum_{d|n} d^\ell$ is the ℓ -th divisor sum. We also require the transformations ($k > 1$)

$$G_{2k}(\tau) = \tau^{-2k} G_{2k}\left(-\frac{1}{\tau}\right) \quad G_2(\tau) = \tau^{-2} G_2\left(-\frac{1}{\tau}\right) + \frac{2\pi i}{\tau}. \quad (2.11)$$

2.3. Euler-Maclaurin summation formula. The Euler-Maclaurin summation formula (see e.g. [18], correcting a sign error) implies that for $\alpha \in \mathbb{R}$ and $f : \mathbb{R} \rightarrow \mathbb{R}$ a C^∞ -function such that f and all its derivatives are of rapid decay at infinity, we have

$$\sum_{n=0}^{\infty} f((n+\alpha)t) = \frac{I_f}{t} - \sum_{n=0}^N \frac{B_{n+1}(\alpha)}{(n+1)!} f^{(n)}(0) t^n + O(t^{N+1}) \quad (t \rightarrow 0^+), \quad (2.12)$$

where $I_f := \int_0^\infty f(x) dx$.

3. ASYMPTOTICS FOR AUXILIARY FUNCTIONS

We next determine the asymptotic behaviour of two partial theta functions. These expansions are well-known to experts; however, for the convenience of the reader, we provide proofs.

The first function we require is (here $j \in \mathbb{N}_0$, $r \in \mathbb{Q}$),

$$\mathcal{F}_{j,r}(t) := 2^{-2j} t^j \sum_{n=0}^{\infty} (-1)^n (n+r)^{2j} e^{-\frac{1}{4}(n+r)^2 t}. \quad (3.1)$$

Then

$$\mathcal{F}_{j,r}(t) = \sum_{n=0}^{\infty} \left(\mathbb{F}_j\left(\left(n + \frac{r}{2}\right) \sqrt{t}\right) - \mathbb{F}_j\left(\left(n + \frac{r+1}{2}\right) \sqrt{t}\right) \right),$$

where $\mathbb{F}_j(x) := x^{2j} e^{-x^2}$. By (2.12), we obtain

$$\mathcal{F}_{j,r}(t) = - \sum_{n=0}^N \frac{B_{n+1}\left(\frac{r}{2}\right) - B_{n+1}\left(\frac{r+1}{2}\right)}{(n+1)!} \mathbb{F}_j^{(n)}(0) t^{\frac{n}{2}} + O\left(t^{\frac{N+1}{2}}\right). \quad (3.2)$$

Noting that \mathbb{F}_j is an even function, we compute

$$\mathcal{F}_{j,r}(t) = - \sum_{n=0}^N \frac{B_{2n+2j+1}\left(\frac{r}{2}\right) - B_{2n+2j+1}\left(\frac{r+1}{2}\right)}{(2n+2j+1)n!} (-1)^n t^{n+j} + O\left(t^{N+j+1}\right). \quad (3.3)$$

This implies that

$$\mathcal{F}_{j,r}(t) \sim -\frac{B_{2j+1}\left(\frac{r}{2}\right) - B_{2j+1}\left(\frac{r+1}{2}\right)}{2j+1} t^j. \quad (3.4)$$

In particular,

$$\mathcal{F}_{0,r}(t) \sim \frac{1}{2}. \quad (3.5)$$

We next study, for $j \in \mathbb{N}$, $r \in \mathbb{Q}$,

$$\mathcal{G}_{j,r}(t) := t^{j-\frac{1}{2}} \sum_{n=0}^{\infty} (n+r)^{2j-1} e^{-(n+r)^2 t}. \quad (3.6)$$

Then

$$\mathcal{G}_{j,r}(t) = \sum_{n=0}^{\infty} \mathbb{G}_j \left((n+r) \sqrt{t} \right),$$

where $\mathbb{G}_j(x) := x^{2j-1} e^{-x^2}$. By (2.12) and the fact that $I_{\mathbb{G}_j} = \frac{(j-1)!}{2}$, we obtain that

$$\mathcal{G}_{j,r}(t) = \frac{I_{\mathbb{G}_j}}{\sqrt{t}} - \sum_{n=0}^N \frac{B_{n+1}(r)}{(n+1)!} \mathbb{G}_j^{(n)}(0) t^{\frac{n}{2}} + O\left(t^{\frac{N+1}{2}}\right) \sim \frac{(j-1)!}{2\sqrt{t}}. \quad (3.7)$$

4. A q -SERIES FOR $F_{\ell,s}$

To prove Theorem 1.1, we first find a q -series representation for $F_{\ell,s}$. For this, we use a theorem from [8]. To state it, we let, for $z_0 \in \mathbb{C}$,

$$P_{z_0} := z_0 + [0, 1) + \tau[0, 1).$$

Let ϕ be a meromorphic function satisfying $(\lambda, \mu \in \mathbb{Z}, \varepsilon \in \{0, 1\}, m \in -\frac{1}{2}\mathbb{N})$

$$\phi(z + \lambda\tau + \mu) = (-1)^{2m\mu + \lambda\varepsilon} e^{-2\pi i m(\lambda^2\tau + 2\lambda z)} \phi(z). \quad (4.1)$$

Moreover, let $S_{z_0}(\tau)$ be the set of poles of ϕ in P_{z_0} , which is discrete due to the fact that ϕ is meromorphic. Furthermore $\mathcal{D}_z := \frac{1}{2\pi i} \frac{\partial}{\partial z}$, and for $M \in \frac{1}{2}\mathbb{N}$, $r \in M + \mathbb{Z}$, $\varepsilon \in \{0, 1\}$, define the partial/false theta functions

$$\vartheta_{r,\varepsilon,M}^+(z) = \vartheta_{r,\varepsilon,M}^+(z; \tau) := \sum_{n=0}^{\infty} (-1)^{n\varepsilon} \zeta^{2Mn-r} q^{\frac{1}{4M}(2Mn-r)^2}. \quad (4.2)$$

For $r \in m + \mathbb{Z}$, let $h_{r,z_0}(\tau)$ be the renormalized r -th Fourier coefficient with respect to z_0

$$h_r(\tau) = h_{r,z_0}(\tau) := q^{-\frac{r^2}{4m}} \int_{z_0}^{z_0+1} \phi(z) e^{-2\pi i r z} dz. \quad (4.3)$$

If ϕ has any poles on the line from z_0 to $z_0 + 1$, then the contour must be deformed. In particular, if there is a pole that is not an endpoint, then we define the path to be the sum of the integrals over the paths deformed to pass above and below the pole, weighted by $\frac{1}{2}$. If there is a pole at an endpoint, we replace the path by $[z_0 - \delta, z_0 + 1 - \delta]$

with δ small, so that there is no pole at an endpoint. We now recall the following Theorem from [8] (slightly rewritten).

Theorem 4.1. *Assume that ϕ satisfies (4.1) and choose $z_0 \in \mathbb{C}$, such that ϕ has no poles on the boundary of P_{z_0} . Then we have, for any $r \in m + \mathbb{Z}$,*

$$h_{r,z_0}(\tau) = 2\pi i \sum_{z_s \in S_{z_0}(\tau)} \operatorname{Res}_{z=z_s} \left(\phi(z, \tau) \vartheta_{r,\varepsilon,-m}^+(z; \tau) \right). \quad (4.4)$$

We use Theorem 4.1 to find a q -series for $F_{\ell,s}$. For this, we define

$$g_\ell(z) = g_\ell(z; \tau) := \frac{\eta(\tau)^{3\ell}}{\vartheta(z; \tau)^\ell}.$$

Theorem 4.2. *We have, for $0 < y < v$,*

$$F_{\ell,s}(q) = i^\ell (q)_\infty^{\ell^2 - 2\ell} q^{-h_s - \frac{\ell}{8}} \sum_{\substack{1 \leq j \leq \ell \\ j \equiv \ell \pmod{2}}} \frac{D_{-j}(\tau)}{(j-1)!} \sum_{n=0}^{\infty} (-1)^{n\ell} \left(\ell n + \frac{\ell}{2} - s \right)^{j-1} q^{\frac{1}{2\ell}(\ell n + \frac{\ell}{2} - s)^2},$$

where the $D_{-j}(\tau)$ are the Laurent coefficients of g_ℓ in z (expanded around 0), so that

$$g_\ell(z; \tau) = \sum_{j=1}^{\ell} \frac{D_{-j}(\tau)}{(2\pi i z)^j} + O(1), \quad (4.5)$$

Proof: To apply (4.4), we rewrite

$$F_{\ell,s}(q) = (q)_\infty^{\ell^2 - 1} \operatorname{coeff}_{\zeta^s} \frac{(q)_\infty}{(\zeta)_\infty^\ell (\zeta^{-1}q)_\infty^\ell} = (q)_\infty^{\ell^2 - 2\ell} f_{\ell,s}(\tau)$$

where

$$f_{\ell,s}(\tau) := (-i)^\ell \operatorname{coeff}_{\zeta^{s+\frac{\ell}{2}}} g_\ell(z; \tau). \quad (4.6)$$

Note that g_ℓ is a Jacobi form of weight ℓ and index $m = -\frac{\ell}{2}$. In particular we have the elliptic transformation ($\lambda, \mu \in \mathbb{Z}$)

$$g_\ell(z + \lambda\tau + \mu) = (-1)^{(\lambda+\mu)\ell} \zeta^{\lambda\ell} q^{\frac{\ell\lambda^2}{2}} g_\ell(z). \quad (4.7)$$

Thus (4.1) holds with $\varepsilon \equiv \ell \pmod{2}$. The poles of $g_\ell(z)$ are the lattice $\mathbb{Z} + \mathbb{Z}\tau$. Moreover, for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{SL}_2(\mathbb{Z})$ we require the modular transformation

$$g_\ell \left(\frac{z}{c\tau + d}; \frac{a\tau + b}{c\tau + d} \right) = (c\tau + d)^\ell e^{-\frac{\pi i \ell c z^2}{c\tau + d}} g_\ell(z; \tau).$$

To rewrite (4.6), we use (4.4), to obtain

$$H_{s+\frac{\ell}{2}}(\tau) = 2\pi i \cdot \operatorname{Res}_{z=\tau} \left(g_\ell(z) \vartheta_{s+\frac{\ell}{2}, \varepsilon, \frac{\ell}{2}}^+(z; \tau) \right),$$

where $H_j(\tau)$ denotes the j -th renormalized $h_{j,z_0}(\tau)$ (see (4.3)) Fourier coefficient of $g_\ell(z; \tau)$ in the fixed range $0 < y < v$, and thus z_0 is chosen such that it satisfies the

conditions of Theorem 4.1 with $\tau \in P_{z_0}$. Using (4.7) with $\lambda = 1$ and $\mu = 0$ (and $z \mapsto z - \tau$), we directly obtain that

$$g_\ell(z) = (-1)^\ell \zeta^\ell q^{-\frac{\ell}{2}} g_\ell(z - \tau).$$

Plugging in, we find that

$$\begin{aligned} H_{s+\frac{\ell}{2}}(\tau) &= (-1)^\ell q^{-\frac{\ell}{2}} 2\pi i \operatorname{Res}_{z=\tau} \left(g_\ell(z - \tau; \tau) \zeta^\ell \vartheta_{s+\frac{\ell}{2}, \varepsilon, \frac{\ell}{2}}^+(z; \tau) \right) \\ &= (-1)^\ell q^{-\frac{\ell}{2}} \sum_{\substack{j=1 \\ j \equiv \ell \pmod{2}}}^{\ell} \frac{D_{-j}(\tau)}{(j-1)!} \left[\mathcal{D}_z^{j-1} \left(\zeta^\ell \vartheta_{s+\frac{\ell}{2}, \varepsilon, \frac{\ell}{2}}^+(z; \tau) \right) \right]_{z=\tau}. \end{aligned}$$

Computing the derivative

$$\left[\mathcal{D}_z^{j-1} \left(\zeta^\ell \vartheta_{s+\frac{\ell}{2}, \varepsilon, \frac{\ell}{2}}^+(z; \tau) \right) \right]_{z=\tau} = q^{\frac{\ell}{2}} \sum_{n=0}^{\infty} (-1)^{n\varepsilon} \left(\ell n + \frac{\ell}{2} - s \right)^{j-1} q^{\frac{1}{2\ell}(\ell n + \frac{\ell}{2} - s)^2}$$

then gives

$$H_{s+\frac{\ell}{2}}(\tau) = (-1)^\ell \sum_{\substack{1 \leq j \leq \ell \\ j \equiv \ell \pmod{2}}} \frac{D_{-j}(\tau)}{(j-1)!} \sum_{n=0}^{\infty} (-1)^{n\varepsilon} \left(\ell n + \frac{\ell}{2} - s \right)^{j-1} q^{\frac{1}{2\ell}(\ell n + \frac{\ell}{2} - s)^2}. \quad (4.8)$$

Noting that

$$F_{\ell, s}(q) = (-i)^\ell (q)_\infty^{\ell^2 - 2\ell} q^{-h_s - \frac{\ell}{8}} H_{s+\frac{\ell}{2}}(\tau) \quad (4.9)$$

finishes the claim. \square

5. PROOF OF THEOREM 1.1

5.1. Asymptotic series for quasimodular forms. Using (2.8) gives that

$$\eta\left(\frac{it}{2\pi}\right) \sim \sqrt{\frac{2\pi}{t}} e^{-\frac{\pi^2}{6t}} \quad (t \rightarrow 0^+). \quad (5.1)$$

Plugging this into (4.9) we then obtain

$$F_{\ell, s}(e^{-t}) \sim (-i)^\ell \left(\frac{2\pi}{t}\right)^{\frac{\ell^2}{2} - \ell} e^{-\frac{\pi^2}{12t}(\ell^2 - 2\ell)} H_{s+\frac{\ell}{2}}\left(\frac{it}{2\pi}\right). \quad (5.2)$$

To determine the asymptotic behavior of $H_{s+\frac{\ell}{2}}$, we first investigate the coefficients in form of D_{-j} in (4.8). Using (2.9), we write

$$g_\ell\left(z; \frac{it}{2\pi}\right) = \frac{1}{(-2\pi z)^\ell} \exp\left(\ell \sum_{k=1}^{\infty} \frac{G_{2k}\left(\frac{it}{2\pi}\right)}{2k} z^{2k}\right). \quad (5.3)$$

From this we see that in $g_\ell(z; \tau)$ every coefficient in front of z^{-j} has the shape

$$D_{-j}(\tau) = \sum_{m,k} c(m, k) G_{k_1}^{m_1}(\tau) \cdots G_{k_r}^{m_r}(\tau), \quad (5.4)$$

where the summation is over all tuples of positive integers (of any finite length) $m = (m_1, \dots, m_r), k = (k_1, \dots, k_r)$ such that $m_1 k_1 + \cdots + m_r k_r = \ell - j$ and the $c(m, k)$ are real constants depending on the tuple. By (2.11), we have that, for $k \geq 2$,

$$G_{2k} \left(\frac{it}{2\pi} \right) = \left(\frac{2\pi}{it} \right)^{2k} \left(-\frac{(2\pi i)^{2k} B_{2k}}{(2k)!} + \frac{2(2\pi i)^{2k}}{(2k-1)!} \sum_{n=1}^{\infty} \sigma_{2k-1}(n) e^{-\frac{4\pi^2 n}{t}} \right).$$

Thus we obtain, for $k \geq 2$,

$$G_{2k} \left(\frac{it}{2\pi} \right) \sim -\frac{\left(\frac{4\pi^2}{t} \right)^{2k} B_{2k}}{(2k)!}, \quad (5.5)$$

with an error term that decays exponentially as a function of $\frac{1}{t}$. For $G_2\left(\frac{it}{2\pi}\right)$ the main term is also given by (5.5), but now the error is only smaller by the polynomial factor t due to the additional term $\frac{4\pi^2}{t}$ in the transformation (2.11). Regardless, if we now plug (5.5) into (5.3), the accumulated error term in (5.4) is smaller than the main term by at least a polynomial factor of t .

Furthermore, note that it is valid to calculate any given Laurent coefficient in (5.3) by restricting z to a segment of the imaginary axis, so $z = iy$ with $0 < y < v = \frac{t}{2\pi}$. This is a subset of the canonical analytic domain $0 < y < v$. This restriction is convenient as it allows us to package all of the main asymptotic terms into a generating function, which can then be evaluated using the identities from Section 2. In particular, for the remainder of this section we write

$$f(z, t) \sim_z g(z, t)$$

for two functions

$$f(z, t) = \sum_{n \in \mathbb{Z}} a_t(n) z^n, \quad g(z, t) = \sum_{n \in \mathbb{Z}} b_t(n) z^n,$$

if for all n

$$a_t(n) \sim b_t(n) \quad (\text{as } t \rightarrow 0^+).$$

Using (2.3) (noting that $\frac{4\pi^2 y}{t} < 2\pi$), we find that for $z = iy$,

$$g_\ell \left(z; \frac{it}{2\pi} \right) \sim_z \frac{1}{(-2\pi z)^\ell} \exp \left(-\ell S \left(\frac{4\pi^2 z}{t} \right) \right) = \frac{1}{(-2\pi z)^\ell} \frac{\left(\frac{4\pi^2 z}{t} \right)^\ell e^{\frac{2\pi^2 \ell z}{t}}}{\left(e^{\frac{4\pi^2 z}{t}} - 1 \right)^\ell}. \quad (5.6)$$

We also require that D_{-j} is a quasimodular form of weight $\ell - j$ and thus satisfies (see (2.11))

$$D_{-j} \left(\frac{it}{2\pi} \right) \ll t^{j-\ell}. \quad (5.7)$$

We now distinguish 2 cases, depending on whether ℓ is even or odd.

5.2. Proof of Theorem 1.1 for ℓ odd. We first assume that ℓ is odd. Plugging the definition (3.1) of \mathcal{F}_j in (4.8) gives

$$H_{s+\frac{\ell}{2}} \left(\frac{it}{2\pi} \right) = - \sum_{j=0}^{\frac{\ell-1}{2}} \frac{D_{-2j-1} \left(\frac{it}{2\pi} \right)}{(2j)!} (2\ell)^j t^{-j} \mathcal{F}_{j, \frac{1}{2} - \frac{s}{\ell}}(2\ell t). \quad (5.8)$$

Now, by (5.7) and (3.4), we obtain that

$$D_{-2j-1} \left(\frac{it}{2\pi} \right) \mathcal{F}_{j, \frac{1}{2} - \frac{s}{\ell}}(2\ell t) t^{-j} \ll t^{-\ell+2j+1}.$$

Thus (if we show that this is not vanishing) the dominant term in (5.8) comes from $j = 0$ and gives, by (3.5),

$$H_{s+\frac{\ell}{2}} \left(\frac{it}{2\pi} \right) \sim -\frac{1}{2} D_{-1} \left(\frac{it}{2\pi} \right). \quad (5.9)$$

To determine the asymptotic behavior of D_{-1} , we compute, using (5.6) and (3.2),

$$\begin{aligned} D_{-1} \left(\frac{it}{2\pi} \right) &= 2\pi i \operatorname{coeff}_{z^{-1}} g_\ell \left(z; \frac{it}{2\pi} \right) \sim 2\pi i \left(-\frac{2\pi}{t} \right)^\ell \operatorname{coeff}_{z^{-1}} \frac{e^{\frac{2\pi^2 \ell z}{t}}}{\left(e^{\frac{4\pi^2 z}{t}} - 1 \right)^\ell} \\ &= 2\pi i \left(-\frac{2\pi}{t} \right)^\ell \frac{t}{4\pi^2} \operatorname{coeff}_{z^{-1}} \frac{e^{\frac{\ell z}{2}}}{(e^z - 1)^\ell} = \frac{-i \left(\frac{2\pi}{t} \right)^{\ell-1}}{(\ell-1)!} B_{\ell-1}^{(\ell)} \left(\frac{\ell}{2} \right). \end{aligned}$$

Thus when ℓ is odd, we have $\mathcal{C}_\ell = (-1)^{\frac{\ell-1}{2}} \frac{B_{\ell-1}^{(\ell)} \left(\frac{\ell}{2} \right)}{2^{(\ell-1)!}}$.

The theorem statement then follows from (5.2) and Proposition A.1. □

5.3. Proof of Theorem 1.1 for ℓ even. We next assume that ℓ is even. Plugging (3.6) in (4.8) gives

$$H_{s+\frac{\ell}{2}} \left(\frac{it}{2\pi} \right) = \sum_{1 \leq j \leq \frac{\ell}{2}} \frac{D_{-2j} \left(\frac{it}{2\pi} \right)}{(2j-1)!} \left(\frac{t}{2\ell} \right)^{\frac{1}{2}-j} \mathcal{G}_{j, \frac{1}{2} - \frac{s}{\ell}} \left(\frac{\ell t}{2} \right).$$

We have, by (3.7),

$$\mathcal{G}_{j, \frac{1}{2} - \frac{s}{\ell}} \left(\frac{\ell t}{2} \right) \sim \frac{(j-1)!}{\sqrt{2\ell t}}.$$

Now, by (5.7),

$$D_{-2j} \left(\frac{it}{2\pi} \right) \mathcal{G}_{j, \frac{1}{2} - \frac{s}{\ell}} \left(\frac{\ell t}{2} \right) t^{\frac{1}{2}-j} \ll t^{-\ell+j}.$$

Thus the dominant term comes from $j = 1$ and gives asymptotically (if non-vanishing)

$$\frac{1}{t} D_{-2} \left(\frac{it}{2\pi} \right). \quad (5.10)$$

Now, by (3.2) and (5.6), we obtain

$$\begin{aligned} D_{-2} \left(\frac{it}{2\pi} \right) &= (2\pi i)^2 \text{coeff}_{z^{-2}} g_\ell \left(z; \frac{it}{2\pi} \right) \sim (2\pi i)^2 \left(\frac{2\pi}{t} \right)^\ell \left(\frac{t}{4\pi^2} \right)^2 \text{coeff}_{z^{-2}} \frac{e^{\frac{\ell z}{2}}}{(e^z - 1)^\ell} \\ &= - \left(\frac{2\pi}{t} \right)^{\ell-2} \frac{B_{\ell-2}^{(\ell)} \left(\frac{\ell}{2} \right)}{(\ell-2)!}. \end{aligned}$$

Thus when ℓ is even, $\mathcal{C}_\ell = (-1)^{\frac{\ell}{2}+1} \frac{1}{2\pi} \frac{B_{\ell-2}^{(\ell)} \left(\frac{\ell}{2} \right)}{(\ell-2)!}$.

Thus the claim follows from (5.10) and Proposition A.1. \square

5.4. Proof of Corollary 1.2. The asymptotic formula (1.3) follows directly from Theorem 1.1 by plugging in (1.2), combined with (5.1).

For the second claim, we recall that

$$\text{qdim}(L_{\mathfrak{g}}(-\Lambda_0(s+1) + s\Lambda_1)) = \lim_{t \rightarrow 0^+} \frac{\text{ch}[L_{\mathfrak{g}}(-\Lambda_0(s+1) + s\Lambda_1)](it)}{\text{ch}[L_{\mathfrak{g}}(-\Lambda_0)](it)}$$

and observe that (1.3) is independent of s . \square

Remarks.

(1) In [1], the following fusion rules formula

$$V_{s_1} \boxtimes V_{s_2} = V_{s_1+s_2},$$

was established, where, for convenience, we let $V_s := L_{\mathfrak{g}}(-\Lambda_0(s+1) + s\Lambda_1)$, for $s \geq 0$, and $V_{-s} := L_{\mathfrak{g}}(-\Lambda_0(-s+1) - s\Lambda_{\ell-1})$ for $s < 0$. In this formula the symbol \boxtimes denotes the *fusion product*, that is, it indicates that the space of intertwining operators for this particular triple of modules is one-dimensional and zero otherwise. In other words the fusion ring generated by isomorphism classes of ordinary irreducible $L_{\mathfrak{g}}(-\Lambda_0)$ modules is isomorphic to $\mathbb{C}[\mathbb{Z}]$. Because of Corollary 1.2, the quantum dimension $\text{qdim}(\cdot)$ defines the trivial (one-dimensional) representation of the fusion ring.

(2) In [7], the first and the third author studied asymptotic properties of the Fourier coefficients of the Bloch-Okounkov n -point functions. They also discussed level¹ $\ell \in \mathbb{N}$ n -point functions $F^{(\ell)}$. As shown in [7, Section 4], the Bloch-Okounkov 1-point function of level ℓ is given by $F^{(\ell)}(z; \tau) = \frac{1}{\Theta(z; \tau)^\ell}$, where $\eta(\tau)^3 \Theta(z; \tau) = \vartheta(z; \tau)$. Theorem 1.1 can be used to derive the leading asymptotics of its Fourier coefficients.

¹Not to be confused with the level appearing in this paper.

After slight adjustments, due to additional Euler factors, we immediately get that the s -th Fourier coefficient $F_s^{(\ell)}$ satisfies (as $t \rightarrow 0^+$)

$$F_s^{(\ell)} \left(\frac{it}{2\pi} \right) \sim \mathcal{C}_\ell,$$

so it has no growing term.

6. A FULL ASYMPTOTIC EXPANSION FOR $\ell = 3$

In this section we specialize to $\ell = 3$ (i.e., $\mathfrak{g} = \mathfrak{sl}_3$) and explicitly work out the full asymptotic expansion in this case.

Proposition 6.1. *For $\Lambda \in \{\Lambda_1, \Lambda_2\}$ we have, as $t \rightarrow 0^+$,*

$$\begin{aligned} & \text{ch}[L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda)] \left(\frac{it}{2\pi} \right) \\ &= e^{-\frac{5\pi^2}{6t}} \left(\frac{2\pi}{t} \right)^{\frac{5}{2}} \left(\left(\frac{\pi^2}{4t^2} + \frac{3}{4t} \right) \sum_{n=0}^N \frac{E_{2n} \left(\frac{1}{2} - \frac{s}{3} \right)}{n!} \left(-\frac{3t}{2} \right)^n \right. \\ & \quad \left. + \frac{3}{8t} \sum_{n=0}^N \frac{E_{2n+2} \left(\frac{1}{2} - \frac{s}{3} \right)}{n!} \left(-\frac{3t}{2} \right)^n + O(t^{N+1}) \right). \end{aligned}$$

In particular, for $s = 0$ we have

$$\begin{aligned} & \text{ch}[L_{\mathfrak{g}}(-\Lambda_0)] \left(\frac{it}{2\pi} \right) \\ &= e^{-\frac{5\pi^2}{6t}} \left(\frac{2\pi}{t} \right)^{\frac{5}{2}} \left(\frac{\pi^2}{4t^2} + \frac{3}{4t} - \frac{1}{4t} \frac{\partial}{\partial t} \right) \left(\sum_{n=0}^N \frac{E_{2n}}{n!} \left(-\frac{3t}{8} \right)^n + O(t^{N+1}) \right). \end{aligned} \quad (6.1)$$

Proof: We have, by (4.9),

$$F_{3,s}(q) = i(q)_\infty^3 q^{-h_s - \frac{3}{8}} H_{s+\frac{3}{2}}(\tau). \quad (6.2)$$

Now, by (5.8)

$$H_{s+\frac{3}{2}} \left(\frac{it}{2\pi} \right) = -D_{-1} \left(\frac{it}{2\pi} \right) \mathcal{F}_{0, \frac{1}{2} - \frac{s}{3}}(6t) - D_{-3} \left(\frac{it}{2\pi} \right) \frac{3}{t} \mathcal{F}_{1, \frac{1}{2} - \frac{s}{3}}(6t). \quad (6.3)$$

We compute the Laurent coefficients D_{-j} , using (2.9) and (4.5),

$$D_{-3}(\tau) = i, \quad D_{-1}(\tau) = -\frac{3i}{8\pi^2} G_2(\tau).$$

Using

$$G_2 \left(\frac{it}{2\pi} \right) = -\frac{4\pi^4}{3t^2} + \frac{4\pi^2}{t} + O\left(t^{-2} e^{-\frac{4\pi^2}{t}}\right)$$

(3.3), and (6.3), we then obtain

$$\begin{aligned} H_{s+\frac{3}{2}}\left(\frac{it}{2\pi}\right) &= \left(-\frac{\pi^2 i}{4t^2} - \frac{3i}{4t}\right) \sum_{n=0}^N \frac{E_{2n}\left(\frac{1}{2} - \frac{s}{3}\right)}{n!} \left(-\frac{3t}{2}\right)^n \\ &\quad - \frac{3i}{8t} \sum_{n=0}^N \frac{E_{2n+2}\left(\frac{1}{2} - \frac{s}{3}\right)}{n!} \left(-\frac{3t}{2}\right)^n + O(t^{N+1}). \end{aligned} \quad (6.4)$$

Here we employ (2.5) with $m = 2$.

To finish the claim, we plug (6.2) into (1.2), to obtain that

$$\text{ch}[L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda_1)](\tau) = \frac{i}{\eta(\tau)^5} H_{s+\frac{3}{2}}(\tau).$$

Using (5.1) and (6.4), then yields the claims. Formula (6.1) follows easily by (2.4). \square

Proposition 6.1 immediately leads to the following asymptotic expansion for the quantum dimension.

Corollary 6.2. *For $\Lambda \in \{\Lambda_1, \Lambda_2\}$ we have, as $t \rightarrow 0^+$,*

$$\frac{\text{ch}[L_{\mathfrak{g}}(-(s+1)\Lambda_0 + s\Lambda)](it)}{\text{ch}[L_{\mathfrak{g}}(-\Lambda_0)](it)} \sim 1 - \frac{s^2(\pi^2 - 1)}{3\pi}t + O(t^2). \quad (6.5)$$

Remark. (Heisenberg patterns) Let $M(1)$ be the rank one Heisenberg vertex operator algebra. For an $M(1)$ -module $M(1, s)$, we easily get

$$\frac{\text{ch}[M(1, s)](it)}{\text{ch}[M(1)](it)} = e^{-2\pi s^2 t} = 1 - 2\pi s^2 t + O(t^2). \quad (6.6)$$

The reader should notice the similarity between (6.6) and Corollary 6.2 as they both have an s^2 in the second term of the asymptotic expansion (incidentally both vertex algebras share “additive” fusion rules [1, 2]). It would be very interesting to find a closed expression for the asymptotic expansion in (6.5).

7. A DECOMPOSITION FOR $F_{\ell, s}(\zeta_1, \dots, \zeta_{\ell-1})$

In this part we prove Theorem 1.3 from the introduction.

Proof of Theorem 1.3: We write

$$F_{\ell, s}(\zeta_1, \dots, \zeta_{\ell-1}) = (-i)^\ell q^{\frac{\ell}{8}} \prod_{j=1}^{\ell-1} \zeta_j^{-\frac{j}{2}} (q)_\infty^{\ell+1} \mathcal{F}_{\ell, s+\frac{\ell}{2}}. \quad (7.1)$$

Here, for $r \in \mathbb{Z} + \frac{\ell}{2}$,

$$\mathcal{F}_{\ell, r} := \text{coeff}_{\zeta_i^r} \mathcal{F}_\ell(z_\ell),$$

where

$$\mathcal{F}_\ell(z_\ell) = \mathcal{F}_\ell(z_1, \dots, z_{\ell-1}, z_\ell) := \frac{(-1)^\ell}{\prod_{j=1}^{\ell} \vartheta(w_j)}.$$

Using (2.7), we have

$$\mathcal{F}_\ell(z_\ell + 1) = (-1)^\ell \mathcal{F}_\ell(z_\ell), \quad \mathcal{F}_\ell(z_\ell - \tau) = (-1)^\ell q^{\frac{\ell}{2}} \mathcal{F}_\ell(z_\ell) \prod_{j=1}^{\ell} \zeta_j^{-j}.$$

We now find a formula for $\mathcal{F}_{\ell,r}$ that is closely related to Theorem 4.1 above (which was proven in [8] by a similar argument). In particular, we compute the following integral in two ways:

$$\int_{\partial P_{z_0}} \mathcal{F}_\ell(w) \vartheta_{r,\varepsilon,\frac{\ell}{2}}^+ \left(w + \frac{1}{\ell} \sum_{j=1}^{\ell-1} j z_j \right) dw. \quad (7.2)$$

Here z_0 is chosen such that $w_j \in P_{z_0}$ for all j (which is possible due to the restrictions on the ζ_j); note that $-v < \text{Im}(z_0) < 0$.

For the first computation, we also use the elliptic shifts (as above, $\varepsilon \equiv \ell \pmod{2}$)

$$\begin{aligned} (-1)^\ell q^{\frac{\ell}{2}} e^{-2\pi i \ell z} \prod_{j=1}^{\ell-1} \zeta_j^{-j} \vartheta_{r,\varepsilon,\frac{\ell}{2}}^+ \left(z + \frac{1}{\ell} \sum_{j=1}^{\ell-1} j z_j - \tau \right) &= \vartheta_{r,\varepsilon,\frac{\ell}{2}}^+ \left(z + \frac{1}{\ell} \sum_{j=1}^{\ell-1} j z_j \right) \\ &= (-1)^\ell q^{\frac{(r+\ell)^2}{2\ell}} e^{-2\pi i (r+\ell)z} \prod_{j=1}^{\ell-1} \zeta_j^{-\frac{j(r+\ell)}{\ell}}, \\ \vartheta_{r,\varepsilon,\frac{\ell}{2}}^+ \left(z + \frac{1}{\ell} \sum_{j=1}^{\ell-1} j z_j \right) &= (-1)^\ell \vartheta_{r,\varepsilon,\frac{\ell}{2}}^+ \left(z + \frac{1}{\ell} \sum_{j=1}^{\ell-1} j z_j + 1 \right), \end{aligned}$$

to obtain that (7.2) equals

$$(-1)^\ell q^{\frac{(r+\ell)^2}{2\ell}} \mathcal{F}_{\ell,r+\ell} \prod_{j=1}^{\ell-1} \zeta_j^{-\frac{(r+\ell)j}{\ell}}. \quad (7.3)$$

On the other hand, the Residue Theorem implies that (7.2) also equals

$$2\pi i \sum_{w \in S_{z_0}(\tau)} \text{Res}_{z=w} \mathcal{F}_\ell(z) \vartheta_{r,\varepsilon,\frac{\ell}{2}}^+ \left(z + \frac{1}{\ell} \sum_{j=1}^{\ell-1} j z_j \right).$$

Now, for $1 \leq j \leq \ell$, \mathcal{F}_ℓ has simple poles at w_j . We compute

$$\text{Res}_{w=w_\nu} \mathcal{F}_\ell(w) = -\frac{1}{2\pi\eta^3} \frac{1}{\prod_{\substack{j=1 \\ j \neq \nu}}^{\ell} \vartheta(w_\nu - w_j)}.$$

Comparing to (7.3), we find that

$$\mathcal{F}_{\ell,r+\ell} = -i(-1)^\ell q^{-\frac{(r+\ell)^2}{2\ell}} \prod_{j=1}^{\ell-1} \zeta_j^{\frac{(r+\ell)j}{\ell}} \eta(\tau)^{-3} \sum_{\nu=1}^{\ell} \frac{\vartheta_{r,\varepsilon,\frac{\ell}{2}}^+ \left(w_\nu - \frac{1}{\ell} \sum_{j=1}^{\ell} w_j \right)}{\prod_{\substack{j=1 \\ j \neq \nu}}^{\ell} \vartheta(w_\nu - w_j)}.$$

Plugging in to (7.1) and setting $r = s - \frac{\ell}{2}$ gives the claim. \square

Remarks.

(1) One could prove related decompositions without the condition that the w_j are distinct. Any repeated values among the w_j require modified calculations of residues in the proof above, leading to partial derivatives of ϑ^+ (as in Theorem 4.2, which is related to the limiting case that all w_j have the repeated value 0).

(2) Theorem 1.3 could be used to determine the asymptotic behavior as $t \rightarrow 0^+$ of $F_{\ell,s}(\zeta_1, \dots, \zeta_{\ell-1})$, so long as the ζ_j are also bounded in the ranges given in the theorem.

(3) We stress that Theorem 1.3 also applies to $\ell = 2$, which is quite different compared to $\ell \geq 3$ because the relevant vertex algebra is not an affine Lie algebra). Furthermore, note that in the case $\ell = 1$, Theorem 1.3 is equivalent to Theorem 4.2.

8. MODULAR-TYPE TRANSFORMATION PROPERTIES OF $F_{\ell,s}$

In this section we discuss general modular transformation properties of $F_{\ell,s}$. The modular transformation properties of the denominator appearing in the formula for $F_{\ell,s}$ are well-understood, so we only have to consider the numerator. We note that in the parlance of [9, 11], $\vartheta_{s+\frac{\ell}{2}, \ell, \frac{\ell}{2}}^+(z; \tau)$ is an example of a regularized (Jacobi) false theta function. The following proposition gives the modular transformation properties of these functions.

Proposition 8.1. *For $z \in \mathbb{C} \setminus \mathbb{R}$ and $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ with $c \neq 0$, we have*

$$\begin{aligned} & \vartheta_{r,\varepsilon,M}^+ \left(z; \frac{a\tau + b}{c\tau + d} \right) \\ &= \sqrt{-\frac{i(c\tau + d)}{2}} \sum_{j=0}^{2c-1} (-1)^{j\varepsilon} \zeta_{4cM}^{a(2Mj-r)^2} \left(\zeta^{-r+2Mj} \int_{\mathbb{R}} \frac{e^{\frac{\pi i(c\tau+d)x^2}{2} - \frac{\pi i}{\sqrt{cM}}(r-2Mj)x}}{1 - \zeta^{4cM} e^{4\pi i \sqrt{cM}x}} dx \right. \\ & \quad + \frac{1}{4\sqrt{cM}} (1 - \mathrm{sgn}(\mathrm{Im}(z))) \zeta_{8cM}^{2Mj-r} e^{2\pi i cM(c\tau+d)(z - \frac{1}{8cM})^2} \\ & \quad \left. \times \vartheta \left(\frac{c\tau + d}{2} \left(z - \frac{1}{8cM} \right) - \frac{1}{2} + \frac{r - 2Mj}{4cM}; \frac{c\tau + d}{8cM} \right) \right). \end{aligned}$$

Proof: We assume without loss of generality that $c > 0$. Plugging in the series representation of ϑ^+ and writing $\frac{a\tau+b}{c\tau+d} = \frac{a}{c} - \frac{1}{c(c\tau+d)}$, we find that

$$\vartheta_{r,\varepsilon,M}^+ \left(z; \frac{a\tau + b}{c\tau + d} \right) = \sum_{j=0}^{2c-1} (-1)^{j\varepsilon} \zeta_{4cM}^{a(2Mj-r)^2} \vartheta_{r-2Mj,0,2cM}^+ \left(z; -\frac{2}{c\tau + d} \right).$$

We next invert each term in the sum by finding an inversion formula for $\vartheta_{r,0,M}^+(z; \frac{i}{w})$. If $\text{Im}(z) > 0$, then we use

$$e^{-\frac{\pi i}{\tau} A^2} = \sqrt{-i\tau} \int_{\mathbb{R}} e^{\pi i \tau x^2 + 2\pi i A x} dx,$$

to obtain that

$$\vartheta_{r,0,M}^+ \left(z; \frac{i}{w} \right) = \sqrt{w} \zeta^{-r} \int_{\mathbb{R}} \frac{e^{-\pi w x^2 - \frac{2\pi i r}{\sqrt{2M}} x}}{1 - \zeta^{2M} e^{2\pi i \sqrt{2M} x}} dx. \quad (8.1)$$

Plugging into (8.1) gives the claim in this case.

If $\text{Im}(z) < 0$, then we complete the theta function, calculating

$$\vartheta_{r,0,M}^+(z; \tau) + \vartheta_{-r-2M,0,M}^+(-z; \tau) = q^{\frac{(M-r)^2}{4M}} \zeta^{M-r} \vartheta \left(2Mz - \frac{1}{2} + (M-r)\tau; 2M\tau \right).$$

For the partial theta function $\vartheta_{-r-2M,0,M}^+(-z; \tau)$ we again use (8.1). which is invariant under $r \mapsto -r - 2M$, $z \mapsto -z$, and multiplication by -1 . From (2.8) we obtain for the theta function

$$\begin{aligned} e^{-\frac{\pi(M-r)^2}{2Mw}} \zeta^{M-r} \vartheta \left(2Mz - \frac{1}{2} + (M-r)\frac{i}{w}; -\frac{2M}{iw} \right) \\ = -i \zeta_{4M}^{M-r} e^{-2\pi M w (z - \frac{1}{4M})^2} \sqrt{\frac{w}{2M}} \vartheta \left(izw - \frac{iw}{4M} - \frac{1}{2} + \frac{r}{2M}; \frac{iw}{2M} \right). \end{aligned}$$

Plugging in gives the claim. \square

As a special case, we recover Proposition 7 of [11].

Corollary 8.2. *For $z \in \mathbb{C} \setminus \mathbb{R}$, we have*

(i) *If ℓ is odd, then we have*

$$\begin{aligned} (-i\tau)^{-\frac{1}{2}} \vartheta_{s+\frac{\ell}{2}, 1, \frac{\ell}{2}}^+ \left(z; -\frac{1}{\tau} \right) \\ = \frac{1}{2} \int_{\mathbb{R}} \frac{e^{\pi i \tau x^2} e^{\frac{2\pi i}{\sqrt{\ell}}(s+\ell)(x-\sqrt{\ell}z)}}{\cos(\sqrt{\ell}\pi(x-\sqrt{\ell}z))} dx + \frac{(1 - \text{sgn}(\text{Im}(z)))}{2\sqrt{\ell}} e^{\pi i \ell \tau z^2} \vartheta \left(\tau z + \frac{s}{\ell}; \frac{\tau}{\ell} \right). \end{aligned}$$

(ii) *If ℓ is even, then we have*

$$\begin{aligned} (-i\tau)^{-\frac{1}{2}} \vartheta_{s+\frac{\ell}{2}, 0, \frac{\ell}{2}}^+ \left(z; -\frac{1}{\tau} \right) = -\frac{i}{2} \int_{\mathbb{R}} \frac{e^{\pi i \tau x^2} e^{\frac{2\pi i}{\sqrt{\ell}}(s+\ell)(x-\sqrt{\ell}z)}}{\sin(\sqrt{\ell}\pi(x-\sqrt{\ell}z))} dx \\ + \frac{(1 - \text{sgn}(\text{Im}(z)))}{2i\sqrt{\ell}} e^{-\frac{\pi i s}{\ell}} e^{\pi i \ell \tau (z - \frac{1}{2\ell})^2} \vartheta \left(\tau z + \frac{s}{\ell} - \frac{\tau}{2\ell}; \frac{\tau}{\ell} \right). \end{aligned}$$

Proof: The claim follows from Proposition 8.1 with $a = d = 0$, $c = -b = 1$, using the easily verified identity

$$\frac{1}{2}e^{-\frac{\pi iz}{2} - \frac{\pi i}{4} + \frac{\pi i \tau}{16}} \left(\vartheta \left(\frac{z}{2} - \frac{\tau}{8} - \frac{1}{4}; \frac{\tau}{4} \right) + i\vartheta \left(\frac{z}{2} - \frac{\tau}{8} + \frac{1}{4}; \frac{\tau}{4} \right) \right) = \vartheta(z; \tau).$$

□

Remark. In view of Theorem 1.3, the modular transformation formulas in this section can be now used to compute modular transformation formulas for $\text{ch}[M](z; \tau)$ for every irreducible module M . This result then can be interpreted as a statement on characters of more general class of irreducible $L_{\mathfrak{g}}(-\Lambda_0)$ -modules parametrized by a continuous parameter. Presumably, such results can be used to formulate a Verlinde type formula for $L_{\mathfrak{g}}(-\Lambda_0)$ -modules. As this analysis is clearly beyond the scope of this paper, it is left for future investigation.

APPENDIX A. COEFFICIENTS OF BERNOULLI POLYNOMIALS

Here we calculate some special values for certain generalized Bernoulli polynomials. In particular, we are interested in the values

$$\mathcal{C}_\ell^* := \begin{cases} (-1)^{\frac{\ell-1}{2}} \frac{B_{\ell-1}^{(\ell)}(\frac{\ell}{2})}{2(\ell-1)!} & \text{if } \ell \text{ is odd,} \\ (-1)^{\frac{\ell}{2}+1} \frac{1}{2\pi} \frac{B_{\ell-2}^{(\ell)}(\frac{\ell}{2})}{(\ell-2)!} & \text{if } \ell \text{ is even.} \end{cases}$$

Proposition A.1. *We have*

$$\mathcal{C}_\ell = \mathcal{C}_\ell^*.$$

Proof: We offer 2 proofs. The first proof uses recurrences. We first assume that ℓ is odd. It is direct that

$$\mathcal{C}_1 = \mathcal{C}_1^* = \frac{1}{2}, \quad \mathcal{C}_{\ell+2} = \frac{\ell}{4(\ell+1)} \mathcal{C}_\ell. \quad (\text{A.1})$$

Thus the identity follows, once we show that \mathcal{C}_ℓ^* also satisfies (A.1) (with \mathcal{C}_ℓ replaced by \mathcal{C}_ℓ^*). For this we use the following integral representation of \mathcal{C}_ℓ^*

$$B_{\ell-1}^{(\ell)} \left(\frac{\ell}{2} \right) = -2^{2-\ell} (\ell-1)! \frac{i^{-\ell}}{2\pi i} \int_{\mathbb{R}} \frac{1}{\cosh(z)^\ell} dz$$

which follows by a direct residue calculation. The recurrence (A.1) now follows, using

$$\frac{d^2}{dz^2} \frac{1}{\cosh(z)^\ell} = \frac{\ell^2}{\cosh(z)^\ell} - \frac{\ell(\ell+1)}{\cosh(z)^{\ell+2}}. \quad (\text{A.2})$$

We next assume that ℓ is even. Then

$$\mathcal{C}_2 = \mathcal{C}_2^* = \frac{1}{2\pi}, \quad \mathcal{C}_{\ell+2} = \frac{\ell}{4(\ell+1)} \mathcal{C}_\ell. \quad (\text{A.3})$$

In this case, we have the representation

$$B_{\ell-2}^{(\ell)}\left(\frac{\ell}{2}\right) = -2^{1-\ell}(\ell-2)!i^{-\ell} \int_{\mathbb{R}} \frac{1}{\cosh(z)^\ell} dz.$$

As above, plugging in (A.2) now shows that (A.3) holds for \mathcal{C}_ℓ^* .

Second proof. The second proof uses the formal Residue Theorem. Writing $w = g(z) \in z\mathbb{C}[[z]]$ with $g'(0) \neq 0$, we have

$$\operatorname{Res}_{w=0} f(w) = \operatorname{Res}_{z=0} f(g(z))g'(z).$$

For ℓ odd, we let $f(w) := \frac{(w+1)^{\frac{\ell}{2}-1}}{w^\ell}$ and $w := g(z) = e^z - 1$. Then

$$\operatorname{Res}_{z=0} \frac{e^{\frac{\ell}{2}z}}{(e^z - 1)^\ell} = \operatorname{Res}_{w=0} \frac{(w+1)^{\frac{\ell}{2}-1}}{w^\ell} = \binom{\frac{\ell}{2}-1}{\ell-1}.$$

This implies the statement.

For ℓ even, we have

$$I_\ell := \operatorname{Res}_{w=0} \frac{\log(1+w)(w+1)^{\frac{\ell}{2}-1}}{w^\ell} = \operatorname{Res}_{z=0} \frac{ze^{\frac{\ell}{2}z}}{(e^z - 1)^\ell}.$$

Expanding $\log(1+w) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{w^n}{n}$, we get

$$I_\ell = \operatorname{Res}_{w=0} \frac{(w+1)^{\frac{\ell}{2}-1}}{w^\ell} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{w^n}{n} = \sum_{j=0}^{\frac{\ell}{2}-1} \binom{\frac{\ell}{2}-1}{j} \frac{(-1)^{\frac{\ell}{2}+1+j}}{\frac{\ell}{2}+j}.$$

To this end, we need the evaluation

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{1}{k+c} = \frac{n!(c-1)!}{(n+c)!}. \quad (\text{A.4})$$

To see (A.4), we integrate the Binomial Theorem to obtain (here $B(a, b)$ is the usual Beta function)

$$\begin{aligned} \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{k+c} &= \int_0^1 \sum_{k=0}^n \binom{n}{k} (-1)^k x^{k+c-1} dx = \int_0^1 x^{c-1} (1-x)^n dx = B(c, n+1) \\ &= \frac{\Gamma(c)\Gamma(n+1)}{\Gamma(c+n+1)}, \end{aligned}$$

by the standard evaluation of the Beta integral.

Letting $n = \frac{\ell}{2} - 1$ and $c = \frac{\ell}{2}$ in (A.4), we then obtain $I_\ell = (-1)^{\frac{\ell}{2}+1} \frac{(\frac{\ell}{2}-1)!(\frac{\ell}{2}-1)!}{(\ell-1)!}$ as desired. \square

REFERENCES

- [1] D. Adamovic and O. Perse, *Fusion rules and complete reducibility of certain modules for affine Lie algebras*, Journal of Algebra and its Applications **13** (2014), 1350062.
- [2] D. Adamovic and O. Perse, *Representations of certain non-rational vertex operator algebras of affine type*, Journal of Algebra **319** (2008), 2434-2450.
- [3] T. Arakawa and K. Kawasetsu, *Quasi-lisse vertex operator algebras and modular linear differential equations*, arXiv:1610.05865.
- [4] T. Arakawa and A. Moreau, *Sheets and associated varieties of affine vertex algebras*, arXiv:1601.05906, to appear in Advances in Mathematics.
- [5] C. Beem, M. Lemos, P. Liendo, W. Peelaers, L. Rastelli, and B. C. van Rees, *Infinite Chiral Symmetry in Four Dimensions*, arXiv:1312.5344.
- [6] K. Bringmann, T. Creutzig, and L. Rolin, *Negative index Jacobi forms and quantum modular forms*, Res. Math. Sci. **1** (2014), 1-32.
- [7] K. Bringmann and A. Milas, *On the Fourier expansion of Bloch-Okounkov n -point function*, Journal of Combinatorial Theory Series A **36** (2015), 201-219.
- [8] K. Bringmann, L. Rolin, and S. Zwegers, *On the Fourier Coefficients of negative index meromorphic Jacobi forms*, Research in the Mathematical Sciences **3** (2016), 1-9.
- [9] T. Creutzig and A. Milas, *False theta functions and the Verlinde formula*, Adv. Math. **262** (2014), 520-545.
- [10] T. Creutzig and A. Milas, *Higher rank partial and false theta functions and representation theory*, Advances in Mathematics **314** (2017), 203-227.
- [11] T. Creutzig, A. Milas, and S. Wood, *On regularised quantum dimensions of the singlet vertex operator algebra and false theta functions*, International Mathematics Research Notices (**2017.5**) 1390-1432.
- [12] I. Frenkel and Y. Zhu, *Vertex operator algebras associated to representations of affine and Virasoro algebras*, Duke Math. J **66** (1992), 123-168.
- [13] Y.Z. Huang, *Vertex operator algebras and the Verlinde conjecture*, Communications in Contemporary Mathematics **10**, 01 (2008): 103-154.
- [14] V. Kac and M. Wakimoto, *Modular invariant representations of infinite-dimensional Lie algebras and superalgebras*, Proceedings of the National Academy of Sciences **85** (1988), 4956-4960.
- [15] V. Kac and M. Wakimoto, *Integrable highest weight modules over affine superalgebras and Appell's function*, Communications in Mathematical Physics **215** (2001): 631-682.
- [16] V. Kac and M. Wakimoto, *On characters of irreducible highest weight modules of negative integer level over affine Lie algebras*, arXiv:1706.08387.
- [17] N.E. Norlund, *Vorlesungen über Differenzenrechnung*, Berlin, 1924.
- [18] D. Zagier, *The Mellin transform and other useful analytic techniques*, Appendix to E. Zeidler, Quantum Field Theory I: Basics in Mathematics and Physics. A Bridge Between Mathematicians and Physicists Springer-Verlag, Berlin-Heidelberg-New York (2006), 305-323.

MATHEMATICAL INSTITUTE, UNIVERSITY OF COLOGNE, WEYERTAL 86-90, 50931 COLOGNE, GERMANY

E-mail address: kbringma@math.uni-koeln.de

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA

E-mail address: mahlburg@math.lsu.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, SUNY-ALBANY, ALBANY, NY 12222, U.S.A.

E-mail address: `amilas@albany.edu`