

PARTIAL THETA FUNCTIONS AND MOCK MODULAR FORMS AS q -HYPERGEOMETRIC SERIES

KATHRIN BRINGMANN, AMANDA FOLSOM, AND ROBERT C. RHOADES

Dedicated to the visionary Ramanujan, on the 125th anniversary of his birth.

ABSTRACT. Ramanujan studied the analytic properties of many q -hypergeometric series. Of those, mock theta functions have been particularly intriguing, and by work of Zwegers, we now know how these curious q -series fit into the theory of automorphic forms. The analytic theory of partial theta functions however, which have q -expansions resembling modular theta functions, is not well understood. Here we consider families of q -hypergeometric series which converge in two disjoint domains. In one domain, we show that these series are often equal to one another, and define mock theta functions, including the classical mock theta functions of Ramanujan, as well as certain combinatorial generating functions, as special cases. In the other domain, we prove that these series are typically not equal to one another, but instead are related by partial theta functions.

1. INTRODUCTION

1.1. **Curious q -series.** Partial theta functions have played a curious role in the theory of partitions (see for instance [1, 2, 10]). For example, Rogers proved the following identity for $|q| < 1$:

$$(1.1) \quad \sum_{n \geq 0} \frac{(-1)^n q^{\frac{n(n+1)}{2}}}{(-q)_n} = \sum_{n \geq 0} \binom{-12}{n} q^{\frac{n^2-1}{24}} =: \psi(q),$$

where $(a)_n = (a; q)_n := \prod_{j=0}^{n-1} (1 - aq^j)$, and (\cdot) denotes the usual Kronecker symbol. Andrews [4] provided a partition theoretic interpretation, giving a relationship between those partitions into distinct parts and those with odd largest part and even largest part. The series defined by $\psi(q)$ in (1.1) is an example of a partial theta function, meaning that while the series expansion resembles that of a theta function, the sum runs only over half of a lattice instead of a full one. Despite a closeness in the q -series expansions of partial theta functions and usual theta functions, which are modular forms, the analytic theory of partial theta functions is not well understood.

Rogers's identity in (1.1) is by no means an isolated example of this type. There are many instances in the literature of q -hypergeometric series that are combinatorial generating functions, partial theta functions, mock theta functions, or modular forms. For example, with $|q| < 1$, we have the combinatorial q -hypergeometric series [3]

$$(1.2) \quad R(w; q) := 1 + \sum_{n \geq 1} \frac{q^{n^2}}{(wq)_n (w^{-1}q)_n} = 1 + \sum_{n \geq 1} \sum_{m \in \mathbb{Z}} R(m, n) w^m q^n,$$

where $R(m, n)$ denotes the number of integer partitions of n with rank m , and Dyson's *rank* [12] of a partition equals the largest part minus its number of parts. A well known theorem, using Durfee squares, gives that $R(1; q) = q^{1/24}/\eta(\tau)$ ($q = e^{2\pi i\tau}$, $\tau \in \mathbb{H}$) is a modular form of weight $-1/2$. Here

2010 *Mathematics Subject Classification.* 33D15, 11F37, 11F27.

Key words and phrases. Ramanujan; modular form; partial theta function; mock theta function; q -hypergeometric series.

$\eta(\tau) := q^{\frac{1}{24}}(q)_\infty$ is Dedekind's eta-function. Moreover, setting $w = -1$ in (1.2), one obtains $f(q)$, Ramanujan's third order mock theta function which he defined in his last letter to Hardy as

$$(1.3) \quad f(q) := 1 + \sum_{n \geq 1} \frac{q^{n^2}}{(-q)_n^2},$$

which, in analogy to Rogers's partial theta function identity (1.1), is a q -hypergeometric series admitting a combinatorial interpretation of f as the generating function for the difference between the number of integer partitions with an even number of parts and those with an odd number of parts. Thanks to the breakthrough of the 2002 doctoral thesis of Zwegers [28], we now finally understand how the mock theta functions fit into the theory of modular forms. Loosely speaking, Zwegers showed that the mock theta functions satisfy modular transformation laws after they are "completed" by the addition of certain non-holomorphic integrals. Zwegers's discovery led to the development of a more overarching theory of *harmonic weak Maass forms*, which are certain nonholomorphic modular forms. Many advances have been made in the last decade in this area, notably by the first author and Ono, and their collaborators. (See for instance, the surveys of Ono [23] and Zagier [27] and the many references therein.) On the other hand, while there are tantalizing hints which suggest a connection between certain partial theta functions and Ramanujan's mock theta functions, the analytic theory of the partial theta functions is not well understood. See, for instance, the works of Hikami, Lawrence-Zagier, and Zagier [17, 19, 26] on topological invariants of 3-manifolds. In this paper, we seek to understand further curiosities evoked by partial theta functions and mock theta functions. In particular, we study relationships between these functions inside *and* outside the unit disc $\mathbb{D} := \{q \in \mathbb{C} : |q| < 1\}$. We first illustrate this with a motivating example.

1.2. An example. Consider the following q -hypergeometric series, defined for $|q| < 1$ by

$$(1.4) \quad f^*(q) := 1 + \sum_{n \geq 1} \frac{(-1)^{n+1} q^n}{(-q)_n}.$$

It is known (see (26.22) of [13] for example) that inside the unit disc $f^*(q)$ equals Ramanujan's mock theta function $f(q)$, defined in (1.3). On the other hand, $f^*(q)$ also converges outside of the unit disc. For $|q| < 1$ we have, using (3.6) below (or equation (4.16) of [17]), as well as (4.20) of [17]

$$(1.5) \quad f^*(q^{-1}) = 1 - \sum_{n \geq 1} (-1)^n \frac{q^{\frac{n(n-1)}{2}}}{(-q)_n} = 2\psi(q).$$

Note that the function $f^*(q)$ defined in [17] is not equal to the function $f^*(q)$ defined in (1.4). Continuing with this example, the q -hypergeometric series $f(q)$ from (1.3) also converges outside of the unit disc. In light of (1.5), one might expect $f(q^{-1})$ to equal $2\psi(q)$ (for $|q| < 1$), but this is not the case. The difference between $f(q^{-1})$ and the expected function $2\psi(q)$ is the series

$$(1.6) \quad S(q) := \frac{1}{(-q)_\infty^2} \sum_{n \geq 0} (-1)^n q^{\frac{n(n+1)}{2}},$$

that is the quotient of a partial theta function and a modular form. This fact is due to an identity found in Ramanujan's "lost" notebook (given in [5], valid for $|q| < 1$):

$$f(q^{-1}) = 2\psi(q) - S(q).$$

We note that this identity may also be derived from (2.6) using (3.7).

Consider now the series

$$(1.7) \quad \phi(q) := \sum_{n \geq 0} \frac{q^{n^2}}{(-q^2; q^2)_n}.$$

Again this series converges inside and outside the unit disc. In the region $|q| < 1$ it is one of Ramanujan's original (third order) mock theta functions. In his "lost" notebook (see [8] p. 235), Ramanujan showed that for $|q| < 1$

$$\phi(-q^{-1}) = \psi(q).$$

Hence, by (1.5), as before we might expect that the two third order mock theta functions satisfy $\phi(q) = \frac{1}{2}f(-q)$. However, this is again not the case. Interestingly, we know that the mock theta functions $\phi(q)$ and $f(-q)$ are related via the Mock Theta Conjectures (see for instance (3.12) of [14]). Namely

$$2\phi(q) = f(-q) + T(q),$$

where

$$(1.8) \quad T(q) := \frac{(q^2; q^2)_\infty^7}{(q)_\infty^3 (q^4; q^4)_\infty^3}.$$

The function $T(q)$ is, up to a power of q , a modular form. We summarize the above discussion with the following diagram ($|q| < 1$):

$$(1.9) \quad \begin{array}{ccccc} f(q) & = & f^*(q) & = & 2\phi(-q) - T(-q) & \text{mock theta} \\ \downarrow & & \downarrow & & \downarrow & \\ f(q^{-1}) + S(q) & = & f^*(q^{-1}) & = & 2\phi(-q^{-1}) & \text{partial theta} \end{array}$$

That is, we have three different q -hypergeometric series $f(q)$, $f^*(q)$, and $\phi(q)$ (that are mock theta functions) which define the same function inside \mathbb{D} , but different functions outside \mathbb{D} . The functions outside \mathbb{D} include the same partial theta function $\psi(q)$, and their difference $S(q)$ is defined by a partial theta function and a modular form.

Remark. Partial theta functions arise naturally in many q -series identities, and are often written in a slightly different form from that of $\psi(q)$ in (1.1). For example, it is not difficult to see that

$$\psi(q) = \sum_{n \geq 0} q^{\frac{n(3n+1)}{2}} (1 - q^{2n+1}) = \sum_{n \geq 0} q^{\frac{n(3n+1)}{2}} - \sum_{n \leq -1} q^{\frac{n(3n+1)}{2}}.$$

From this perspective, such series are often called *false theta functions* (see [4]).

1.3. q -hypergeometric families and mock theta functions. In this paper, we study families of q -hypergeometric series inside and outside \mathbb{D} , that include the classical mock theta functions of Ramanujan, as well as certain combinatorial generating functions, as special cases. We show that in more generality, curiosities as depicted by (1.9) also hold for the aforementioned families of functions. Throughout we denote

$$(a_1, \dots, a_r)_n = (a_1, \dots, a_r; q)_n := (a_1)_n \cdot \dots \cdot (a_r)_n.$$

When $|q| < 1$, we consider the following families of q -hypergeometric series:

$$(1.10) \quad g_2(w; q) := \sum_{n \geq 0} \frac{(-q)_n q^{\frac{n(n+1)}{2}}}{(w, w^{-1}q)_{n+1}},$$

$$(1.11) \quad g_3(w; q) := \sum_{n \geq 0} \frac{q^{n(n+1)}}{(w, w^{-1}q)_{n+1}},$$

$$(1.12) \quad K(w; q) := \sum_{n \geq 0} \frac{(-1)^n q^{n^2} (q; q^2)_n}{(wq^2, w^{-1}q^2; q^2)_n},$$

which define functions (in certain ranges of w which we will specify later) for both $|q| < 1$ and $|q| > 1$. Hickerson, and later Gordon and McIntosh [14, 15, 16], noticed that all of Ramanujan’s original mock theta functions can be written in terms of $g_2(w; q)$ and $g_3(w; q)$ for $|q| < 1$. Hence, these functions are called *universal mock theta functions*. The function $K(w; q)$ appears in Ramanujan’s “lost” notebook [5], and is related by modular transformation to the universal mock theta function (1.10) [21]. The families of q -hypergeometric series discussed here are essentially “mock Jacobi forms”, which first occurred in Zwegers’s thesis [28] and were systematically studied by the first author and Richter [11]. For the purpose of this paper, we will not require this perspective.

The remainder of the paper is structured as follows. In §2 we provide background and useful tools and results from q -combinatorial analysis. In §3 - §5 we introduce and study many infinite families of q -hypergeometric series inside and outside \mathbb{D} in analogy to the curiosities observed in (1.9).

2. q -COMBINATORIAL ANALYSIS

In this section, we provide some combinatorial identities which we require for the proofs of the main theorems. Many of the series studied in this paper are related to the *basic hypergeometric series*, defined by

$$F(a, b; t) = F(a, b; t, q) := \sum_{n \geq 0} \frac{(aq)_n t^n}{(bq)_n}.$$

We refer the reader to [13] for a detailed discussion on these functions, including questions of convergence. These series satisfy various difference equations, including the following (see (2.4) of [13])

$$(2.1) \quad F(a, b; t) = \frac{1-b}{1-t} + \frac{b-atq}{1-t} F(a, b; tq),$$

which lead to many interesting identities. One identity, the “Rogers-Fine identity,” that we will make use of here is as follows (see equation (1) of [24]):

$$(2.2) \quad F(a/q, b/q; t) = \sum_{n \geq 0} \frac{(a, atq/b)_n b^n t^n q^{n^2-n} (1-atq^{2n})}{(b)_n (t)_{n+1}}.$$

Iteration methods also give rise to the identities (see (12.2) and (12.3) of [13])

$$(2.3) \quad (1-t)F(a, b; t) = \sum_{n \geq 0} \frac{\left(\frac{b}{a}\right)_n}{(bq)_n (tq)_n} (-at)^n q^{\frac{n(n+1)}{2}},$$

$$(2.4) \quad (1-t)F(0, b; t) = \sum_{n \geq 0} \frac{(bt)^n q^{n^2}}{(bq)_n (tq)_n},$$

where (2.4) is derived from (2.3) by letting $a \rightarrow 0$. In special cases, these q -hypergeometric series may directly reduce to simpler expressions, such as (see (14.31) of [13])

$$(2.5) \quad (1-a)F(a, -a; a) = 1 + 2 \sum_{n \geq 1} (-1)^n a^{2n} q^{n^2}.$$

On the other hand, the following more involved result of Andrews (see Theorem 1 of [7]), when combined with the Rogers-Fine identity, is very useful in establishing many partial theta function identities

$$(2.6) \quad \sum_{n \geq 0} \frac{(B, -Abq)_n}{(-aq, -bq)_n} q^n = -a^{-1} \frac{(B, -Abq)_\infty}{(-aq, -bq)_\infty} \sum_{m \geq 0} \frac{(A^{-1})_m}{(-Ba^{-1})_{m+1}} (Aba^{-1}q)^m \\ + (1+a^{-1})(1+b) \sum_{m \geq 0} \frac{(-a^{-1}q, -a^{-1}ABq)_m}{(-Ba^{-1}, Aba^{-1}q)_{m+1}} (-b)^m.$$

Similarly, we have the following results of Ramanujan (see (1.1)_R of [7], (3.1) of [9], and (3.11)_R of [7]):

$$(2.7) \quad \sum_{n \geq 0} \frac{q^n}{(-aq, -a^{-1}q)_n} = (1+a) \sum_{n \geq 0} a^{3n} q^{\frac{n(3n+1)}{2}} (1-a^2 q^{2n+1}) - \frac{a}{(-aq, -a^{-1}q)_\infty} \sum_{n \geq 0} (-1)^n a^{2n} q^{\frac{n(n+1)}{2}},$$

$$(2.8) \quad \sum_{n \geq 0} \frac{(-1)^n a^{2n} q^{\frac{n(n+1)}{2}}}{(-aq)_n} = \sum_{n \geq 0} a^{3n} q^{\frac{n(3n+1)}{2}} (1-a^2 q^{2n+1}),$$

$$(2.9) \quad (1+a^{-1}) \sum_{n \geq 0} \frac{q^{2n+1} (q; q^2)_n}{(-aq, -a^{-1}q; q^2)_{n+1}} = \sum_{n \geq 0} (-a)^n q^{\frac{n(n+1)}{2}} - (q)_\infty \sum_{n \geq 0} q^{3n^2+n} a^{3n} (1-a^2 q^{4n+2}) \left(\sum_{m \in \mathbb{Z}} q^{m^2} a^m \right)^{-1}.$$

3. THE UNIVERSAL MOCK THETA FUNCTION g_3

In this section, we study the universal mock theta function $g_3(w; q)$ defined in (1.11). Theorem 3.1 gives three different q -hypergeometric series representations of $g_3(w; q)$ inside the unit disc. Theorem 3.2 shows that these series represent different functions outside \mathbb{D} that are related to each other by partial theta functions. To state our results, define for $|q| < 1$ and $w \in \mathbb{C} \setminus \{0\}$ with $w \neq q^\ell$ ($\ell \in \mathbb{Z}$)

$$g_{3,1}(w; q) := g_3(w; q), \\ g_{3,2}(w; q) := -\frac{1}{w} + \frac{1}{w(1-w)} R(w; q).$$

Additionally, for $w \neq q^{-\ell}$ ($\ell \in \mathbb{N}$) with $|w| > |q|$, we let

$$g_{3,3}(w; q) := \sum_{n \geq 0} \frac{w^{-n} q^n}{(w)_{n+1}}.$$

Theorem 3.1. *For $|q| < 1$ and $w \in \mathbb{C}$ with the same restrictions as above, we have that*

$$g_{3,1}(w; q) = g_{3,2}(w; q) = g_{3,3}(w; q).$$

To describe our next result concerning these representations outside the unit disc (for certain ranges of w), we require further functions. Generalizing the partial theta function $\psi(q)$ defined in (1.1), and the function $S(q)$, given in (1.6), we define for $|q| < 1$ the (two-variable) functions

$$(3.1) \quad \begin{aligned} \psi_1(w; q) &:= -w - w^2 \sum_{n \geq 0} \binom{12}{n} w^{\frac{n-1}{2}} q^{\frac{n^2-1}{24}}, \\ \psi_2(w; q) &:= \sum_{n \geq 0} w^n q^{\frac{n(n+1)}{2}}. \end{aligned}$$

Moreover if $w \neq q^\ell$ ($\ell \in \mathbb{Z}$), let

$$S_2(w; q) := \frac{w^2}{(wq, w^{-1})_\infty} \psi_2(-w^2; q).$$

Remark. Using the fact that $(-1)^{(n-1)/2} = \binom{-1}{n}$, we have that $\psi_1(-1; q) = 1 - \psi(q)$. Moreover $2S_2(-1; q) = S(q)$.

Theorem 3.2. *For $|q| < 1$ and $w \in \mathbb{C} \setminus \{0\}$ with $w \neq q^\ell$ ($\ell \in \mathbb{Z}$) we have*

$$(3.2) \quad g_{3,2}(w; q^{-1}) = \psi_1(w^{-1}; q) + S_2(w^{-1}; q).$$

Moreover if $w \neq q^\ell$ ($\ell \in \mathbb{N}_0$) and $w \neq 0$, we have that

$$(3.3) \quad g_{3,3}(w; q^{-1}) = \psi_1(w^{-1}; q).$$

Proof of Theorem 3.1. To prove the first equality stated in Theorem 3.1, we use the identities

$$(3.4) \quad R(w; q) = \frac{1-w}{(q)_\infty} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{\frac{3n^2+n}{2}}}{1-wq^n},$$

$$(3.5) \quad g_3(w; q) = \frac{1}{(q)_\infty} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{\frac{3n^2+3n}{2}}}{1-wq^n},$$

the first of which is contained in the proof of Theorem 7.1 of [27], and the second of which may be found in (3.3) of [14]. Using these, and the fact that $\sum_{n \in \mathbb{Z}} (-1)^n q^{(3n^2+n)/2} = (q)_\infty$, we obtain that

$$g_{3,1}(w; q) - g_{3,2}(w; q) = \frac{1}{w} - \frac{1}{w(q)_\infty} \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{3n^2+n}{2}} = 0.$$

To prove the second equality stated in Theorem 3.1, we first rewrite

$$(1-w)g_{3,3}(w; q) = F\left(0, w; \frac{q}{w}\right).$$

We then use (2.4) to deduce that $g_{3,3}(w; q) = g_{3,1}(w; q)$. □

Proof of Theorem 3.2. Set $\rho = q^{-1}$. We use the fact that for $n \in \mathbb{N}_0$ we have

$$(3.6) \quad (a; \rho)_n = (a^{-1}; q)_n (-a)^n \rho^{\frac{n(n-1)}{2}}.$$

We obtain, employing (2.7) with $a = -w^{-1}$,

$$\begin{aligned} g_{3,2}(w; \rho) &= -w^{-1} + \frac{1}{w(1-w)} \sum_{n \geq 0} \frac{\rho^{n^2}}{(w\rho, w^{-1}\rho; \rho)_n} = -w^{-1} + \frac{1}{w(1-w)} \sum_{n \geq 0} \frac{q^n}{(wq, w^{-1}q)_n} \\ &= -w^{-1} - w^{-2} \sum_{n \geq 0} (-1)^n w^{-3n} q^{\frac{n(3n+1)}{2}} (1 - w^{-2} q^{2n+1}) + S_2(w^{-1}; q). \end{aligned}$$

We note that the second equality also yields convergence in the claimed domain. Questions of convergence follow similar in all cases, therefore we will not mention them anymore.

Identity (3.2) is now deduced from the easily verified identity

$$(3.7) \quad \sum_{n \geq 0} (-1)^n w^{-3n} q^{\frac{n(3n+1)}{2}} (1 - w^{-2} q^{2n+1}) = \sum_{n \geq 0} \binom{12}{n} w^{-\frac{n+1}{2}} q^{\frac{n^2-1}{24}}.$$

To prove (3.3), we use again (3.6) to obtain

$$(3.8) \quad g_{3,3}(w; \rho) = \sum_{n \geq 0} \frac{w^{-n} \rho^n}{(w; \rho)_{n+1}} = \sum_{n \geq 0} \frac{w^{-2n-1} (-1)^{n+1} q^{\frac{n^2-n}{2}}}{(w^{-1})_{n+1}}.$$

Using (2.1), (2.8), and (3.7), we find that

$$\begin{aligned} \sum_{n \geq 0} \frac{(-1)^n w^{-2n} q^{\frac{n^2-n}{2}}}{(w^{-1}q)_n} &= \lim_{x \rightarrow \infty} F\left(x, w^{-1}; \frac{w^{-2}}{qx}\right) = \lim_{x \rightarrow \infty} \left(\frac{1-w^{-1}}{1-\frac{w^{-2}}{qx}} + \frac{w^{-1}-w^{-2}}{1-\frac{w^{-2}}{qx}} F\left(x, w^{-1}; \frac{w^{-2}}{x}\right) \right) \\ &= (1-w^{-1}) + w^{-1} (1-w^{-1}) \sum_{n \geq 0} \frac{(-1)^n w^{-2n} q^{\frac{n(n+1)}{2}}}{(w^{-1}q)_n} \\ &= (1-w^{-1}) \left(1 + w^{-1} \sum_{n \geq 0} \binom{12}{n} w^{-\frac{n+1}{2}} q^{\frac{n^2-1}{24}} \right) = (1-w) \psi_1(w^{-1}; q). \end{aligned}$$

Inserting this into (3.8) now yields the result. \square

Remark. As is suggested in [19], the partial theta function $\psi_1(\zeta; q)$, for ζ a root of unity, is closely related to the *shadow* of the mock theta function $g_3(\zeta; q)$ (see [27], e.g., for a definition of this term and more on mock modular forms and [18] for the shadow of g_3).

4. THE UNIVERSAL MOCK THETA FUNCTION g_2

Just as in the case of the universal mock theta function $g_3(w; q)$ described in §3, we can give alternate q -hypergeometric series expressions for the universal mock theta function $g_2(w; q)$ inside \mathbb{D} , and moreover, show that these different expressions are related by partial theta functions outside \mathbb{D} . One particularly nice expression involves a combinatorial rank generating function. To describe this, we let for $|q| < 1$, and $w \in \mathbb{C} \setminus \{0\}$ with $w \neq q^\ell$ ($\ell \in \mathbb{Z} \setminus \{0\}$),

$$O_2(w; q) := \sum_{n \geq 0} \frac{(-1)_n q^{\frac{n(n+1)}{2}}}{(wq, w^{-1}q)_n},$$

denote the overpartition rank generating function (defined in Proposition 1.1 of [20]). We point out that the overpartition function $O_2(w; q)$ is equal to McIntosh's function $K_2(w; q)$, and the universal mock theta function $g_2(w; q)$ is equal to McIntosh's function $H(w; q)$ (see [21]).

Analogous to the series $g_{3,j}(w; q)$, $1 \leq j \leq 3$, we define for $|q| < 1$ and $w \in \mathbb{C} \setminus \{0\}$ with $w \neq q^\ell$ ($\ell \in \mathbb{Z}$)

$$\begin{aligned} g_{2,1}(w; q) &:= g_2(w; q), \\ g_{2,2}(w; q) &:= \frac{1+w}{2w(1-w)} O_2(w; q) - \frac{1}{2w}. \end{aligned}$$

We also define for $|q| < 1$ and $w \neq q^{-\ell}$ ($\ell \in \mathbb{N}$) and $|w| > 1$ the function

$$g_{2,3}(w; q) := -\frac{1+w}{2w^2} \sum_{n \geq 0} \frac{(-wq)_n}{(wq)_n} w^{-n} - \frac{1}{2w}.$$

Our next result is an analogue to Theorem 3.1 for $g_2(w; q)$ when $|q| < 1$.

Theorem 4.1. *For $|q| < 1$ and $w \in \mathbb{C}$, assuming the same restrictions as above, we have that*

$$(4.1) \quad g_{2,1}(w; q) = g_{2,2}(w; q) = g_{2,3}(w; q).$$

Next we will analyze these functions outside the unit disc. We introduce partial theta functions for $|q| < 1$, defined for $w \in \mathbb{C}$ as follows:

$$(4.2) \quad \psi_3(w; q) := \sum_{n \geq 0} (-1)^{n+1} w^{2n+1} q^{n^2},$$

$$(4.3) \quad \psi_4(w; q) := \sum_{n \geq 1} w^{3n-2} q^{\frac{3n^2-n}{2}} (1 - wq^n).$$

Moreover if $w \neq 0$ and $w \neq -q^\ell$ ($\ell \in \mathbb{Z}$), we define

$$(4.4) \quad S_1(w; q) := \frac{(-q)_\infty}{(-w^{-1}, -wq)_\infty} (w^{-1} \psi_1(w; q) + 1),$$

and for $w \neq 0$ with $w \neq q^\ell$ ($\ell \in \mathbb{Z}$), we let

$$(4.5) \quad S_4(w; q) := \frac{(-q)_\infty}{(wq, w^{-1})_\infty} \psi_4(w; q).$$

In analogy to Theorem 3.2, we have the following relations between the series in Theorem 4.1 and partial theta functions.

Theorem 4.2. *For $|q| < 1$ and $w \in \mathbb{C} \setminus \{0\}$ with $w \neq q^\ell$ ($\ell \in \mathbb{Z}$), we have*

$$(4.6) \quad g_{2,1}(w; q^{-1}) = -w^{-2} \psi_3(w; q) - w^{-1} + S_4(w; q),$$

$$(4.7) \quad g_{2,2}(w; q^{-1}) = \psi_3(w^{-1}; q) + S_1(-w^{-1}; q).$$

Moreover for $|q| < 1$ and $w \in \mathbb{C}$ with $w \neq q^\ell$ ($\ell \in \mathbb{N}$) and $|w| > 1$, we have

$$(4.8) \quad g_{2,3}(w; q^{-1}) = \psi_3(w^{-1}; q).$$

Remark. It is not difficult to see that

$$\psi_4(w; q) = w^{-1} - w^{-2} + w^{-3} \psi_1(-w; q).$$

Remark. Note that

$$\psi_3(w^{-1}; q) + w^{-2}\psi_3(w; q) + w^{-1} = \sum_{n \in \mathbb{Z}} (-1)^{n+1} w^{2n-1} q^{n^2}$$

is the classical Jacobi theta function. In this way, by Theorem 4.1, we have two series, $g_{2,1}(w; q)$ and $g_{2,2}(w; q)$, that are equal to each other, and define a mock theta function inside the unit disc. On the other hand, by Theorem 4.2, outside of the disc these series are essentially each equal to half of a Jacobi theta function, but not the same half!

Proof of Theorem 4.1. By (3.2) of [21], we have that

$$g_{2,1}(w; q) = \frac{(1+w)}{2w(1-w)} O_2(w; q) - \frac{1}{2w} = g_{2,2}(w; q).$$

On the other hand, (2.3) yields

$$\frac{w}{w-1} \sum_{n \geq 0} \frac{(-1)_n q^{\frac{n(n+1)}{2}}}{(wq, w^{-1}q)_n} = F\left(-w, w; \frac{1}{w}\right) = \sum_{n \geq 0} \frac{(-wq)_n}{(wq)_n} w^{-n},$$

which can be used to show that $g_{2,2}(w; q) = g_{2,3}(w; q)$. \square

Proof of Theorem 4.2. To establish (4.6), we obtain, again using (3.6),

$$(4.9) \quad g_2(w; \rho) = \sum_{n \geq 0} \frac{(-\rho; \rho)_n \rho^{\frac{n(n+1)}{2}}}{(w, w^{-1}\rho; \rho)_{n+1}} = \frac{q}{(1-wq)(1-w^{-1})} \sum_{n \geq 0} \frac{(-q)_n q^n}{(w^{-1}q, wq^2)_n}.$$

Next we apply (2.6) with $B = -q$, $a = -1/w$, and $b = -wq$, and let $A \rightarrow 0$. This gives that the rightmost expression in (4.9) equals

$$(4.10) \quad \frac{q}{(1-w^{-1})} \sum_{n \geq 0} \frac{(w)_{n+1} (wq)^n}{(-wq)_{n+1}} + \frac{wq(-q)_\infty}{(wq, w^{-1})_\infty} \sum_{n \geq 0} \frac{(-1)^n w^{2n} q^{\frac{n^2+3n}{2}}}{(-wq)_{n+1}}.$$

To the first summand in (4.10) we apply (2.2) with $a = wq$, $b = -wq^2$ and $t = wq$ to obtain

$$(4.11) \quad -qw(1+qw)^{-1} F(w, -wq; wq) = \sum_{n \geq 1} (-1)^n w^{2n-1} q^{n^2}.$$

Next we use (2.3) to realize the second summand in (4.10) as

$$\frac{wq(-q)_\infty}{(wq, w^{-1})_\infty} F(-w, 0; -wq).$$

To this we apply (2.2) with $a = -wq$ and $t = -wq$, and let $b \rightarrow 0$, yielding

$$(4.12) \quad \frac{(-q)_\infty}{(wq, w^{-1})_\infty} \sum_{n \geq 1} w^{3n-2} q^{\frac{3n^2-n}{2}} (1-wq^n).$$

Combining (4.11) and (4.12) gives (4.6).

Next, to establish (4.7), we use (3.6) and find

$$(4.13) \quad g_{2,2}(w; \rho) = \frac{1+w}{2w(1-w)} \sum_{n \geq 0} \frac{(-1; \rho)_n \rho^{\frac{n(n+1)}{2}}}{(w\rho, w^{-1}\rho; \rho)_n} - \frac{1}{2w} = \frac{1+w}{2w(1-w)} \sum_{n \geq 0} \frac{(-1)_n q^n}{(wq, w^{-1}q)_n} - \frac{1}{2w}.$$

Then we apply an identity of Ramanujan, similar to (2.9), namely (3.18)_R of [7] with $c = -q^{-1}$, $a = -w$, and $b = -w^{-1}$ to see that

$$(4.14) \quad \sum_{n \geq 0} \frac{(-1)_n q^n}{(wq, w^{-1}q)_n} = \frac{w-1}{w+1} \sum_{n \geq 0} \frac{(-1)^n q^{\frac{n(n+1)}{2}} (-1)_n w^{-2n}}{(w^{-2}q^2; q^2)_n} + \frac{(-1)_\infty}{w(wq, w^{-1}q)_\infty} \sum_{n \geq 0} \frac{(-1)^n q^{\frac{n(n+1)}{2}} w^{-2n}}{(-w^{-1})_{n+1}}.$$

Using (2.8) and (3.7), we have that the second q -hypergeometric series on the right hand side equals

$$(1+w^{-1})^{-1} \sum_{n \geq 0} \binom{12}{n} (-w)^{\frac{1-n}{2}} q^{\frac{n^2-1}{24}}.$$

Next we use (2.3) followed by (2.5) to see that

$$(4.15) \quad \sum_{n \geq 0} \frac{(-1)^n q^{\frac{n(n+1)}{2}} (-1)_n w^{-2n}}{(w^{-2}q^2; q^2)_n} = (1-w^{-1}) F(w^{-1}, -w^{-1}; w^{-1}) = 1 + 2 \sum_{n \geq 1} (-1)^n w^{-2n} q^{n^2}.$$

Using (4.14) and (4.15) in (4.13) together with (3.1) now gives (4.7).

Equation (4.8) follows by first using (3.6) to write

$$(4.16) \quad g_{2,3}(w; \rho) = -\frac{(1+w)}{2w^2} \sum_{n \geq 0} \frac{(-w\rho; \rho)_n w^{-n}}{(w\rho; \rho)_n} - \frac{1}{2w} = -\frac{(1+w)}{2w^2} \sum_{n \geq 0} \frac{(-1)^n w^{-n} (-w^{-1}q)_n}{(w^{-1}q)_n} - \frac{1}{2w}.$$

After applying (2.5) to the rightmost series expression in (4.16) and simplifying we obtain (4.8). \square

5. THE UNIVERSAL MOCK THETA FUNCTION K

In this section, we study the universal mock theta function $K(w; q)$ as defined in (1.12). Analogous to the series $g_{k,j}(w; q)$, $2 \leq k \leq 3$, $1 \leq j \leq 3$ defined in §3 and §4, we consider for $|q| < 1$ and $w \neq 0$ and $w \neq q^{2\ell+1}$ ($\ell \in \mathbb{Z}$)

$$K_1(w; q) := \sum_{n \geq 1} \frac{(-1)^{n-1} q^{n^2} (q; q^2)_{n-1}}{(wq, w^{-1}q; q^2)_n}.$$

Moreover for $|q| < 1$ and $w \neq q^{-2\ell+1}$ ($\ell \in \mathbb{N}$) and $|w| > |q|$, we define

$$\kappa(w; q) := \sum_{n \geq 0} \frac{q^{n+1} w^{-n} (wq^2; q^2)_n}{(wq; q^2)_{n+1}}.$$

The functions $K(w; q)$ and $K_1(w; q)$ are discussed at length in relation to the function $K_2(w; q) = O_2(w; q)$ from §4 in [21]. We point out that $K(w; q)$, as given in (1.12), is defined for $|q| < 1$, and $w \in \mathbb{C} \setminus \{0\}$, where $w \neq q^{2\ell}$ ($\ell \in \mathbb{Z} \setminus \{0\}$).

We define the following partial theta function:

$$\psi_5(w; q) := \sum_{n \geq 0} \binom{n}{3} (-w)^{n-1} q^{\frac{n^2-1}{3}}.$$

If $w \neq 0$ and $w \neq q^{2\ell+1}$ ($\ell \in \mathbb{Z}$) let

$$s_1(w; q) := \frac{(q; q^2)_\infty}{w(wq, w^{-1}q; q^2)_\infty} (w^{-1} \psi_1(w; q^2) + 1).$$

Moreover if $w \neq q^{2\ell} (\ell \in \mathbb{Z} \setminus \{0\})$, we define

$$S_5(w; q) := \frac{w(q; q^2)_\infty}{(wq^2, w^{-1}q^2; q^2)_\infty} \psi_5(w; q).$$

We offer the following results for these functions inside and outside the disc.

Theorem 5.1. *For $|q| < 1$, and under the same restrictions as above, we have*

$$(5.1) \quad \kappa(w; q) = K_1(w; q) = -\frac{w}{(1-w)^2} \left(K(w; q) - \frac{(q; q^2)_\infty^3 (q^2; q^2)_\infty}{(wq, w^{-1}q)_\infty} \right).$$

Theorem 5.2. *For $|q| < 1$ we have for $w \neq q^{2\ell+1} (\ell \in \mathbb{N}_0)$ with $|w| > 1$*

$$(5.2) \quad \kappa(w; q^{-1}) = \frac{1}{1-w} \psi_2(w^{-1}; q).$$

If $w \neq 0$ and $w \neq q^{2\ell+1} (\ell \in \mathbb{Z})$, we have

$$(5.3) \quad K_1(w; q^{-1}) = \frac{w}{w-1} (\psi_2(w; q) + s_1(w; q)).$$

Moreover if $w \neq 0$ and $w \neq q^{2\ell} (\ell \in \mathbb{Z} \setminus \{0\})$, we obtain

$$(5.4) \quad K(w; q^{-1}) = (1-w)\psi_2(w; q) + S_5(w; q).$$

Proof of Theorem 5.1. The fact that the rightmost expression stated in (5.1) equals $K_1(w; q)$ may be found in (5.3) of [18]. To prove that $K_1(w; q) = \kappa(w; q)$, we modify the proof given in [6] of (3.6)_R to accommodate the parameter w . We have, applying (10.1) of [25] with q replaced by q^2 , $a = q^3/x$, $b = q$, $c = q^2$, $e = wq^3$, $f = w^{-1}q^3$, and letting $x \rightarrow 0$.

$$K_1(w; q) = \frac{q}{(1-wq)(1-w^{-1}q)} \lim_{x \rightarrow 0} \sum_{n \geq 0} \frac{(q^3/x, q, q^2; q^2)_n x^n}{(wq^3, w^{-1}q^3, q^2; q^2)_n} = \sum_{n \geq 0} \frac{q^{n+1} w^{-n} (wq^2; q^2)_n}{(wq; q^2)_{n+1}}.$$

□

Proof of Theorem 5.2. To prove (5.2), we find, using (3.6), that

$$(5.5) \quad \kappa(w; \rho) = \sum_{n \geq 0} \frac{w^{-n} \rho^{n+1} (w\rho^2; \rho^2)_n}{(w\rho; \rho^2)_{n+1}} = -\frac{1}{w} \sum_{n \geq 0} \frac{w^{-n} (w^{-1}q^2; q^2)_n}{(w^{-1}q; q^2)_{n+1}}.$$

We next apply the Rogers-Fine identity (2.2) with $q \rightarrow q^2$, and $a = w^{-1}q^2$, $b = w^{-1}q^3$, $t = w^{-1}$. From this and (5.5) we find that

$$\begin{aligned} \kappa(w; \rho) &= \frac{1}{1-w} \sum_{n \geq 0} q^{2n^2+n} w^{-2n} (1 + q^{2n+1} w^{-1}) \\ &= \frac{1}{1-w} \left(\sum_{n \in 2\mathbb{N}_0} q^{\frac{n(n+1)}{2}} w^{-n} + \sum_{n \in 2\mathbb{N}_0+1} q^{\frac{n(n+1)}{2}} w^{-n} \right) \\ &= \frac{1}{1-w} \psi_2(w^{-1}; q). \end{aligned}$$

Next, to prove (5.3), we note that by (3.6),

$$(5.6) \quad K_1(w; \rho) = \sum_{n \geq 0} \frac{(-1)^n \rho^{(n+1)^2} (\rho; \rho^2)_n}{(w\rho, w^{-1}\rho, \rho^2)_{n+1}} = \sum_{n \geq 0} \frac{q^{2n+1} (q; q^2)_n}{(wq, w^{-1}q; q^2)_{n+1}}.$$

We apply (2.9) to see that the rightmost expression in (5.6) equals

$$\frac{w}{w-1} \left(\sum_{n \geq 0} w^n q^{\frac{n(n+1)}{2}} - \frac{(q)_\infty \sum_{n \geq 0} (-1)^n w^{3n} q^{3n^2+n} (1-w^2 q^{4n+2})}{\sum_{n \in \mathbb{Z}} (-1)^n w^n q^{n^2}} \right).$$

Equation (5.3) now follows from $(q)_\infty / (q^2; q^2)_\infty = (q; q^2)_\infty$, (3.7), and the identity

$$\sum_{n \in \mathbb{Z}} (-1)^n w^n q^{n^2} = (q^2; q^2)_\infty (wq; q^2)_\infty (w^{-1}q; q^2)_\infty,$$

which follows from the Jacobi triple product identity (see [22] for example).

Finally, to prove (5.4), we have again using (3.6)

$$K(w; \rho) = \sum_{n \geq 0} \frac{(-1)^n \rho^{n^2} (\rho; \rho^2)_n}{(w\rho^2, w^{-1}\rho^2; \rho^2)_n} = \sum_{n \geq 0} \frac{q^{2n} (q; q^2)_n}{(wq^2, w^{-1}q^2; q^2)_n}.$$

To this we apply $(3.14)_R$ of [7] to find that

$$K(w; \rho) = (1-w)\psi_2(w; q) + \frac{w(q; q^2)_\infty}{(wq^2, w^{-1}q^2; q^2)_\infty} \sum_{n \geq 0} (-1)^n w^{3n} q^{3n^2+2n} (1+wq^{2n+1}).$$

It is straightforward to see that

$$\sum_{n=0}^{\infty} (-1)^n w^{3n} q^{3n^2+2n} (1+wq^{2n+1}) = \sum_{n \geq 0} \binom{n}{3} (-w)^{n-1} q^{\frac{n^2-1}{3}},$$

and the result now follows. \square

Remark. We note that $\psi_2(w; q) + w^{-1}\psi_2(w^{-1}; q) = \sum_{n \in \mathbb{Z}} w^n q^{\frac{n(n+1)}{2}}$ is a Jacobi theta function.

ACKNOWLEDGEMENTS

The authors thank Jeremy Lovejoy for several insightful comments. The first author was partially supported by the Alfried Krupp-prize, and the second author is grateful for the support of National Science Foundation grant DMS-1049553.

REFERENCES

- [1] K. Alladi, *A partial theta identity of Ramanujan and its number-theoretic interpretation*. Ramanujan J. 20 (2009), 329-339.
- [2] K. Alladi, *A combinatorial study and comparison of partial theta identities of Andrews and Ramanujan*. Ramanujan J. 23 (2010), 227-241,
- [3] A. Atkin and P. Swinnerton-Dyer, *Some properties of partitions*. Proc. London Math. Soc. 4 (1954), 84-106.
- [4] G. Andrews, *Partitions: Yesterday and Today*, New Zealand Mathematical Society, Wellington, (1979).
- [5] G. Andrews, *An introduction to Ramanujan's "lost" notebook*. Amer. Math. Monthly 86 (1979), 89-108.
- [6] G. Andrews, *Mordell integrals and Ramanujan's "lost" notebook*. Analytic number theory (Philadelphia, Pa., 1980), pp. 10-18, Lecture Notes in Math., 899, Springer, Berlin-New York, 1981.
- [7] G. Andrews, *Ramanujan's "lost" notebook. I. Partial θ functions*, *Adv. Math.* 41 (1981), 137-172.
- [8] G. Andrews and B. Berndt, *Ramanujan's Lost Notebook, Part I*, Springer, New York 2005.
- [9] B. Berndt and A. Yee, *Combinatorial proofs of identities in Ramanujan's lost notebook associated with the Rogers-Rine identity and false theta functions*. Annals of Combinatorics 7 (2003), 409-423.
- [10] B. Berndt, B. Kim and A. Yee, *Ramanujan's lost notebook: combinatorial proofs of identities associated with Heine's transformation or partial theta functions*. J. Combin. Theory Ser. A 117 (2010), 857-973.
- [11] K. Bringmann and O. Richter, *Zagier-type dualities and lifting maps for harmonic Maass-Jacobi forms*. Adv. Math. 225 (2010), 2298-2315.
- [12] F. Dyson, *Some guesses in the theory of partitions*. Eureka (Cambridge) 8 (1944), 10-15.

- [13] N. Fine, *Basic hypergeometric series and applications*. Mathematical Surveys and Monographs, 27. American Mathematical Society, Providence, RI, (1988).
- [14] B. Gordon and R. McIntosh, *A survey of classical mock theta functions*. To appear in *Developments in Mathematics*, Springer, (2011).
- [15] D. Hickerson, *A proof of the mock theta conjectures*. *Invent. Math.* 94 (1988), 639-660.
- [16] D. Hickerson, *On the seventh order mock theta functions*. *Invent. Math.* 94 (1988), 661-677.
- [17] K. Hikami, *Mock (false) theta functions as quantum invariants*. *Regul. Chaotic Dyn.* 10 (2005), 509-530.
- [18] S. Kang, *Mock Jacobi forms in basic hypergeometric series*. *Comp. Math.* 145 (2009), 553-565.
- [19] R. Lawrence and D. Zagier, *Modular forms and quantum invariants of 3-manifolds*. *Asian J. Math.* 3 (1999), 93-107.
- [20] J. Lovejoy, *Rank and conjugation for the Frobenius representation of an overpartition*. *Ann. Comb.* 9 (2005), 321-334.
- [21] R. McIntosh, *The H and K family of mock theta functions*. To appear in *Canad. J. Math.*, (2011-12).
- [22] K. Ono, *The web of modularity: arithmetic of the coefficients of modular forms and q -series*. CBMS Regional Conference Series in Mathematics, 102. American Mathematical Society, Providence, RI, (2004).
- [23] K. Ono, *Unearthing the visions of a master: harmonic Maass forms and number theory*. Proceedings of the 2008 Harvard-MIT Current Developments in Mathematics Conference, International Press, Somerville, MA, 2009, 347-454.
- [24] L. Rogers, *On two theorems of combinatorial analysis and some allied identities*. *Proc. London Math. Soc.* (2) 16 (1917), 315-336.
- [25] D. Sears, *On the transformation theory of basic hypergeometric functions*. *Proc. London Math. Soc.* (2) 53 (1951), 158-180.
- [26] D. Zagier, *Vassiliev invariants and a strange identity related to the Dedekind eta-function*. *Topology* 40 (2001), 945-960.
- [27] D. Zagier, *Ramanujan's mock theta functions and their applications (after Zwegers and Ono-Bringmann)*, Séminaire Bourbaki. Vol. 2007/2008. Astérisque No. 326 (2009), Exp. No. 986, vii-viii, 143-164 (2010).
- [28] S. Zwegers, *Mock theta functions*. Ph.D. Thesis, Universiteit Utrecht, (2002).

MATHEMATICAL INSTITUTE, UNIVERSITY OF COLOGNE, WEYERTAL 86-90, 50931 COLOGNE, GERMANY
E-mail address: `kbringma@math.uni-koeln.de`

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06520-8283 U.S.A.
E-mail address: `amanda.folsom@yale.edu`

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305, U.S.A.
E-mail address: `rhoades@math.stanford.edu`