

OVERPARTITIONS AND CLASS NUMBERS OF BINARY QUADRATIC FORMS

KATHRIN BRINGMANN AND JEREMY LOVEJOY

ABSTRACT. We show that the Zagier-Eisenstein series shares its non-holomorphic part with certain weak Maass forms whose holomorphic parts are generating functions for overpartition rank differences. This has a number of consequences, including exact formulas, asymptotics, and congruences for the rank differences as well as q -series identities of the mock theta type.

1. INTRODUCTION AND STATEMENT OF RESULTS

In this paper we relate overpartition analogues of Ramanujan's mock theta function $f(q)$ to the generating function for Hurwitz class numbers $H(n)$ of binary quadratic forms of discriminant $-n$. The generating function for $H(n)$ is the holomorphic part of the Zagier-Eisenstein series $\mathcal{F}(z)$ [15, 20], where

$$\mathcal{F}(z) := -\frac{1}{12} + \sum_{\substack{n \geq 1 \\ n \equiv 0,3 \pmod{4}}} H(n)q^n + \frac{(1+i)}{16\pi} \int_{-\bar{z}}^{i\infty} \frac{\Theta(\tau)}{(z+\tau)^{\frac{3}{2}}} d\tau. \quad (1.1)$$

Here $\Theta(z) := \sum_{n=-\infty}^{\infty} q^{n^2}$ is the classical theta function, where as usual $q = e^{2\pi iz}$ and $z = x + iy$. The series $\mathcal{F}(z)$ transforms like a weight $\frac{3}{2}$ modular form on $\Gamma_0(4)$, but it is non-holomorphic. It is the original example of a class of functions now called *weak Maass forms* [9] (see Section 2 for their definition).

Building on an idea of Zwegers [21], the first author and Ono [6, 7] have recently constructed infinite families of weak Maass forms arising from the generating function for Dyson's rank. Recall that Dyson [14] defined the *rank* of a partition to be the largest part minus the number of parts. A special case of the results in [6] says that Ramanujan's mock theta function

$$\begin{aligned} f(q) &= 1 + \sum_{n=1}^{\infty} \alpha(n)q^n := 1 + \sum_{n=1}^{\infty} \frac{q^{n^2}}{(1+q)^2(1+q^2)^2 \cdots (1+q^n)^2} \\ &= 1 + q - 2q^2 + 3q^3 - 3q^4 + 3q^5 - 5q^6 + \cdots, \end{aligned}$$

which counts the number of partitions with even rank minus the number of partitions with odd rank, is the holomorphic part of a weight $\frac{1}{2}$ weak Maass form.

Viewing this function in the framework of weak Maass forms has led to many applications, including an exact formula for the coefficients of $f(q)$ [6], asymptotics for the number of partitions of n with fixed rank m [4], and identities for rank differences [8]. For example, in [6] the first author

Date: October 9, 2008.

and Ono proved an exact formula for $\alpha(n)$ conjectured by Andrews and Dragonette [1, 12],

$$\alpha(n) = \frac{\pi}{(24n-1)^{\frac{1}{4}}} \sum_{k=1}^{\infty} \frac{(-1)^{\lfloor \frac{k+1}{2} \rfloor} A_{2k} \left(n - \frac{k(1+(-1)^k)}{4} \right)}{k} \cdot I_{\frac{1}{2}} \left(\frac{\pi \sqrt{24n-1}}{12k} \right). \quad (1.2)$$

Here $A_{2k}(n)$ denotes a Kloosterman-type sum and $I_{\frac{1}{2}}(x)$ is the usual Bessel function of order $\frac{1}{2}$.

In this paper we consider two analogues of Ramanujan's mock theta function $f(q)$ in the setting of overpartitions. For the coefficients of these two series we give exact formulas which are of a completely different nature than (1.2). Namely, we will exhibit formulas in terms of the Hurwitz class number $H(n)$.

Recall that an *overpartition* of n is a partition in which the first occurrence of a number may be overlined. For example, the 14 overpartitions of 4 are

$$4, \bar{4}, 3+1, \bar{3}+1, 3+\bar{1}, \bar{3}+\bar{1}, 2+2, \bar{2}+2, 2+1+1, \bar{2}+1+1, 2+\bar{1}+1, \bar{2}+\bar{1}+1, 1+1+1+1, \bar{1}+1+1+1.$$

As in the case of classical partitions, we define the rank of an overpartition to be its largest part minus the number of its parts, ignoring whether or not the parts are overlined.

Next recall the $M2$ -rank, which was introduced by the second author [17] based on the work in [3]. To define it, we use the notation $\ell(\cdot)$ to denote the largest part of an object, $n(\cdot)$ to denote the number of parts, and λ_o for the subpartition of an overpartition consisting of the odd non-overlined parts. Then the $M2$ -rank of an overpartition λ is

$$M2\text{-rank}(\lambda) := \left\lfloor \frac{\ell(\lambda)}{2} \right\rfloor - n(\lambda) + n(\lambda_o) - \chi(\lambda),$$

where $\chi(\lambda) = 1$ if the largest part of λ is odd and non-overlined and $\chi(\lambda) = 0$ otherwise. For example, the $M2$ -rank of the overpartition $5 + \bar{4} + 4 + \bar{3} + 1 + 1$ is $3 - 6 + 3 - 1 = -1$.

Now let $\bar{p}_e(n)$ (resp. $\bar{p}_o(n)$, $M2_e(n)$, $M2_o(n)$) denote the number of overpartitions of n with even rank (resp. odd rank, even $M2$ -rank, odd $M2$ -rank). For instance we have $\bar{p}_e(4) = 2$, $\bar{p}_o(4) = 12$, $M2_e(4) = 6$, and $M2_o(4) = 8$. We shall be concerned with the rank differences

$$\begin{aligned} \bar{\alpha}(n) &:= \bar{p}_e(n) - \bar{p}_o(n), \\ \bar{\alpha}_2(n) &:= M2_e(n) - M2_o(n), \end{aligned}$$

whose generating functions are

$$\begin{aligned} \bar{f}(q) &:= \sum_{n=0}^{\infty} \bar{\alpha}(n) q^n = 1 + 2q - 4q^2 + 8q^3 - 10q^4 + 8q^5 - 8q^6 + \dots, \\ \bar{f}_2(q) &:= \sum_{n=0}^{\infty} \bar{\alpha}_2(n) q^n = 1 + 2q + 4q^2 - 2q^4 + 8q^5 + 8q^6 + \dots. \end{aligned}$$

In [5] the authors showed that $\bar{f}(q)$ is the holomorphic part of a weak Maass form of weight $\frac{3}{2}$ (note the different weight from the partition case). Similar arguments will be used to show that $\bar{f}_2(q)$ is also the holomorphic part of a weight $\frac{3}{2}$ weak Maass form. Then we shall see that the non-holomorphic parts corresponding to $\bar{f}(q)$ and $\bar{f}_2(q)$ essentially match that of $\mathcal{F}(z)$. It turns out that the modular forms resulting from cancelling these non-holomorphic parts can also be written in terms of $H(n)$, yielding exact formulas for $\bar{\alpha}(n)$ and $\bar{\alpha}_2(n)$.

We use $\mathcal{H}(q)$ to denote the generating function for the Hurwitz class numbers (i.e. the holomorphic part of $\mathcal{F}(z)$):

$$\mathcal{H}(q) := -\frac{1}{12} + \sum_{\substack{n \geq 1 \\ n \equiv 0,3 \pmod{4}}} H(n)q^n = -\frac{1}{12} + \frac{1}{3}q^3 + \frac{1}{2}q^4 + q^7 + q^8 + q^{11} + \dots.$$

Theorem 1.1. *We have*

(i)

$$\bar{f}(-q) = -16\mathcal{H}(q) - \frac{1}{3}\Theta^3(z),$$

(ii)

$$\bar{f}_2(q) = -8\mathcal{H}(q) + \frac{1}{3}\Theta^3(z).$$

Now to express the coefficients $\bar{\alpha}(n)$ and $\bar{\alpha}_2(n)$ in terms of class numbers, we recall that Gauss proved that if we define $r(n)$ by

$$\sum_{n \geq 0} r(n)q^n := \Theta^3(z) = 1 + 6q + 12q^2 + 8q^3 + 6q^4 + \dots,$$

then we have

$$r(n) = \begin{cases} 12H(4n) & \text{if } n \equiv 1, 2 \pmod{4}, \\ 24H(n) & \text{if } n \equiv 3 \pmod{8}, \\ r(n/4) & \text{if } n \equiv 0 \pmod{4}, \\ 0 & \text{if } n \equiv 7 \pmod{8}. \end{cases} \quad (1.3)$$

The following formulas are then immediate:

Corollary 1.2. *We have*

(i)

$$(-1)^n \bar{\alpha}(n) = \begin{cases} -4H(4n) & \text{if } n \equiv 1, 2 \pmod{4}, \\ -24H(n) & \text{if } n \equiv 3 \pmod{8}, \\ -16H(n) & \text{if } n \equiv 7 \pmod{8}, \\ -16H(n) - \frac{1}{3}r(n/4) & \text{if } n \equiv 0 \pmod{4}. \end{cases} \quad (1.4)$$

(ii)

$$\bar{\alpha}_2(n) = \begin{cases} 4H(4n), & \text{if } n \equiv 1, 2 \pmod{4} \\ 0 & \text{if } n \equiv 3 \pmod{8} \\ -8H(n), & \text{if } n \equiv 7 \pmod{8} \\ -8H(n) + \frac{1}{3}r(n/4) & \text{if } n \equiv 0 \pmod{4}. \end{cases} \quad (1.5)$$

Four remarks.

1) Theorem 1.1 further emphasizes the role that rank differences play in linking partitions to automorphic forms which are not classical modular forms. In addition to the mock theta function $f(q)$ described earlier, we recall that the generating function for the number of partitions into distinct parts with even rank minus the number of partitions into distinct parts with odd rank gives rise to a Maass waveform [2, 11].

2) The method of proof yielding exact formulas for $\bar{\alpha}(n)$ and $\bar{\alpha}_2(n)$ is completely different from the one used in [6] to obtain (1.2). In [6] the authors use Maass-Poincaré series whereas here we employ relations between non-holomorphic parts of weak Maass forms.

3) One obvious application of (1.4) and (1.5) is to use facts about class numbers to learn about rank differences. For example, the above equations imply that $\bar{\alpha}(n)$ and $\bar{\alpha}_2(n)$ only grow polynomially (like n^k where $1/2 - \epsilon < k < 1/2 + \epsilon$, to be precise), whereas the coefficients $\alpha(n)$ grow exponentially. From a combinatorial perspective, there are many surprising consequences. Even before appealing to the vast knowledge about class numbers, there are immediate relations, such as

$$\bar{p}_e(4n+1) - \bar{p}_o(4n+1) = M2_e(4n+1) - M2_o(4n+1),$$

which seem rather unlikely given the disparate definitions of Dyson's rank and the $M2$ -rank.

4) It would be interesting to see whether combinatorial properties of overpartitions could be employed to obtain information about class numbers. For example, is there some natural involution on overpartitions which changes the parity of the rank (or $M2$ -rank) but which is not defined on a subset whose size clearly corresponds to class numbers?

In Corollaries 1.3 and 1.4 below we give a brief indication of what can be said about overpartitions by combining class number formulas with (1.4) and (1.5). Corollary 1.3 contains some exact formulas at p^{2k} and $2p^{2k}$, while Corollary 1.4 gives a couple of congruences in arithmetic progressions whose members are highly divisible by powers of 4. We state these for $\bar{f}(q)$, but similar statements hold for $\bar{f}_2(q)$, and of course these are just a few among endless possibilities.

Corollary 1.3. *If p is an odd prime, then we have*

$$\bar{p}_e(p^{2k}) - \bar{p}_o(p^{2k}) = \begin{cases} 2p^k, & p \equiv 1 \pmod{4}, \\ \frac{2p^k(p+1)-4}{p-1}, & p \equiv 3 \pmod{4}, \end{cases} \quad (1.6)$$

$$\bar{p}_e(2p^{2k}) - \bar{p}_o(2p^{2k}) = \begin{cases} -4p^k, & p \equiv 1, 3 \pmod{8}, \\ \frac{-4p^k(p+1)+8}{p-1}, & p \equiv 5, 7 \pmod{8}. \end{cases} \quad (1.7)$$

Corollary 1.4. *We have the following congruences:*

(i) *For any $a \geq 1$, $n \geq 0$, and $t = 1$ or 2 , we have*

$$\bar{p}_e(4^{a+1}n + t4^a) \equiv \bar{p}_o(4^{a+1}n + t4^a) \pmod{2^{a+2} - 3}.$$

(ii) *For any $a \geq 1$ and $n \geq 0$, we have*

$$\bar{p}_e(2 \cdot 4^{a+1}n + 3 \cdot 4^a) \equiv \bar{p}_o(2 \cdot 4^{a+1}n + 3 \cdot 4^a) \pmod{2^{a+1} - 1}.$$

For example, if we take $a = 4$ and $t = 1$ in (i) of Corollary 1.4, then we obtain

$$\bar{p}_e(1024n + 256) \equiv \bar{p}_o(1024n + 256) \pmod{61}.$$

Another application is to relate overpartition rank differences to other partition-theoretic functions which have connections to class numbers. To give one example, using work of Ono and Sze [19] we may deduce the following:

Corollary 1.5. *Let $C_4(n)$ denote the number of partitions of n which are 4-cores. If $8n + 5$ is square-free, then we have*

$$\begin{aligned} 8C_4(n) &= \bar{p}_e(8n+5) - \bar{p}_o(8n+5) \\ &= M2_e(8n+5) - M2_o(8n+5). \end{aligned} \quad (1.8)$$

Thus all of the congruences and identities for $C_4(n)$ in [19] apply to $\bar{p}_e(8n+5) - \bar{p}_o(8n+5)$ and $M2_e(8n+5) - M2_o(8n+5)$ as well.

As a final application of Theorem 1.1, we use known generating functions for $\bar{f}(q)$, $\bar{f}_2(q)$, and class numbers to deduce q -series identities. We give four examples.

Corollary 1.6. *We have*

(i)

$$\frac{4}{\Theta(z+1/2)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+n}}{(1+q^n)^2} - \frac{8}{\Theta(z)} \sum_{n \geq 0} \frac{nq^{n^2}(1-q^{2n})}{(1+q^{2n})} = \Theta^3(z+1/2), \quad (1.9)$$

(ii)

$$\frac{4}{\Theta(z+1/2)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+2n}}{(1+q^{2n})^2} + 4 \prod_{n=1}^{\infty} \frac{(1-q^{4n-2})}{(1-q^{4n})} \sum_{n \geq 0} \frac{q^{n^2+3n+1}}{(1-q^{2n+1})^2} = \Theta^3(z), \quad (1.10)$$

(iii)

$$\frac{4}{\Theta(z)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+n}}{(1+(-q)^n)^2} - \frac{8}{\Theta(z+1/2)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+2n}}{(1+q^{2n})^2} = -\Theta^3(z), \quad (1.11)$$

(iv)

$$\frac{1}{\Theta(z)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+n}}{(1+(-q)^n)^2} + \frac{1}{\Theta(z+1/2)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+2n}}{(1+q^{2n})^2} = -6\mathcal{H}(q). \quad (1.12)$$

2. WEAK MAASS FORMS AND THE PROOF OF THEOREM 1.1

Here we prove Theorem 1.1. Let us first recall the notion of a weak Maass form. If $k \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$, $z = x + iy$ with $x, y \in \mathbb{R}$, then the weight k hyperbolic Laplacian is given by

$$\Delta_k := -y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) + ik y \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right). \quad (2.1)$$

If v is odd, then define ϵ_v by

$$\epsilon_v := \begin{cases} 1 & \text{if } v \equiv 1 \pmod{4}, \\ i & \text{if } v \equiv 3 \pmod{4}. \end{cases} \quad (2.2)$$

Moreover we let χ be a Dirichlet character. A (*harmonic*) *weak Maass form of weight k with Nebentypus χ on a subgroup $\Gamma \subset \Gamma_0(4)$* is any smooth function $g : \mathbb{H} \rightarrow \mathbb{C}$ satisfying the following:

(1) For all $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ and all $z \in \mathbb{H}$, we have

$$g(Az) = \left(\frac{c}{d} \right) \epsilon_d^{-2k} \chi(d) (cz+d)^k g(z).$$

(2) We have that $\Delta_k g = 0$.

(3) The function $g(z)$ has at most linear exponential growth at all the cusps of Γ .

In [5] the authors related $\bar{f}(q)$ to a weak Maass form. To be more precise, define the function

$$\bar{\mathcal{M}}(z) := \bar{f}(q) - \bar{\mathcal{N}}(z),$$

with

$$\overline{\mathcal{N}}(z) := \frac{\sqrt{2}}{\pi i} \int_{-\bar{z}}^{i\infty} \frac{\Theta(\tau + \frac{1}{2})}{(-i(\tau + z))^{\frac{3}{2}}} d\tau.$$

Then the function $\overline{\mathcal{M}}(z)$ is a weak Maass form of weight $\frac{3}{2}$ on $\Gamma_0(16)$. (If no mention is made of the Nebentypus, then it is trivial.) We have a similar result for the function $\overline{f}_2(q)$. For this we let

$$\overline{\mathcal{M}}_2(z) := \overline{f}_2(q) - \overline{\mathcal{N}}_2(z),$$

where

$$\overline{\mathcal{N}}_2(z) := -\frac{i}{\sqrt{2}\pi} \int_{-\bar{z}}^{i\infty} \frac{\Theta(\tau)}{(-i(\tau + z))^{\frac{3}{2}}} d\tau.$$

We show

Theorem 2.1. *The function $\overline{\mathcal{M}}_2(z)$ is a weak Maass form of weight $\frac{3}{2}$ on $\Gamma_0(16)$.*

Idea of proof of Theorem 2.1. The proof is quite similar to the case of $\mathcal{M}(z)$ treated in [5]. From [17] and the Watson-Whipple transformation, we have that

$$\overline{f}_2(q) = \frac{4}{\Theta(z + \frac{1}{2})} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+2n}}{(1 + q^{2n})^2}. \quad (2.3)$$

Next define the function

$$\overline{M}_{2,r}(z) = \overline{M}_{2,r}(q) := \frac{4}{\Theta(z + \frac{1}{2})} \sum_{n \in \mathbb{Z}} \frac{(-1)^{n+1} q^{n^2}}{1 + e^{2\pi i r} q^{2n}}.$$

This function is related to $\overline{f}_2(q)$ via

$$\overline{f}_2(q) = \frac{1}{2\pi i} \frac{\partial}{\partial r} (\overline{M}_{2,r}(q)) \Big|_{r=0}.$$

One first determines a transformation law of $\overline{M}_{2,r}(q)$ under inversion. Let

$$\eta(z) := q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$$

be the Dedekind η -function.

Lemma 2.2. *We have*

$$\overline{M}_{2,r} \left(-\frac{1}{z} \right) = -\sqrt{2}i(-iz)^{\frac{1}{2}} \overline{O}_{2,-irz}(z) - 4\sqrt{2}(-iz)^{-\frac{1}{2}} I_r(z),$$

where

$$\begin{aligned} \overline{O}_{2,r}(z) &:= \frac{\eta(4z)}{\eta^2(8z)} \sum_{\substack{m \in \mathbb{Z} \\ m \text{ odd}}} \frac{q^{\frac{m^2}{2}}}{1 - e^{\pi i r} q^{2m}}, \\ I_r(z) &:= \int_{\mathbb{R}} \frac{e^{-\frac{2\pi i x^2}{z}}}{1 - e^{2\pi i r - \frac{4\pi i x}{z}}} dx. \end{aligned}$$

Now let

$$I(z) := \frac{1}{2\pi i} \frac{\partial}{\partial r} (I_r(z)) \Big|_{r=0}.$$

We relate $I(z)$ to a theta integral. As in [7] or [5], one can prove:

Lemma 2.3. *We have*

$$I(z) = \frac{z^2}{8\sqrt{2\pi}} \int_0^\infty \frac{\Theta\left(\frac{iu}{4}\right)}{(-i(iu+z))^{\frac{3}{2}}} du.$$

Differentiating Lemma 2.2 and using Lemma 2.3 yields

Lemma 2.4. *We have*

$$\bar{f}_2\left(-\frac{1}{z}\right) = -\frac{(-iz)^{\frac{3}{2}}}{\sqrt{2}} \bar{O}_2\left(\frac{z}{8}\right) + \frac{(-iz)^{\frac{3}{2}}}{4} \int_0^\infty \frac{\Theta\left(\frac{iu}{4}\right)}{(-i(iu+z))} du,$$

where

$$\bar{O}_2(z) := \frac{\eta(4z)}{\eta^2(8z)} \sum_{\substack{m \in \mathbb{Z} \\ m \text{ odd}}} \frac{q^{\frac{m^2}{2}+2m}}{(1-q^{2m})^2}.$$

It is not hard to see that $\bar{\mathcal{N}}_2(z)$ introduces the same error integral as $\bar{f}_2(q)$ under inversion. Now one can finish the proof as in [6]. \square

We next turn to the proof of Theorem 1.1. For this define the function

$$\bar{g}(z) := \bar{f}(-q) + 16\mathcal{H}(z).$$

We must show that $\bar{g}(z) = -\frac{1}{3}\Theta^3(z)$.

We start by showing that $\bar{g}(z)$ is a weakly holomorphic modular form on $\Gamma_0(16)$. For this we compute that $\bar{\mathcal{M}}\left(z + \frac{1}{2}\right)$ is a weak Maass form of weight $\frac{3}{2}$ on $\Gamma_0(16)$ whose holomorphic part is given by $\bar{f}(-q)$. Thus $\bar{g}(z)$ is the holomorphic part of the weak Maass form $\bar{\mathcal{M}}\left(z + \frac{1}{2}\right) + 16\mathcal{F}(z)$. We now check that this weak Maass form doesn't have a non-holomorphic part. Indeed, the non-holomorphic part of $\bar{\mathcal{M}}\left(z + \frac{1}{2}\right)$ is given by

$$\frac{-\sqrt{2}}{\pi i} \int_{-\bar{z}}^{i\infty} \frac{\Theta(\tau)}{(-i(\tau+z))^{\frac{3}{2}}} d\tau. \quad (2.4)$$

This agrees with the non-holomorphic part of $-16\mathcal{F}(z)$.

We next claim that $\bar{g}(z)$ is actually a holomorphic modular form. It is known (see [13, Prop 1.2.4], for example) that if the coefficients of a weakly holomorphic modular form grow at most polynomially, then it is a holomorphic modular form. To take advantage of this fact in the case of $\bar{g}(z)$, we multiply by $\Theta(z)$. Then we have

$$\bar{g}(z)\Theta(z) = 4 \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+n}}{(1+(-q)^n)^2} + 16\mathcal{H}(z)\Theta(z), \quad (2.5)$$

the first term on the right hand side following from the generating function [5]

$$\bar{f}(q) = \frac{4}{\Theta(z+1/2)} \sum_{n \in \mathbb{Z}} \frac{(-1)^n q^{n^2+n}}{(1+q^n)^2}. \quad (2.6)$$

Now the coefficients in this first term of (2.5) can be expressed in terms of sum-of-divisor functions, and hence grow at most polynomially. Moreover, it is well-known that the Hurwitz class numbers grow polynomially, thus so do the coefficients in the product of $\mathcal{H}(z)$ and $\Theta(z)$.

Hence we have that $\bar{g}(z)\Theta(z)$ is a holomorphic modular form of weight 2 on $\Gamma_0(16)$. To show that it is equal to $\frac{-1}{3}\Theta^4(z)$ we compute that the q -expansions agree up to q^4 . This then completes the proof of part (i) of Theorem 1.1.

The proof of part (ii) is essentially the same. There we use Theorem 2.1 to cancel the non-holomorphic parts of $\overline{\mathcal{M}}_2(z)$ and $\mathcal{F}(z)$, and for the polynomial growth we appeal to the generating function for the $M2$ -rank (2.3).

3. SKETCHES OF PROOFS OF THE COROLLARIES

Proof of Corollary 1.3 and 1.4. These are simple calculations using Corollary 1.2, equation and the following fact [10, p.273]: If $-n = Df^2$ where D is a fundamental discriminant, then

$$H(n) = \frac{h(D)}{w(D)} \sum_{d|f} \mu(d) \left(\frac{D}{d}\right) \sigma_1(f/d). \quad (3.1)$$

Here $h(D)$ is the class number of $\mathbb{Q}(\sqrt{D})$, $w(D)$ is half the number of units in the ring of integers of $\mathbb{Q}(\sqrt{D})$, $\sigma_1(n)$ is the sum of the divisors of n , and $\mu(n)$ is the Möbius function. \square

Proof of Corollary 1.5. It is shown in [19] that if $8n + 5$ is square-free, then

$$C_4(n) = \frac{1}{2}h(-32n - 20).$$

Invoking (3.1) shows that in this case

$$\bar{p}_e(8n + 5) - \bar{p}_o(8n + 5) = 4h(-32n - 20),$$

thus establishing (1.8). \square

Proof of Corollary 1.6. Kronecker [16], Mordell [18], and others have given nice generating functions involving certain class numbers $F(n)$ and $G(n)$. The definitions of these class numbers are not important here, only that $H(n) = G(n) - F(n)$ and $r(n) = 24F(n) - 12G(n)$. Thus we may recast the main theorem in terms of these class numbers,

$$\bar{f}(q) = -16 \sum_{n \geq 0} F(n)(-q)^n + \Theta^3(z + 1/2)$$

and

$$\bar{f}_2(q) = -8 \sum_{n \geq 0} F(n)q^n + \Theta^3(z).$$

Then (1.9) follows from (2.6) and Mordell's generating function for $F(n)$ [18], while (1.10) follows from (2.3) and Kronecker's generating function for $F(n)$ [16, Eq. (XI)]. Equations (1.11) and (1.12) follow from Theorem 1.1 and equations (2.6) and (2.3). \square

REFERENCES

- [1] G. Andrews, On the theorems of Watson and Dragonette for Ramanujan's mock theta functions, *Amer. J. Math.* **88** (1966), pages 454-490.
- [2] G.E. Andrews, F.J. Dyson, and D. Hickerson, Partitions and indefinite quadratic forms, *Invent. Math.* **91** (1988), 391-407.
- [3] A. Berkovich and F.G. Garvan, Some observations on Dyson's new symmetries of partitions, *J. Comb. Theory Ser. A* **100** (2002), 61-93.
- [4] K. Bringmann, Asymptotics for rank partition functions, *Trans. Amer. Math. Soc.*, accepted for publication.
- [5] K. Bringmann and J. Lovejoy, Dyson's rank, overpartitions, and weak Maass forms, *Int. Math. Res. Not.* (2007), rnm063.
- [6] K. Bringmann and K. Ono, The $f(q)$ mock theta function conjecture and partition ranks, *Invent. Math.* **165** (2006), 243-266.
- [7] K. Bringmann and K. Ono, Dyson's rank and weak Maass forms, *Ann. Math.*, to appear.
- [8] K. Bringmann, K. Ono, and R. Rhoades, Eulerian series as modular forms, *J. Amer. Math. Soc.* **21** (2008), pages 1085-1104.
- [9] J.H. Bruinier and J. Funke, On two geometric theta lifts, *Duke Math. J.* **125** (2004), 45-90.
- [10] H. Cohen, Sums involving the values at negative integers of L -functions of quadratic characters, *Math. Ann.* **217** (1975), 271-285.
- [11] H. Cohen, q -identities for Maass waveforms, *Invent. Math.* **91** (1988), 409-422.
- [12] L. Dragonette, Some asymptotic formulae for the mock theta series of Ramanujan, *Trans. Amer. Math. Soc.* **72** (1952), pages 474-500.
- [13] F. Diamond and J. Shurman, *A first course in modular forms*, Graduate Texts in Mathematics **228**, Springer-Verlag, New York, 2005.
- [14] F. Dyson, Some guesses in the theory of partitions, *Eureka (Cambridge)* **8** (1944), pages 10-15.
- [15] F. Hirzebruch and D. Zagier, Intersection numbers of curves on Hilbert modular surfaces and modular forms of Nebentypus, *Invent. Math.* **36** (1976), 57-113.
- [16] M. Kronecker, Über die Anzahl der verschiedenen Classen quadratischer Formen von negativer Determinante, *J. reine Angew. Math.* **57** (1860), 248-255.
- [17] J. Lovejoy, Rank and conjugation for a second Frobenius representation of an overpartition, *Ann. Comb.*, **12** (2008), 101-113.
- [18] L.J. Mordell, Note on class relation formulae, *Mess. Math.* **45** (1915), 76-80.
- [19] K. Ono and L. Sze, 4-core partitions and class numbers, *Acta Arith.* **65** (1997), 249-272.
- [20] D. Zagier, Nombres de classes et formes modulaires de poids $3/2$, *C.R. Acad. Sci. Paris Sér. A-B* **281** (1975), 883-886.
- [21] S. P. Zwegers, *Mock theta-functions and real analytic modular forms*, q -series with applications to combinatorics, number theory, and physics (Urbana, IL, 2000), 269-277, *Contemp. Math.*, **291**, Amer. Math. Soc., Providence, RI, 2001.

MATHEMATISCHES INSTITUT, UNIVERSITÄT KÖLN, WEYERTAL 86-90, D-50931 KÖLN, GERMANY

E-mail address: `kbringma@math.uni-koeln.de`

CNRS, LIAFA, UNIVERSITÉ DENIS DIDEROT - PARIS 7, CASE 7014, 75205 PARIS CEDEX 13, FRANCE

E-mail address: `lovejoy@liafa.jussieu.fr`