

# On the classification of lattices over $\mathbb{Q}(\sqrt{-3})$ , which are even unimodular $\mathbb{Z}$ -lattices

by

MICHAEL HENTSCHEL, ALOYS KRIEG

Lehrstuhl A für Mathematik, RWTH Aachen University, D-52056 Aachen  
michael.hentschel@mathA.rwth-aachen.de, krieg@mathA.rwth-aachen.de

GABRIELE NEBE

Lehrstuhl D für Mathematik, RWTH Aachen University, D-52056 Aachen  
gabriele.nebe@math.rwth-aachen.de

# 1 Introduction

The classification of even unimodular  $\mathbb{Z}$ -lattices is explicitly known only in the cases of rank 8, 16 and 24 (cf. [CS]). Given an imaginary quadratic number field  $K$  with discriminant  $d_K$  and ring of integers  $\mathcal{O}_K$ , Cohen and Resnikoff [CR] showed that there exists a free  $\mathcal{O}_K$ -module  $\Lambda$  of rank  $r$ , which is even and satisfies  $\det \Lambda = (2/\sqrt{d_K})^r$  if and only if  $r \equiv 0 \pmod{4}$ , where explicit examples can also be found in [DK]. Each such  $\mathcal{O}_K$ -module is an even unimodular  $\mathbb{Z}$ -lattice of rank  $2r$  and the associated Hermitian theta series is a Hermitian modular form of weight  $r$  for the full modular group. An explicit description of the isometry classes of these lattices has so far only been obtained for the Gaussian number field  $K = \mathbb{Q}(i)$  and  $r = 4, 8$  by Schiemann [S], where 1 resp. 3 isometry classes exist, and for  $r = 12$  by Kitazume and Munemasa [KM], where 28 isometry classes exist. It should be noted that similar results over the Hurwitz order can be found in [Q] and [BN].

In this paper we derive analogous results for  $\mathcal{O}_K$ -modules whenever  $K = \mathbb{Q}(i\sqrt{3})$  is Eisenstein's number field. There is exactly one isometry class for  $r = 4$  and  $r = 8$  and surprisingly there exist just 5 isometry classes if  $r = 12$ . In the proof we make use of the Niemeier classification and verify which of these lattices have got the structure of an  $\mathcal{O}_K$ -module. The main idea is that the orthogonal automorphism group of such a lattice must contain an element with minimal polynomial  $p(X) = X^2 - X + 1$ , which corresponds to  $\frac{1}{2}(1 + i\sqrt{3})I$ . Moreover we investigate the filtration of cusp forms spanned by the associated Hermitian theta series.

## 2 The mass of the Eisenstein lattices

Throughout the paper let  $K = \mathbb{Q}(i\sqrt{3})$  be Eisenstein's number field of discriminant  $-3$  with attached Dirichlet character

$$\chi_K(n) = \left(\frac{n}{3}\right), \quad n \in \mathbb{Z},$$

and ring of integers

$$\mathcal{O} := \mathbb{Z} + \mathbb{Z}\omega, \quad \omega = (1 + i\sqrt{3})/2,$$

which is Euclidean with respect to the norm  $N(a) = a\bar{a}$ . We consider  $K^r$ ,  $r \in \mathbb{N}$ , with the standard Hermitian scalar product

$$K^r \times K^r \rightarrow K, \quad (x, y) \mapsto \langle x, y \rangle := \bar{x}^{tr} y.$$

Let  $\mathcal{U}(r)$  denote the unitary group of rank  $r$ .

**Definition 1.**  $\Lambda \subset K^r$  is called an *Eisenstein lattice* of rank  $r$  if there exist linearly independent  $b_1, \dots, b_r \in K^r$  such that

- (i)  $\Lambda = \mathcal{O}b_1 + \dots + \mathcal{O}b_r$ ,
- (ii)  $\det(\langle b_j, b_k \rangle) = (2/i\sqrt{3})^r$ ,
- (iii)  $\langle \lambda, \lambda \rangle \in 2\mathbb{Z}$  for all  $\lambda \in \Lambda$ .

Two such lattices  $\Lambda, \Lambda'$  are called *isometric* if there exists an *isometry*  $U \in \mathcal{U}(r)$  such that

$$\Lambda' = U\Lambda.$$

The (unitary) *automorphism group* of  $\Lambda$  is defined by

$$\text{Aut}(\Lambda) := \{U \in \mathcal{U}(r); U\Lambda = \Lambda\}.$$

The Eisenstein lattices form a genus and its *mass* is

$$\mu_r := \sum_{\Lambda} \frac{1}{\#\text{Aut}(\Lambda)},$$

where we sum over representatives of the isometry classes of Eisenstein lattices of rank  $r$ .

Considering the symmetric bilinear form

$$K^r \times K^r \rightarrow \mathbb{Q}, \quad (x, y) \mapsto \text{Re}\langle x, y \rangle,$$

we may also consider  $\Lambda$  as an even unimodular  $\mathbb{Z}$ -lattice of rank  $2r$

$$\Lambda_{\mathbb{Z}} = \mathbb{Z}b_1 + \mathbb{Z}\omega b_1 + \dots + \mathbb{Z}b_r + \mathbb{Z}\omega b_r.$$

This yields an embedding of  $\text{Aut} \Lambda$  into the (orthogonal) automorphism group  $\text{Aut}_{\mathbb{Z}}(\Lambda)$  of  $\Lambda_{\mathbb{Z}}$ . Hence it is clear that Eisenstein lattices exist only if  $r$  is a multiple of 4.

**Theorem 1.** *The mass of the genus of Eisenstein lattices of rank  $r$ ,  $r \in \mathbb{N}$ ,  $4 \mid r$ , is given by*

$$\mu_r = \frac{1}{2^{r-1} \cdot r!} \cdot \prod_{j=1}^{r/2} |B_{2j} \cdot B_{2j-1, \chi_K}|,$$

where  $B_j$  (resp.  $B_{j, \chi_K}$ ) is the (generalized) Bernoulli number. In particular one has

$$\mu_4 = \frac{1}{2^7 \cdot 3^5 \cdot 5}, \quad \mu_8 = \frac{1}{2^{15} \cdot 3^{10} \cdot 5^2}, \quad \mu_{12} = \frac{691 \cdot 809 \cdot 1847}{2^{22} \cdot 3^{17} \cdot 5^3 \cdot 7 \cdot 11 \cdot 13}.$$

*Proof.* Hashimoto and Koseki [HaK] computed the mass  $\mu_r^*$  of odd unimodular  $\mathcal{O}$ -modules of rank  $r$ ,  $4 \mid r$ , to be

$$\mu_r^* = \frac{3^{r/2} + 1}{2^r \cdot r!} \cdot \prod_{j=1}^{r/2} |B_{2j} \cdot B_{2j-1, \chi_K}|,$$

where  $B_j$  (resp.  $B_{j, \chi_K}$ ) are the (generalized) Bernoulli numbers (cf. [Z], resp. [F]). Using the same counting argument as in [BN], Proposition 2.4, we obtain

$$\mu_r = \mu_r^* \cdot \frac{c_1}{c_2},$$

where  $c_1$  is the number of maximal isotropic subspaces of the orthogonal  $\mathbb{F}_3$ -vector space  $\mathbb{F}_3^r$  and where  $c_2$  is the number of maximal isotropic subspaces of the symplectic  $\mathbb{F}_3$ -vector space  $\mathbb{F}_3^r$ . These numbers can be found in [T], p. 78 resp. p. 174, and are equal to

$$c_1 = \prod_{j=1}^{r/2} (3^{j-1} + 1), \quad c_2 = \prod_{j=1}^{r/2} (3^j + 1).$$

This yields the first claim. The special values are obtained from an explicit calculation of the Bernoulli numbers exploiting the formulas in [Z].  $\square$

### 3 Classification of Eisenstein lattices

In order to obtain a classification of the Eisenstein lattices we make use of Theorem 1. First of all we use the fact from [DK] that there exists an Eisenstein lattice  $\Lambda_4$  of rank 4 given by its Gram matrix

$$\begin{pmatrix} 2I & C \\ -C & 2I \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad C = \frac{2i}{\sqrt{3}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

An easy computer calculation yields

$$\# \text{Aut}(\Lambda_4) = 2^7 \cdot 3^5 \cdot 5 = \frac{1}{\mu_4}.$$

In view of

$$\# \text{Aut}(\Lambda_4 \oplus \Lambda_4) = 2 \cdot (\# \text{Aut}(\Lambda_4))^2 = \frac{1}{\mu_8}$$

Theorem 1 implies

**Corollary 1.** *There is exactly one isometry class of Eisenstein lattices of rank 4 and of rank 8. Representatives are  $\Lambda_4$  and  $\Lambda_4 \oplus \Lambda_4$ .*

In the case of rank 12 we apply the Niemeier classification of even unimodular 24-dimensional  $\mathbb{Z}$ -lattices (cf. [CS]). We use the fact that  $\omega I$  is always contained in the automorphism group of an Eisenstein lattice. Its minimal polynomial over  $\mathbb{Q}$  is  $p(X) = X^2 - X + 1$ .

**Proposition 1.** *Each matrix  $B \in \mathbb{Z}^{2r \times 2r}$  with minimal polynomial  $p(X) = X^2 - X + 1$  is in  $\text{GL}(2r; \mathbb{Z})$  conjugate to*

$$\text{diag}(A, \dots, A), \quad A = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}.$$

*Proof.* Following Newman [N], p. 54 and Theorem III.12, the claim basically follows from the fact that  $\mathcal{O} = \mathbb{Z}[\omega]$  has class number 1.  $\square$

Now we are going to investigate which Niemeier lattices possess an automorphism with minimal polynomial  $p(X) = X^2 - X + 1$ .

**Lemma 1.** *Let  $L$  be a  $\mathbb{Z}$ -lattice, which has a decomposition into orthogonally indecomposable non-isometric lattices*

$$L = R_1^{n_1} \oplus \dots \oplus R_s^{n_s}.$$

*Then one has*

- a)  $\text{Aut}(L) = \times_{j=1}^s \text{Aut}(R_j^{n_j})$ .
- b)  $\text{Aut}(R_j^{n_j}) = \{\text{diag}(\phi_1, \dots, \phi_{n_s})\sigma_j; \phi_k \in \text{Aut}(R_j), \sigma_j \in S_{n_j}\}$ .
- c) *If  $\phi \in \text{Aut}(L)$  exists with minimal polynomial  $p(X) = X^2 - X + 1$ , then  $\phi$  belongs to the kernel of the group epimorphism  $\text{Aut}(L) \rightarrow S_{n_1} \times \dots \times S_{n_s}$  in a) and b).*

*Proof.* a), b) The results are well-known (cf. [Kn]).

c) Let  $P_\sigma$  be the integral orthogonal  $n \times n$  matrix associated to  $\sigma \in S_n$  in the  $j$ -component. Then  $p(\phi) = 0$  shows that  $P_\sigma^2 - P_\sigma$  is diagonal. Hence  $\sigma = id$  follows.  $\square$

Thus we have reduced the problem to orthogonally indecomposable  $\mathbb{Z}$ -lattices. In the case of root lattices (cf. [CS], Chapter 4) we obtain the well-known

**Lemma 2.** a) If  $n \geq 2$  one has

$$\text{Aut}(A_n) \cong S_{n+1} \times \{\pm 1\}.$$

b) If  $n \neq 4$  one has

$$\text{Aut}(D_n) \cong S_n \times \{\pm 1\}^n.$$

Considering our situation we obtain

**Corollary 2.** If

$$\Lambda = A_n, \quad n \neq 2 \quad \text{or} \quad \Lambda = D_n, \quad n \neq 4 \quad \text{or} \quad \Lambda = E_7$$

then  $\text{Aut}(\Lambda)$  does not contain an automorphism with minimal polynomial  $p(X) = X^2 - X + 1$ .

*Proof.* The claim is clear for  $A_1 = \mathbb{Z}$ . Suppose that  $\phi \in \text{Aut}(\Lambda)$  with minimal polynomial  $p(X) = X^2 - X + 1$  exists. Hence  $\phi^3 = -id$  follows. Considering  $A_n$ ,  $n \geq 3$ , and the action of  $S_{n+1}$  on  $A_n$  we conclude that  $A_n$  contains a sublattice of rank 3 which is  $\phi$ -invariant. As 3 is odd  $\phi$  must possess a real eigenvalue which contradicts  $p(X) = X^2 - X + 1$ .

$\text{Aut}(D_n)$ ,  $n = 1, 2$ , does not contain an element of order 3. Hence let  $n = 3$  or  $n \geq 5$  and  $\Lambda = D_n$ . Again we can conclude that  $D_n$  contains a sublattice of rank 3 which is  $\phi$ -invariant. Hence the same argument as above yields the contradiction.

In view of rank  $E_7 = 7$  we may proceed in the same way as in the last case.  $\square$

A verification yields

**Lemma 3.** The lattices  $A_2, D_4, E_6, E_8$  and the Leech lattice  $L_0$  possess an automorphism with minimal polynomial  $p(X) = X^2 - X + 1$ . This automorphism is unique up to conjugacy.

*Proof.* The existence and the uniqueness are verified by a direct computation for  $A_2$  and  $D_4$ . Considering  $E_6$ ,  $E_8$  and  $L_0$  the result follows from [CC].  $\square$

Hence we get

**Theorem 2.** *There are exactly 5 isometry classes of Eisenstein lattices of rank 12. The underlying  $\mathbb{Z}$ -lattices, the orders of the unitary automorphism groups and their index in the orthogonal automorphism groups are given by the following table*

<i>root system of <math>\Lambda</math></i>	$\# \text{Aut}(\Lambda)$	$[\text{Aut}_{\mathbb{Z}}(\Lambda) : \text{Aut}(\Lambda)]$
$3E_8$	$2^{22} \cdot 3^{16} \cdot 5^3$	$2^{21} \cdot 5^3 \cdot 7^3$
$4E_6$	$2^{16} \cdot 3^{17}$	$2^{16} \cdot 5^4$
$6D_4$	$2^{21} \cdot 3^9 \cdot 5$	$2^{19}$
$12A_2$	$2^7 \cdot 3^{15} \cdot 5 \cdot 11$	$2^{12}$
$\emptyset$	$2^{14} \cdot 3^8 \cdot 5^2 \cdot 11 \cdot 13$	$2^8 \cdot 3 \cdot 5^2 \cdot 7 \cdot 23$

*Proof.* Apply Lemma 3 and the Niemeier classification in [CS]. After conjugation we may assume that

$$B = \text{diag}(A, \dots, A), \quad A = \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix},$$

belongs to  $\text{Aut}(\Lambda)$  due to Proposition 1. Using

$$AZ^2 \cong \mathcal{O}$$

we have got an  $\mathcal{O}$ -structure on  $\Lambda$ . Moreover we have

$$\text{Aut}(\Lambda) \cong \{U \in \text{Aut}_{\mathbb{Z}}(\Lambda); UB = BU\}.$$

Now  $\# \text{Aut}(\Lambda)$  and  $[\text{Aut}_{\mathbb{Z}}(\Lambda) : \text{Aut}(\Lambda)]$  are computed explicitly using [PS]. Using Theorem 1 we conclude that we have already found all the isometry classes.  $\square$

**Remark 1.** a) Explicit examples of the lattices in Theorem 2 given by their Gram matrices can be found in [H1].

b) Theorem 1 yields

$$\mu_{16} = \frac{13 \cdot 47 \cdot 419 \cdot 691 \cdot 809 \cdot 1847 \cdot 3617 \cdot 16519}{2^{31} \cdot 3^{22} \cdot 5^4 \cdot 11 \cdot 17} \approx 0,002.$$

So far we have constructed 5 isometry classes of Eisenstein lattices  $\Lambda'_j$  of rank 16, namely the orthogonal sum from those in Theorem 2 with  $\Lambda_4$ . But they have got only a small proportion of the whole mass

$$\sum_{j=1}^5 \frac{1}{\#\text{Aut}(\Lambda'_j)} / \mu_{16} < 5 \cdot 10^{-14}.$$

c) The Eisenstein lattices associated with  $3E_8$ ,  $6D_4$  and  $L_0$  have also got the structure of a module over the Hurwitz order (cf. [BN]). The quaternionic automorphism group has the index  $3^{12}$  resp.  $3^6$  resp.  $2 \cdot 3^5 \cdot 11$  in the unitary automorphism group  $\text{Aut}(\Lambda)$ .

d) Analogous results for other imaginary quadratic number fields and  $r = 4, 8$  are achieved in [H2].

## 4 The Hermitian theta series

We consider the Hermitian half-space of degree  $n$

$$\mathcal{H}_n = \{Z \in \mathbb{C}^{n \times n}; \frac{1}{2i}(Z - \overline{Z}^{tr}) > 0\}.$$

The Hermitian modular group

$$\Gamma_n := \{M \in \text{SL}(2n; \mathcal{O}); MJ\overline{M}^{tr} = J\}, \quad J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix},$$

acts on  $\mathcal{H}_n$  by the usual fractional linear transformation. The space  $[\Gamma_n, r]$  of Hermitian modular forms of degree  $n$  and weight  $r$  consists of all holomorphic  $f : \mathcal{H}_n \rightarrow \mathbb{C}$  satisfying

$$f(M\langle Z \rangle) = \det(CZ + D)^r \cdot f(Z) \quad \text{for all } M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_n$$

with the additional condition of boundedness for  $n = 1$  (cf. [B]). The subspace  $[\Gamma_n, r]_0$  of cusp forms is defined by the kernel of the Siegel  $\phi$ -operator

$$f | \phi \equiv 0.$$



**Proposition 2.** ([CR]) *Let  $\Lambda$  be an Eisenstein lattice of rank  $r$  with Gram matrix  $H$ . Then the associated Hermitian theta series*

$$\begin{aligned}\Theta^{(n)}(Z, H) &:= \sum_{G \in \mathcal{O}^{r \times n}} e^{\pi i \operatorname{trace}(\overline{G}^{tr} H G \cdot Z)}, \quad Z \in \mathcal{H}_n, \\ &= \sum_{T \geq 0} \sharp(H, T) e^{\pi i \operatorname{trace}(TZ)},\end{aligned}$$

with the Fourier coefficients

$$\sharp(H, T) := \sharp\{G \in \mathcal{O}^{r \times n}; \overline{G}^{tr} H G = T\},$$

belongs to

$$[\Gamma_n, r].$$

Just as in the case of Siegel modular forms we obtain the analytic version of Siegel's main theorem involving the Siegel Eisenstein series in  $[\Gamma_n, r]$  (cf. [B], [K]).

**Corollary 3.** *Let  $H_1, \dots, H_s$  be Gram matrices of the lattices  $\Lambda_1, \dots, \Lambda_s$ , which are representatives of the isometry classes of Eisenstein lattices of rank  $r$ . If  $r > 2n$  one has*

$$\frac{1}{\mu_r} \sum_{j=1}^s \frac{1}{\sharp \operatorname{Aut}(\Lambda_j)} \Theta^{(n)}(Z, H_j) = E_r^{(n)}(Z) = \sum_{\begin{pmatrix} A & B \\ C & D \end{pmatrix} : \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \setminus \Gamma_n} \det(CZ + D)^{-r}.$$

If  $r = 4$  or  $r = 8$  the Eisenstein series therefore coincides with a Hermitian theta series. Let  $H_1, \dots, H_5$  be fixed Gram matrices of the Eisenstein lattices associated with  $3E_8, 4E_6, 6D_4, 12A_2$  resp.  $L_0$ . Then we can describe the filtration analogous to [NV].

**Theorem 3.** a) *Let*

$$\begin{aligned}F^{(n)}(Z) &:= \Theta^{(n)}(Z, H_1) - 30\Theta^{(n)}(Z, H_2) + 135\Theta^{(n)}(Z, H_3) - 160\Theta^{(n)}(Z, H_4) \\ &\quad + 54\Theta^{(n)}(Z, H_5) \in [\Gamma_n, 12].\end{aligned}$$

*Then  $F^{(4)} \not\equiv 0$  and either  $F^{(3)} \equiv 0$  or  $F^{(3)}$  is a non-trivial cusp form.*

b) *The form*

$$G^{(3)}(Z) = \Theta^{(3)}(Z, H_2) - 6\Theta^{(3)}(Z, H_3) + 8\Theta^{(3)}(Z, H_4) - 3\Theta^{(3)}(Z, H_5)$$

is a non-trivial cusp form of degree 3.

c) The form

$$H^{(2)}(Z) = \Theta^{(2)}(Z, H_3) - 2\Theta^{(2)}(Z, H_4) + \Theta^{(2)}(Z, H_5)$$

is a non-trivial cusp form of degree 2.

d) The form

$$J^{(1)}(Z) = \Theta^{(1)}(Z, H_4) - \Theta^{(1)}(Z, H_5)$$

is a non-trivial cusp form of degree 1.

*Proof.* The functions above are modular forms of weight 12 due to Proposition 2.

a) The modular form  $F^{(4)}$  does not vanish identically, since this is already true for the restriction to the Siegel half-space, where  $F^{(4)}$  is the analogous linear combination of the Siegel theta series. In this case we may use the table of Fourier coefficients in [BFW].

We know from [DK] that

$$[\Gamma_2, 12] = \mathbb{C}E_4^{(2)3} + \mathbb{C}E_6^{(2)2} + \mathbb{C}E_{12}^{(2)}.$$

Calculating a few Fourier coefficients we obtain  $F^{(2)} \equiv 0$ .

b) The analogous computations show that  $G^{(2)} \equiv 0$ . If  $n = 3$  the Fourier coefficient of

$$\begin{pmatrix} 4 & 4\omega/3 & 1 + \omega/3 \\ 4\bar{\omega}/3 & 4 & -2\omega/3 \\ 1 + \bar{\omega}/3 & -2\bar{\omega}/3 & 4 \end{pmatrix}$$

is 5184, hence  $G^{(3)} \not\equiv 0$ .

c) We have

$$H^{(2)} = c \cdot \left( E_{12}^{(2)} - \frac{441}{691} \left( E_4^{(2)} \right)^3 - \frac{250}{691} \left( E_6^{(2)} \right)^2 \right) \in [\Gamma_2, 12]_0, \quad c \neq 0.$$

d) This claim is clear as we deal with elliptic modular forms and elliptic theta series, hence

$$J^{(1)} = 720\Delta,$$

where  $\Delta$  is the normalized discriminant. □

**Remark 2.** a) Just as in the case of quaternionic modular forms (cf. [HK]) there is no analog of the Schottky form over  $K$ .

b) A computation of thousands of Fourier coefficients leads to the conjecture that  $F^{(3)} \equiv 0$ . In this case the space of cusp forms of weight 12 spanned by Hermitian theta series is one dimensional in the cases of degree  $n = 0, 1, 2, 3, 4$ . One knows that

$$[\Gamma_4, 12]_0 \neq \{0\}$$

from [I]. Hence it is conjectured that  $F^{(4)}$  is a Hermitian Ikeda lift.

c) All the theta series associated with Eisenstein lattices of rank  $\leq 12$  are symmetric Hermitian modular forms, i.e.  $f(Z^{tr}) = f(Z)$ , because the lattice  $\overline{\Lambda}$  is isometric to  $\Lambda$ .

## References

- [BN] C. Bachoc and G. Nebe: Classification of Two Genera of 32-dimensional Lattices of rank 8 over the Hurwitz Order. *Exp. Math.* **6**, 151-162 (1997).
- [BFW] R. E. Borcherds, E. Freitag and R. Weissauer: A Siegel cusp form of degree 12 and weight 12. *J. Reine Angew. Math.* **494**, 141-153 (1998).
- [B] H. Braun: Hermitian modular functions III. *Ann. Math.* **53**, 143-160 (1951).
- [CR] D. M. Cohen and H. L. Resnikoff: Hermitian quadratic forms and Hermitian modular forms. *Pacific J. Math.* **76**, 329-337 (1978).
- [CC] J. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson: Atlas of finite groups. Clarendon Press, Oxford 1985.
- [CS] J. Conway and N. J. A. Sloane: Sphere Packings, Lattices and Groups, 3rd ed. Springer-Verlag, New York 1999.
- [DK] T. Dern and A. Krieg: Graded rings of Hermitian modular forms of degree 2. *manuscr. math.* **110**, 251-272 (2003).

- [F] W. Feit: Some Lattices over  $\mathbb{Q}(\sqrt{-3})$ . *J. Algebra* **52**, 248-263 (1978).
- [HaK] K. I. Hashimoto and H. Koseki: Class numbers of definite unimodular hermitian forms over the rings of imaginary quadratic number fields. *Tohoku Math. J.* **41**, 1-30 (1989).
- [H1] M. Hentschel: The Eisenstein Lattices of rank 12 over  $\mathbb{Q}(\sqrt{-3})$ . <http://www.mathA.rwth-aachen.de/de/mitarbeiter/hentschel/>
- [H2] M. Hentschel: On Hermitian theta series and modular forms. PhD thesis, RWTH Aachen 2009.
- [HK] M. Hentschel and A. Krieg: A Hermitian analog of the Schottky form. In : S. Böcherer, T. Ibukiyama, M. Kaneko and F. Sato (ed.): Automorphic Forms and Zeta Functions. Proceedings of the Conference 'In Memory of Tsuneo Arakawa'. World Scientific, New Jersey, 140-159 (2005).
- [I] T. Ikeda: On the lifting of Hermitian modular forms. *Compos. Math.* **144**, 1107-1154 (2008).
- [KM] M. Kitazume and A. Munemasa: Even Unimodular Gaussian Lattices of Rank 12. *J. Number Theory* **95**, 77-94 (2002).
- [Kn] M. Kneser: Quadratische Formen. Springer-Verlag, Berlin-Heidelberg-New York 2002.
- [K] A. Krieg: Modular Forms on the Half-Spaces of Quaternions. Lect. Notes Math. **1143**, Springer-Verlag, Berlin-Heidelberg-New York 1985.
- [NV] G. Nebe and B. Venkov: On Siegel modular forms of weight 12. *J. Reine Angew. Math.* **531**, 49-60 (2001).
- [N] M. Newman: Integral Matrices. Academic Press, New York-London 1972.
- [PS] W. Plesken and B. Souvignier: Computing isometries of lattices. *J. Symb. Comput.* **24**, 327-334 (1997).
- [Q] H.-G. Quebbemann: An application of Siegel's formula over quaternion orders. *Mathematica* **31**, 12-16 (1984).

- [S] A. Schiemann: Classification of Hermitian forms with the neighbour method. *J. Symb. Comput.* **26**, 487-508 (1998).  
<http://www.math.uni-sb.de/ag/schulze/Hermitian-lattices/>
- [T] D. E. Taylor: The Geometry of Classical Groups. Heldermann Verlag, Berlin 1992.
- [Z] D. B. Zagier: Zetafunktionen und quadratische Zahlkörper. Springer-Verlag, Berlin-Heidelberg-New York 1981.