

ASYMPTOTICS OF PARTIAL DENSITY FUNCTIONS FOR DIVISORS

JULIUS ROSS AND MICHAEL SINGER

ABSTRACT. We study the asymptotic behaviour of the partial density function associated to sections of a positive hermitian line bundle that vanish to a particular order along a fixed divisor Y . Assuming the data in question is invariant under an S^1 -action (locally around Y) we prove that this density function has a distributional asymptotic expansion that is in fact smooth upon passing to a suitable real blow-up. Moreover we recover the existence of the “forbidden region” R on which the density function is exponentially small, and prove that it has an “error-function” behaviour across the boundary ∂R . As an illustrative application, we use this to study a certain natural function that can be associated to a divisor in a Kähler manifold.

1. INTRODUCTION

For motivation, consider as a toy example the space of polynomials on \mathbb{C}^n with inner product

$$(p, q) := \int_{\mathbb{C}^n} p(z)\overline{q(z)}e^{-k|z|^2}d\lambda,$$

where $k \in \mathbb{N}$ and $d\lambda$ is $(2\pi)^{-1}$ times the Lebesgue measure. For given $\varepsilon > 0$, let $\{p_\alpha\}$ be an orthonormal basis of polynomials that vanish to order at least εk along the hyperplane $\{z_1 = 0\} \subset \mathbb{C}^n$. Then the *partial density function* associated to these data is the smooth function

$$\rho_k^\varepsilon(z) := \sum_\alpha |p_\alpha(z)|^2 e^{-k|z|^2}.$$

This is independent of choice of orthonormal basis, and our interest lies in its asymptotic behaviour as k tends to infinity. By radial symmetry, $\rho_k^\varepsilon = \rho_k^\varepsilon(x)$ where $x = |z_1|^2$, and we may take the p_α to be monomial. Then, after elementary calculation,

$$k^{-n}\rho_k^\varepsilon(x) = e^{-kx} \sum_{j \geq \varepsilon k} \frac{(kx)^j}{j!}, \quad (1.1)$$

which asymptotically behaves as

$$k^{-n}\rho_k^\varepsilon(x) \sim \frac{1}{\sqrt{2\pi x}} \int_{-\infty}^{\sqrt{k}(x-\varepsilon)} e^{-\frac{t^2}{2x}} dt \quad \text{for } k \gg 0. \quad (1.2)$$

This can be seen, for instance, through the Central Limit Theorem applied to k independent Poisson random variables with parameter x . Thus $k^{-n}\rho_k^\varepsilon$ is asymptotically a standard error-function centred at $x = \varepsilon$; in particular it tends to zero exponentially fast on the set $R = \{x < \varepsilon\}$, and tends to 1 on $\{x > \varepsilon\}$ as k tends to infinity.

In this paper we study the analogous partial density function associated to sections of high powers of a positive hermitian line bundle that vanish to a particular order along a fixed divisor. Similar density functions have found a wide range of uses, including the study of random matrices (e.g. Shiffman–Zelditch [27], Berman [2]), in Kähler geometry (e.g. Pokorný–Singer [21], Ross–Witt–Nyström [24]) and in dimension $n = 1$ is closely related to the Laplacian growth (e.g. Hedenmalm–Makarov [12, 13]).

In the work of Shiffman–Zelditch [27] it is shown, essentially in the toric case, that there is a subset in which the partial density function is exponentially small (and it is here that this is given the name “forbidden region”). Through work of Berman [1] it is known (at least when the base is compact) that there is again an open forbidden region R containing the divisor such

that asymptotically the partial density function is exponentially small on compact subsets of R and is equal to the usual density function on compact subsets of the complement of \overline{R} . This has been studied again in detail in the toric case [21] but other than this rather little is known about the behaviour of the partial density function near the boundary of R .

Our results below give an essentially complete description of the partial density function when all the data in question are invariant under a local holomorphic S^1 -action and ε is sufficiently small. Roughly speaking it states that there is a natural way in which the partial density function has a globally defined asymptotic expansion in powers of $k^{1/2}$ whose terms depend on the curvature of the hermitian metric, all of which are (in principle) computable. Moreover, by working out the leading term we recover the existence of this forbidden region and show that the partial density function has the same error-function behaviour across its boundary as it does in the model case (1.2).

Before stating precise theorems we return once more to the toy example above. A convenient way to express the existence of an asymptotic expansion is through the semi-classical variable $\hbar = k^{-1/2}$. Then to say that a function of the form $k^{-n}\rho_k^\varepsilon(x)$ for $x \in \mathbb{R}_{>0}$ admits a smooth asymptotic expansion in powers of $k^{1/2}$ is to say that the function

$$\hat{\rho}(\hbar, x) := k^{-n}\rho_k^\varepsilon(x) \text{ for } \hbar = \frac{1}{\sqrt{k}} \text{ and } k \in \mathbb{N}$$

extends to a smooth function on $\mathbb{R}_{\geq 0} \times \mathbb{R}_{>0}$, i.e. it extends to a smooth function right up to the boundary on which $\hbar = 0$. Now if ρ_k^ε is the partial density function in (1.1) then $\hat{\rho}$ cannot extend smoothly to the entire boundary $\{\hbar = 0\}$. For as we have seen, its leading order term in \hbar is

$$\frac{1}{\sqrt{2\pi x}} \int_{-\infty}^{\frac{x-\varepsilon}{\hbar}} e^{-\frac{t^2}{2x}} dt$$

which does not extend smoothly over the point $(x, \hbar) = (\varepsilon, 0)$ due to the presence of the term $\xi := \frac{x-\varepsilon}{\hbar}$. However, we can formally circumvent this by considering $\hat{\rho}$ instead as a function of ξ and \hbar , at which point its smoothness is immediate. This is made precise by an approach advocated by Melrose: we consider instead the lift of $\hat{\rho}$ to the real blow-up of $\mathbb{R}_{\geq 0} \times \mathbb{R}_{>0}$ at the point $(0, \varepsilon)$ which does extend to a smooth function across the boundary.

With this in mind we state our main results. Let X be a compact complex manifold and L be a holomorphic line bundle with a positive smooth hermitian metric h . These induce an L^2 -inner product on the space of sections $H^0(L^k)$ and if Y is a smooth divisor in X then for $\varepsilon \geq 0$ the partial density function is defined to be

$$\rho_k^\varepsilon = \sum_{\alpha} |s_{\alpha,k}|_{h^k}^2$$

where $\{s_{\alpha,k}\}$ is an L^2 -orthonormal basis for the space of holomorphic sections of L^k that vanish to order at least εk along Y . When $\varepsilon = 0$ we denote this simply by ρ_k which is the usual density function (often called the Bergman function). We suppose that there is a neighbourhood U of Y that admits a holomorphic S^1 -action on X which is standard in local coordinates around Y . That is,

$$e^{i\theta} \cdot (z, w) = (e^{i\theta} z, w) \text{ for } e^{i\theta} \in S^1 \tag{1.3}$$

where Y is locally defined by $z = 0$. We also assume that the restrictions of all the initial data to U (i.e. the line bundle and metric) are invariant under this action. Denote by $\mu: X \rightarrow \mathbb{R}$ the Hamiltonian of the action, normalized so that $\mu^{-1}(0) = Y$, and let v be the vector field generating the S^1 -action.

Theorem 1.1. *Given the above data, for sufficiently small ε , we have*

$$\rho_k^\varepsilon \sim \begin{cases} O(k^{-\infty}) & \text{on } \mu^{-1}[0, \varepsilon), \\ \rho_k + O(k^{-\infty}) & \text{on } X \setminus \mu^{-1}[0, \varepsilon]. \end{cases} \tag{1.4}$$

By this statement we mean that equality holds on any given compact subset of $\mu^{-1}[0, \varepsilon)$ (respectively $X \setminus \mu^{-1}[0, \varepsilon]$). Thus $\mu^{-1}[0, \varepsilon)$ is precisely the forbidden region described above.

Theorem 1.2. Let ρ_k^ε be the partial density function and set

$$\hat{\rho}(\hbar, x) = k^{-n} \rho_k^\varepsilon \text{ for } \hbar = k^{-1/2}.$$

Then $\hat{\rho}$ has a distributional asymptotic expansion on X . In fact this distribution is the push-forward by the blow-down map β of a smooth function on the real blow-up of $X \times [0, \infty)$ along $\mu^{-1}(\varepsilon) \times \{0\}$. Its leading order term is given by


$$\hat{\rho}(z, \hbar) = \frac{1}{\sqrt{2\pi|v(z)|^2}} \int_{-\infty}^{\frac{\mu(z)-\varepsilon}{\hbar}} e^{-\frac{t^2}{2|v(z)|^2}} dt + O(\hbar) \quad (1.5)$$

for (z, \hbar) such that

$$\frac{\mu(z) - \varepsilon}{\hbar}$$

is bounded.

We refer the reader Appendix A for a summary of this real-blowup that is denoted $[X \times [0, \infty); \mu^{-1}(\varepsilon) \times \{0\}]$. Roughly speaking it is obtained by replacing the submanifold $\mu^{-1}(\varepsilon)$ with a “half-cylinder” as in the following picture, and allows the use of well-defined “polar coordinates” centered at points in $\mu^{-1}(\varepsilon)$.



So using again the substitution $\hbar = 1/\sqrt{k}$ we can interpret this as the statement that $\rho_{1/\hbar^2}^\varepsilon$ is a smooth function on the half-space $X \times [0, \infty)$ (or really the restriction of such a smooth function for values of \hbar with \hbar^{-2} a positive integer). In fact, there is almost certainly an interpretation of density functions for all real powers of L using Melrose’s ideas [18], which would give a natural interpretation of the (partial) density function for all values of the semi-classical parameter \hbar .

The idea of the proof is as follows. Consider first the case that X is one dimensional. Then looking locally we are essentially interested in the case X is the unit disc with coordinate z and Y is the origin. Since all the data are S^1 -invariant, the set of monomials $z^{\varepsilon k}, z^{\varepsilon k+1}, \dots$ give an orthogonal set of local sections that vanish to the right order along Y . Thus the partial density function can be calculated by summing the pointwise norm of these functions, once they have been normalised to have unit L^2 -length. Using the moment-variable (given by a Legendre transform of the potential defining the metric) this can be done with a combination of standard techniques, namely Laplace’s method to calculate the integrals defining the length of these functions, and then a combination of Laplace’s method and the Euler-Maclaurin formula to expand the resulting sum in powers of k . This gives a local formula for a quantity that ought to be the partial density function, and one argues that this is in fact the case up to insignificant terms.

Our real interest is the case of higher dimension. Here we argue similarly, replacing the powers of z with powers of a choice of defining section for Y . We then use a parameterized version of the Legendre transform to deal with the directions normal to Y . In fact, we interpret this

Legendre transform as giving a family of hermitian metrics on certain line bundles on Y which, along with a family of volume forms that we construct, gives a family of density functions on Y (that play the role of the normalising of the sections z^j to have unit length). Together these can be summed to give a quantity that ought to be the partial density function, and again one proves this is the case up to insignificant terms. Then, using essentially the same techniques as in the one-dimensional case, we prove that it has the desired asymptotic expansion in powers of k . We refer the reader to Section 5 for a more detailed summary.

Acknowledgements: During this work JR was supported by an EPSRC Career Acceleration Fellowship (EP/J002062/1) and MS by the Leverhulme Trust. The authors wish to thank Bo Berndtsson, David Witt Nyström, Håkan Hedenmalm and Steve Zelditch for helpful conversations.

2. PARTIAL BERGMAN KERNELS

The density function of a hermitian line bundle is the restriction to the diagonal of the Bergman kernel, which is the reproducing kernel for the L^2 -projection to the space of holomorphic sections. Here we shall discuss the simple extension of this concept in which we impose a certain vanishing of the sections along a fixed submanifold.

2.1. Definition of Partial Bergman Kernels. We start by recalling some standard notation and terminology. Let L be a holomorphic line bundle on a complex manifold X of dimension n , and h be a hermitian metric on L . Given a local holomorphic trivialization ζ of L we can write $|\zeta(z)|_h^2 = e^{-2\varphi_\zeta(z)}$ for some smooth potential function φ_ζ . By standard abuse of notation we let φ denote this potential (even though it is not globally defined) and we confuse φ with the metric $h = e^{-\varphi}$. Thus $h^k = e^{-k\varphi}$ is the induced hermitian metric on $L^k := L^{\otimes k}$ for $k \in \mathbb{N}$. In terms of transition functions, if $\zeta_\alpha = \lambda_{\alpha\beta}\zeta_\beta$ then $\varphi_{\zeta_\beta} = \varphi_{\zeta_\alpha} + \log|\lambda_{\alpha\beta}|$. From this one can extend the notion of a potential to include the case of \mathbb{Q} -line bundles.

If s is a section of L then by abuse of notation we let s also denote its local representative in given a trivialization: thus $|s(z)|_\varphi = |s(z)|e^{-\varphi(z)}$. The associated curvature form¹

$$\omega_\varphi = dd^c\varphi = \frac{1}{2\pi}dJd\varphi = \frac{i}{\pi}\partial\bar{\partial}\varphi \quad (2.1)$$

is a well defined $(1, 1)$ -form and the hermitian metric $e^{-\varphi}$ is said to be (strictly) positive if ω_φ is a (strictly) positive form, which occurs if and only if φ_ζ is a (strictly) plurisubharmonic function for all local trivializations ζ . When φ is strictly plurisubharmonic we let $\omega_\varphi^{[n]} = \frac{1}{n!}\omega_\varphi^n$ be the associated volume form.

The abuse of notation $h = e^{-\varphi}$ also leads us to use the terminology ‘ φ is a (plurisubharmonic potential) for L ’ instead of h is a metric on L (with positive curvature).

The hermitian metric $e^{-\varphi}$ gives a pairing $(\cdot, \cdot)_\varphi: L_z \otimes L_z \rightarrow \mathbb{C}$ that is linear in the first variable and conjugate linear in the second. The induced L^2 -inner product on two smooth sections s, t of L is given by

$$\langle s, t \rangle_\varphi := \int_X (s, t)_\varphi \omega_\varphi^{[n]} = \int_X s(z)\bar{t}(z)e^{-2\varphi(z)}\omega_\varphi^{[n]}$$

and we shall write $\|s\|_\varphi$ for the corresponding L^2 -norm. We denote by $L_{k\varphi}^2 := L_{k\varphi}^2(X)$ the space of sections of L^k with finite L^2 -norm, and let $H_{k\varphi} := H_{k\varphi}(X) \subset L_{k\varphi}^2$ denote the subspace of holomorphic sections.

We write \bar{L} for the bundle L with the conjugate complex structure, so if s is a (local) holomorphic section of L then \bar{s} is a (local) holomorphic section of \bar{L} . If V, W are auxiliary holomorphic vector bundles on X then we let $(\cdot, \cdot)_\varphi$ also denote the naturally induced pairing $(L \otimes V) \otimes (L \otimes W) \rightarrow V \otimes \bar{W}$ and extend our other notations accordingly.

¹Note the factor of 2π

Assume now that φ is a smooth strictly plurisubharmonic potential on L . Since we will be interested in sections of L^k asymptotically as k tends to infinity we make this part of our definition of the partial Bergman kernel as follows. Let $Y \subset X$ be a compact complex submanifold and $\varepsilon \in \mathbb{Q}_{\geq 0}$.

Definition 2.1. Denote by $H_{k\varphi}^{\varepsilon k}$ the subspace of holomorphic sections of L^k vanishing to order at least εk along Y .

Definition 2.2. The *partial Bergman kernel* (PBK) for (ε, Y) is a sequence of sections K_k^ε of $\overline{L}^k \boxtimes L^k$ on $X \times X$ for $k, \varepsilon k \in \mathbb{N}$ such that

- (1) For each fixed z , the section $z' \mapsto K_{k,z}^\varepsilon(z') := K_k^\varepsilon(z, z')$ is in $\overline{L}_z^k \boxtimes H_{k\varphi}^{\varepsilon k}$ and
- (2) We have

$$s(z) = \langle s, K_{k,z}^\varepsilon \rangle_{k\varphi} = \int_X s(z') \overline{K_k^\varepsilon(z, z')} e^{-2k\varphi(z')} \omega_{\varphi, z'}^{[n]} \text{ for all } z \in X \text{ and } s \in H_{k\varphi}^{\varepsilon k} \quad (2.2)$$

where the notation $\omega_{\varphi, z'}^{[n]}$ indicates the integral is being taken with respect to the variable z' .

Definition 2.3. The *partial density function* (PDF) ρ_k^ε is the norm of the restriction of the PBK to the diagonal, i.e.

$$\rho_k^\varepsilon(z) := |K_k^\varepsilon(z, z)|_{k\varphi} = K_k^\varepsilon(z, z) e^{-2k\varphi(z)} \quad (2.3)$$

which is a smooth real valued function on X .

Of course if $\varepsilon = 0$ then this definition recovers what is commonly referred to as the Bergman kernel and density function for L^k and we write these simply as K_k and ρ_k . We remark that all of these definitions can equally be made with $\omega_\varphi^{[n]}$ replaced by a given smooth volume form dV on X .

2.2. Existence and Basic Properties. The existence of the PBK follows from standard functional analysis considerations. Suppose that X is either a compact Kähler manifold, or else a bounded domain, with smooth boundary, in a Kähler manifold.

As is well known, $H_{k\varphi}^{\varepsilon k}$ is a closed subspace of $L_{k\varphi}^2$, so in particular it is a Hilbert space. Moreover, if $z \in X$, the evaluation map $s \mapsto s(z)$ is a bounded linear functional $H_{k\varphi}^{\varepsilon k} \rightarrow L_z^k$. So, by the Riesz representation theorem, there is an element

$$K_{k,z}^\varepsilon \in \overline{L}_z^k \otimes H_{k\varphi}^{\varepsilon k} \text{ with } f(z) = \langle f, K_{k,z}^\varepsilon \rangle_{k\varphi} \text{ for all } f \in H_{k\varphi}^{\varepsilon k}. \quad (2.4)$$

Clearly $\langle u, K_{k,z}^\varepsilon \rangle_{k\varphi} = 0$ if $u \in L_{k\varphi}^2$ is orthogonal to $H_{k\varphi}^{\varepsilon k}$, so the kernel

$$K_k^\varepsilon(z, z') := K_{k,z}^\varepsilon(z') \quad (2.5)$$

is the Schwartz kernel of the orthogonal projection $\mathcal{K}_k^\varepsilon : L_{k\varphi}^2 \rightarrow H_{k\varphi}^{\varepsilon k}$ in the sense that

$$(\mathcal{K}_k^\varepsilon u)(z) = \langle u, K_{k,z}^\varepsilon \rangle_{k\varphi} = \int_X u(z') \overline{K_k^\varepsilon(z, z')} e^{-2k\varphi(z')} \omega_{z'}^{[n]} \text{ for } u \in L_{k\varphi}^2. \quad (2.6)$$

In particular this gives uniqueness of K_k^ε . Because $\mathcal{K}_k^\varepsilon$ is orthogonal, it is self-adjoint, and so (2.5) is hermitian, i.e.

$$\overline{K_{k,z'}^\varepsilon(z)} = K_{k,z}^\varepsilon(z'). \quad (2.7)$$

Thus we can also write (2.6) as

$$\mathcal{K}_k^\varepsilon u(z) = \int_X u(z') K_k^\varepsilon(z', z) e^{-2k\varphi(z')} \omega_{z'}^{[n]}. \quad (2.8)$$

In particular

$$\overline{\partial_{z'} K_{k,z}^\varepsilon(z')} = 0 \text{ and } \overline{\partial_z K_{k,z}^\varepsilon(z')} = 0, \quad (2.9)$$

so K_k^ε is holomorphic as a section of $\overline{L}^k \boxtimes L^k$.

Remark 2.4. If X is a compact Kähler manifold the spaces $H_{k\varphi}^{\varepsilon k}$ are finite-dimensional and the definition above agrees with that in the introduction. For if $s_{\alpha,k}$ is an L^2 -orthonormal basis for $H_{k\varphi}^{\varepsilon k} = H^0(L^k \otimes \mathcal{I}_Y^{\varepsilon k})$ then one easily checks that

$$K_k^\varepsilon(z, z') = \sum_\alpha \overline{s_{\alpha,k}(z)} \boxtimes s_{\alpha,k}(z') \quad (2.10)$$

has the characteristic properties of the PBK, so

$$\rho_k^\varepsilon = \sum_\alpha |s_{\alpha,k}|_{k\varphi}^2. \quad (2.11)$$

Remark 2.5. When $\varepsilon = 0$ the difference $v := u - \mathcal{K}_k[u]$ is the $L_{k\varphi}^2$ -minimal solution of the equation $\bar{\partial}v = \bar{\partial}u$. This follows directly from the orthogonality of the projection.

2.3. Bounds for Bergman Kernels. For convenience we recall here some basic estimates related to Bergman kernels. The fundamental ‘ L^2 implies L^∞ bound’ for holomorphic functions (Proposition 2.6 below) allows us to estimate the norm of the evaluation map and hence by the discussion in §2.2, the norms of the PBK and PDF themselves. These estimates will be refined using *extremal envelopes* in the next section.

On a Kähler manifold X we denote by $B_z(\delta)$ the geodesic ball of radius δ centred at z (taken with respect to the given Kähler metric). As usual L will be a holomorphic line bundle on X with smooth strictly plurisubharmonic potential φ .

Proposition 2.6. *Let X be a complex manifold and $W \subset X$ be relatively compact. Then there is a constant C_W such that for all k sufficiently large*

$$|f(z)|_{k\varphi} \leq C_W k^{n/2} \int_{B_z(k^{-1/2})} |f|_{k\varphi}^2 \omega^{[n]} \leq C_W k^{n/2} \|f\|_{k\varphi} \quad (2.12)$$

for all $f \in H_{k\varphi}$ and all $z \in W$.

Proof. For $z \in W$ we may choose local coordinates on a chart so that $z = 0$, and by a change of gauge that $\varphi(z) = O(|z|^2)$. In this chart the geodesic metric $\rho(\cdot, \cdot)$ is equivalent to the Euclidean metric, so there is a constant $c > 0$ such that $c|z - z'| \leq \rho(z, z') \leq c^{-1}|z - z'|$ for points z, z' in this chart.

Let $\chi(t): \mathbb{R} \rightarrow [0, 1]$ be a smooth non-negative cut-off function equal to 1 for $t \leq 1/2$ and equal to 0 for $t \geq 1$ and set

$$\chi_k(t) = \chi(c^{-1}k^{1/2}t)$$

We recall the *Bochner–Martinelli–Koppelman formula* [22, Ch. IV]

$$u(z) = - \int_{\mathbb{C}^n} \bar{\partial}u(w) \wedge B(w, z), \quad u \in C_0^\infty(\mathbb{C}^n) \quad (2.13)$$

where the Bochner–Martinelli kernel is given by the formula

$$B(z, w) := c_n \frac{1}{|w - z|^{2n}} \eta(w, z) \quad (2.14)$$

where c_n is a universal constant and

$$\eta(w, z) = \iota_{(\bar{w}-\bar{z})\bar{\partial}_w} d\bar{w}_1 \wedge \cdots \wedge d\bar{w}_n \wedge dw_1 \wedge \cdots \wedge dw_n. \quad (2.15)$$

We apply this with $u = \chi_k(|z|)f(z)$. Then $\bar{\partial}u = f\bar{\partial}\chi_k$ is supported in a spherical shell of radius $O(k^{-1/2})$ and $\bar{\partial}\chi_k = O(k^{1/2})$. So

$$|\bar{\partial}(\chi_k)B(w, 0)| = O(k^n). \quad (2.16)$$

Hence (2.13) gives

$$f(0) = - \int_{|w| \leq ck^{-1/2}} f(w) \bar{\partial}\chi_k B(0, z) \quad (2.17)$$

where k is to be taken large enough so the support of χ_k lies in this chart. We are assuming $\varphi(0) = 0$, so $|f(0)|^2 = |f(0)|_{k\varphi}^2$. Thus using the Cauchy-Schwarz inequality,

$$|f(0)|_{k\varphi}^2 \leq C \int_{|w| \leq ck^{-1/2}} |f(w)|_{k\varphi}^2 \omega^{[n]} \int e^{k\varphi} |\bar{\partial}\chi_\delta B(w, 0)|^2, \quad (2.18)$$

as $\omega^{[n]}$ is smooth so is equivalent to the euclidean volume form. Now in the first integral we may replace the Euclidean ball $\{|w| < ck^{-1/2}\}$ by the geodesic ball $B_0(k^{-1/2})$ and only improve the inequality. On the other hand, $k\varphi = O(k|z|^2)$ on this chart so $e^{k\varphi}$ is uniformly bounded, and the set in which χ_k is supported has volume $O(k^{-n})$. Thus (2.16) implies the second integral is of order $O(k^n)$. This proves the result for a single point, and the uniform estimate follows an obvious covering argument. \square

Recall that $K_{k,z}^\varepsilon \in \bar{L}_z^k \otimes H_{k\varphi}^{\varepsilon k}$ so with our convention of using the same notation for a section of a line bundle and its representation in a local frame we have

$$\|K_{k,z}^\varepsilon\|_{k\varphi}^2 = \int_X |K_{k,z}^\varepsilon(w)|^2 e^{-2k(\varphi(w)+\varphi(z))} \omega_w^{[n]}.$$

Corollary 2.7. *Suppose X is compact. Then there is a constant C such that*

$$\|K_{k,z}^\varepsilon\|_{k\varphi} \leq Ck^{n/2} \quad (2.19)$$

for all $z \in X$ and all $\varepsilon \geq 0$.

Proof. Essentially by definition, $\|K_{k,z}^\varepsilon\|_{k\varphi}$ is equal to the operator norm of the evaluation map $H_{k\varphi}^{\varepsilon k} \rightarrow L_z^k$ given by $s \mapsto s(z)$. But the previous theorem says precisely that this operator norm is bounded by $Ck^{n/2}$. \square

Corollary 2.8. *Suppose X is compact. Then there is a constant C such that*

$$|K_k^\varepsilon(z, z')|_{k\varphi} \leq Ck^n \text{ for all } z', z \in X. \quad (2.20)$$

Proof. Apply the above to the function $y \mapsto K_{k,z}(y)$ itself to deduce that $|K_{k,z}(y)|_{k\varphi}$ is bounded by $O(k^{n/2})\|K_{k,z}^\varepsilon\|_{k\varphi} = O(k^n)$. \square

2.4. Decay away from the Diagonal. As further notation let $\rho(z, z')$ denote the geodesic distance between points $z, z' \in X$ taken with respect to the given Kähler metric. We wish to discuss the well known fact that the Bergman kernel decays exponentially fast away from the diagonal (see, for instance, [16, Proposition 9], [6, Prop 4.1], [8], [19, Thm 0.1]).

Theorem 2.9 (Decay away from the diagonal). *Suppose X is compact. Then K_k decays exponentially fast away from the diagonal, in the following sense: there are constants $C, c > 0$ such that*

$$|K_k(x, y)|_{k\varphi} \leq Ck^n e^{-c\sqrt{k}\rho(x,y)} \text{ for all } x, y \in X.$$

For convenience of the reader we give a proof of this fact, based on a variant of the ‘‘Donnelly-Fefferman trick’’ essentially due to Bo Berndtsson [4] (and whom we thank for pointing us in this direction).

Theorem 2.10. *Let f be a $\bar{\partial}$ -closed $(0, 1)$ form with values in L and v be the L^2 -minimal solution to the equation $\bar{\partial}v = f$ and let $x \in X$. Then there are constants c, C (independent of f, k, x) such that*

$$\int_X |v(z)|_{k\varphi}^2 e^{-2c\sqrt{k}\rho(z,x)} \omega_z^{[n]} \leq \frac{C}{k} \int_X |f(z)|_{k\varphi}^2 e^{-\frac{c}{2}\sqrt{k}\rho(z,x)} \omega_z^{[n]}.$$

In particular if $f = \bar{\partial}g$ for some g then this holds for $v = g - \mathcal{H}_k(g)$.

We remark that the above statement may well be suboptimal but is sufficient for our purpose (the statement in [4, Theorem 2.1] replaces the terms $2c$ and the $\frac{c}{2}$ in the left and right hand side respectively with the same constant c when X is a domain in \mathbb{C}^n). The proof we give now is essentially the same as [4, Theorem 2.1], but replaces the Euclidean distance function on \mathbb{C}^n

with the geodesic distance function $\rho(\cdot, x)$ on X , and requires some extra care to deal with the non-smoothness $\rho(\cdot, x)$ near the cut-locus of x .

Proof. The squared distance function $\rho^2 : X \times X \rightarrow \mathbb{R}$ is continuous and can fail to be smooth only on the cut-locus

$$B := \{(z, z') \in X \times X : z \text{ and } z' \text{ are conjugate points}\}$$

of X . In particular ρ^2 is smooth in a fixed neighbourhood N of the diagonal. To deal with the non-smoothness at B , choose a non-negative function $\tilde{\rho}$ which is equal to ρ in N , smooth on $X \times X \setminus N$ and which satisfies

$$\frac{1}{2}\rho(z, z') \leq \tilde{\rho}(z, z') \leq 2\rho(z, z') \text{ for all } (z, z') \in X \times X.$$

Then $\tilde{\rho}^2$ is smooth on $X \times X$ and is equal to ρ^2 in N .

Now fix f and $x \in X$, and simplify notation by writing $\tilde{\rho}(z)$ for $\tilde{\rho}(x, z)$. Then for $k \gg 0$, in the region $\tilde{\rho} \leq \frac{1}{\sqrt{k}}$ we have $\tilde{\rho}(z) = \rho(z, x)$. It will be clear from the following argument that our constants will be uniform in x and f but we leave the reader to keep track of this.

Now for a small $c > 0$ set

$$\chi_k(z) := \begin{cases} c \left(\frac{k}{2} \tilde{\rho}(z)^2 + \frac{1}{2} \right) & \text{if } \tilde{\rho}(z) \leq \frac{1}{\sqrt{k}} \\ c\sqrt{k}\tilde{\rho}(z) & \text{otherwise.} \end{cases}$$

One checks easily that

$$\frac{c\sqrt{k}}{2}\rho(z, x) - c \leq \chi_k(z) \leq 2c\sqrt{k}\rho(z, x) + c \text{ for all } z \in X. \quad (2.21)$$

Furthermore on the region $\tilde{\rho} \leq \frac{1}{\sqrt{k}}$ we have $dd^c\chi_k = O(ck)$, whereas on the complement to this region $dd^c\chi_k = O(c\sqrt{k})$. So we may pick $c \ll 1$ small enough so that the potential

$$\zeta_k := k\varphi - \frac{1}{2}\chi_k$$

has strictly positive curvature growing at rate $O(k)$, say

$$dd^c\zeta_k \geq \frac{k}{2}dd^c\varphi. \quad (2.22)$$

Now set

$$v_k := ve^{-\chi_k}$$

Since v is orthogonal to the the space holomorphic sections with respect to the L^2 -inner product defined by $e^{-k\varphi}$, we see that v_k is orthogonal to this space with respect to the L^2 -inner product defined by $e^{-\zeta_k}$ (all of this is taken with respect to the volume form $\omega_\varphi^{[n]}$ which is fixed). Hence v_k is the L^2 -minimal solution to the equation

$$\bar{\partial}v_k = fe^{-\chi_k} - ve^{-\chi_k}\bar{\partial}\chi_k$$

with respect to the L^2 -inner product induced by $e^{-\zeta_k}$. Thus by the Hörmander estimate

$$\begin{aligned} \int_X |v|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]} &= \int_X |v_k|^2 e^{-2k\varphi + \chi_k} \omega_\varphi^{[n]} \\ &= \int_X |v_k|_{\zeta_k}^2 \omega_\varphi^{[n]} \\ &\leq O(1/k) \int_X |\bar{\partial}v_k|_{\zeta_k}^2 \omega_\varphi^{[n]} \\ &= O(1/k) \int_X |\bar{\partial}v_k|_{k\varphi}^2 e^{\chi_k} \omega_\varphi^{[n]}. \end{aligned} \quad (2.23)$$

We remark that the $O(1/k)$ term is bounded independent of c sufficiently small by (2.22).

Now the right hand side of (2.23) is bounded by

$$O(1/k) \int_X (|f|_{k\varphi}^2 e^{-\chi_k} + |v|_{k\varphi}^2 e^{-\chi_k} |\bar{\partial}\chi_k|^2) \omega_\varphi^{[n]} = \frac{A}{k} \int_X |f|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]} + Zc^2 \int_X |v|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]} \quad (2.24)$$

where A and Z are uniform constants. This follows because $\bar{\partial}\chi_k = O(c\sqrt{k})$. Inserting (2.24) into (2.23) we have

$$\int_X |v|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]} \leq \frac{A}{k} \int_X |f|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]} + Zc^2 \int_X |v|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]}. \quad (2.25)$$

Choosing c so small that $Zc^2 < 1/2$, say, we can move the second integral to the other side, giving

$$\int_X |v|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]} \leq \frac{2A}{k} \int_X |f|_{k\varphi}^2 e^{-\chi_k} \omega_\varphi^{[n]}.$$

The statement of the theorem now follows from (2.21). \square

Proof of Theorem 2.9. We know that $w \mapsto K_{k,y}(w)$ is holomorphic for fixed y , so by the proof of Proposition 2.6,

$$|K_{k,x}(y)|_{k\varphi}^2 \leq Ck^n \int_{B_\delta(y)} |K_{k,x}(z)|_{k\varphi}^2 \omega_{\varphi,z}^{[n]} \quad (2.26)$$

which is valid as long as δ is of order greater than or equal to $2/\sqrt{k}$. Observe that if $\rho(x,y) \leq \frac{1}{\sqrt{k}}$ then the bound we want follows from the bound $|K_k^{\varepsilon Y}(x,y)|_{k\varphi} = O(k^n)$ from (2.8). So we may assume that

$$\rho(x,y) \geq \delta := \frac{2}{\sqrt{k}}.$$

Fix a smooth non-negative cut-off function χ that is identically 1 on $B_{\delta/4}(y)$ supported in $B_{\delta/2}(y)$ and such that $|\bar{\partial}\chi| = O(\delta^{-1})$. Observe that $\rho(x,y) \geq \delta$ implies $\chi(x) = 0$. Then we can clearly replace (2.26) by

$$|K_{k,x}(y)|_{k\varphi}^2 = |K_{k,y}(x)|_{k\varphi}^2 \leq O(k^n) \int |K_{k,x}(z)|_{k\varphi}^2 \chi(z) \omega_{\varphi,z}^{[n]} \quad (2.27)$$

$$= O(k^n) (\chi K_{k,x}, K_{k,x})_{k\varphi} = O(k^n) \mathcal{K}_k[\chi K_{k,x}](x), \quad (2.28)$$

where, we recall, \mathcal{K}_k is the projection onto the holomorphic sections. Now set

$$f := \bar{\partial}(\chi K_{k,x}) = (\bar{\partial}\chi)K_{k,x}$$

and

$$v := \chi K_{k,x} - \mathcal{K}_k(\chi K_{k,x}).$$

Then

$$\mathcal{K}_k[\chi K_{k,x}](x) = \chi(x)K_{k,x}(x) - v(x) = -v(x) \quad (2.29)$$

as $\chi(x) = 0$. On the other hand, by Theorem 2.10, there is a $c > 0$ such that

$$\int |v(z)|_{k\varphi}^2 e^{-2c\sqrt{k}\rho(z,x)} \omega_z^{[n]} \leq O(1/k) \int |f(z)|_{k\varphi}^2 e^{-\frac{c}{2}\sqrt{k}\rho(z,x)} \omega_z^{[n]}. \quad (2.30)$$

Note that f is supported in $B_{\delta/2}(y)$ and if $z \in B_{\delta/2}(y)$ then $\rho(x,y) \leq \rho(z,x) + \frac{1}{\sqrt{k}}$. Thus

$$\int |v(z)|_{k\varphi}^2 e^{-2c\sqrt{k}\rho(z,x)} \omega_z^{[n]} \leq O(1/k) O(\delta^{-2}) \|K_{k,x}\|_{k\varphi}^2 e^{-\frac{c}{2}\sqrt{k}\rho(x,y)} \quad (2.31)$$

$$= O(k^n) e^{-\frac{c}{2}\sqrt{k}\rho(x,y)} \quad (2.32)$$

since $\|K_{k,x}\|_{k\varphi}^2 = O(k^n)$ by Corollary 2.7. On the other hand by same argument with the the Bochner–Martinelli formula again, we have

$$|v(x)|_{k\varphi}^2 \leq O(k^n) \int_{B_\delta(x)} |v|_{k\varphi}^2 \omega_\varphi^{[n]}. \quad (2.33)$$

Notice the quantity $e^{-2c\sqrt{k}\rho(z,x)}$ is bounded above and below by a constant on the ball $B_\delta(x)$, so this gives

$$\begin{aligned} |v(x)|_{k\varphi}^2 &\leq O(k^n) \int_{B_\delta(x)} |v|_{k\varphi}^2 e^{-2c\sqrt{k}\rho(z,x)} \omega_\varphi^{[n]} \\ &\leq O(k^{2n}) e^{-\frac{c}{2}\sqrt{k}\rho(x,y)}. \end{aligned}$$

Putting this with (2.27), (2.29) gives the desired estimate. \square

2.5. Extremal Envelopes. Let $Y \subset X$ be a compact complex submanifold. Following Berman [1] we make the following definition:

Definition 2.11. Given X, Y, L and $\varepsilon \geq 0$, the *extremal envelope*, φ_ε , is defined as

$$\varphi_\varepsilon := \sup\{\gamma \text{ a potential on } L : dd^c\gamma \geq 0, \gamma \leq \varphi \text{ and } \nu_Y(\gamma) \geq \varepsilon\}. \quad (2.34)$$

Here the notation means that $e^{-\gamma}$ is a possibly singular hermitian metric on L with non-negative curvature form; and $\nu_Y(\gamma)$ denotes the Lelong number of γ along Y . Since the upper semicontinuous regularisation of φ_ε lies in the set on the RHS of (2.34), it follows that φ_ε is itself semicontinuous and thus plurisubharmonic. We will always take ε to be sufficiently small so that there exists such a γ , and so φ_ε is not identically $-\infty$. Then by passing to the blowup of X along Y one can prove easily that $\nu_Y(\varphi_\varepsilon) = \varepsilon$.

Definition 2.12. Define the *forbidden region* to be the set

$$D_\varepsilon := \{z \in X : \varphi_\varepsilon(z) < \varphi(z)\},$$

and the *equilibrium set* to be the complement

$$X \setminus D_\varepsilon = \{z \in X : \varphi_\varepsilon(z) = \varphi(z)\}.$$

Then clearly D_0 is empty, and for all $\varepsilon > 0$ it is a neighbourhood of Y , and if $\varepsilon < \varepsilon'$ then $D_{\varepsilon'} \subset D_\varepsilon$. Again following [1], we now show the sets D_ε govern the PDF in that for large k the function ρ_k^ε is exponentially small on any compact subset of D_ε . The following statement refines the bounds (2.19) and (2.20), using the envelope φ_ε .

Proposition 2.13. *Assume X is compact. Then there exists a constant C such that*

$$\|K_{k,z}^\varepsilon\|_{k\varphi} \leq Ck^{n/2} e^{-k(\varphi(z) - \varphi_\varepsilon(z))} \text{ for } z \in X$$

and

$$|K_k^\varepsilon(z, z')|_{k\varphi} \leq Ck^n e^{-k(\varphi(z) - \varphi_\varepsilon(z))} e^{-k(\varphi(z') - \varphi_\varepsilon(z'))} \text{ for } z, z' \in X.$$

In particular, $K_{k,z}^\varepsilon$ is exponentially small on D_ε .

Proof. Note first that since φ and φ_ε are both potentials for L , their difference is a genuine function on X , so the right hand side of these estimates are well-defined. Let $s \in H_{k\varphi}^{\varepsilon k}$. Then by Proposition 2.6 there is a constant C such that

$$|s(w)|_{k\varphi}^2 \leq Ck^n \|s\|_{k\varphi}^2 \text{ for all } w \in X.$$

Taking the logarithm,

$$\frac{1}{2k} \ln |s(w)|^2 \leq \varphi(w) + \alpha_k \text{ for all } w \in X \quad (2.35)$$

where

$$\alpha_k := \frac{\ln(Ck^n \|s\|_{k\varphi}^2)}{2k}.$$

As s vanishes to order at least εk along Y the plurisubharmonic potential $\ln |s|$ has Lelong number at least εk along Y . Combining this with (2.35), the potential $\frac{1}{2k} \ln |s|^2 - \alpha_k$ is a candidate for the envelope defining φ_ε so

$$\frac{1}{2k} \ln |s(z)|^2 - \alpha_k \leq \varphi_\varepsilon(z) \text{ for } z \in X.$$

So multiplying by $2k$ and then exponentiating gives

$$|s(z)|_{k\varphi}^2 \leq Ck^n e^{-2k(\varphi(z) - \varphi_\varepsilon(z))} \|s\|_{k\varphi}^2 \text{ for } z \in X.$$

Thus if $z \in X$, the square of the norm of the evaluation map $H_{k\varphi}^{\varepsilon k} \rightarrow L_z$ given by $s \mapsto s(z)$ is bounded by $Ck^n e^{-k(\varphi(z) - \varphi_\varepsilon(z))}$ which gives the first statement of the Proposition. The second statement follows by applying this to the section $s := K_{k,z}^{\varepsilon Y}$ to get that for $z, z' \in X$

$$|K_{k,z}^{\varepsilon}(z')|_{k\varphi} \leq Ck^{n/2} e^{-k(\varphi(z') - \varphi_\varepsilon(z'))} \|K_{k,z}^{\varepsilon}\|_{k\varphi} \leq O(k^n) e^{-k(\varphi(z) - \varphi_\varepsilon(z))} e^{-k(\varphi(z') - \varphi_\varepsilon(z'))}.$$

□

Remark 2.14. Berman [1] proves that φ_ε is $C^{1,1}$ although we will not use that here. The continuity of φ_ε is enough to imply that for large k , $K_{k,z}^{\varepsilon k}$ is exponentially small on any given compact subset of D_ε . Berman also proves that ρ_k^ε is asymptotically close to ρ_k on any given compact subset of $X \setminus \overline{D_\varepsilon}$. Thus the interest lies in the behaviour across the boundary of D_ε which will be the study of the rest of this paper.

3. LOCAL BERGMAN KERNELS

We next adapt a key idea of Berman–Berndtsson–Sjöstrand [3] who introduced the concept of a local Bergman kernel which has the reproducing property up to a term that is negligible for large k . Our aim is to give a suitable definition of a local PBK that is defined on a neighbourhood of Y .

Again suppose that X is a complex manifold of dimension n and L a holomorphic line bundle with given strictly positive hermitian metric $e^{-\varphi}$. Also fix an open $U \subset X$ containing Y and a smooth positive function χ that has compact support in U and is identically equal to 1 on some open subset $W \subset U$ containing Y .

Definition 3.1. We say a sequence of holomorphic sections B_k^ε of $L|_U \boxtimes \overline{L}|_U$ over $U \times U$ is a *local partial Bergman kernel* (local PBK) on W of order $N \geq 0$ if

(1) (Almost Reproducing) We have

$$\left| f(z) - \int_U \chi(z') f(z') \overline{B}_k^\varepsilon(z, z') e^{-2k\varphi(z')} \omega_{\varphi, z'}^{[n]} \right|_{k\varphi} = O(k^{-N}) \|f\|_{k\varphi} \quad (3.1)$$

uniformly over $f \in H_{k\varphi}^{\varepsilon k}(U)$ and $z \in W$.

(2) (Decay away from the Diagonal) There exist constants $C, c > 0$ such that

$$|B_k^\varepsilon(z, z')|_{k\varphi} \leq Ck^n e^{-c \operatorname{dist}(z, z')} \text{ for all } z, z' \in U. \quad (3.2)$$

Remark 3.2. Our terminology differs slightly from that of Berman–Berndtsson–Sjöstrand in that we include the decay away from the diagonal as part of the definition. It is sometimes convenient to allow $N = \infty$ in which case the $O(k^{-\infty})$ term is understood as meaning the bound holds for any given $N \in \mathbb{N}$. One could relax the assumption that B_k^ε be holomorphic and instead assume that it is “almost holomorphic” as in [3], but we will have no need for this. When necessary we shall refer to this as a local PBK with respect to W or χ if we need to emphasise the dependence on these data.

The point of this definition is that, as proved in [3], a local PBK approximates the globally defined PBK in a neighbourhood of the diagonal of $W' \times W'$ for any $W' \Subset W$.

Theorem 3.3. (*Glueing Local Partial Bergman Kernels*) *With notation above, suppose B_k^ε is a local PBK of order N on W and suppose that ε is sufficiently small so that the forbidden region D_ε is relatively compact in W . Then if $W' \Subset W$ we have for all $r \geq 0$*

$$K_k^\varepsilon(x, y) = B_k^\varepsilon(x, y) + O_{Cr}(k^{n+r/2-N}) \text{ for all } x, y \in W'. \quad (3.3)$$

Proof. The argument here is similar to that of [3]. Let $x, y \in W'$ and $f := K_{k,y}^\varepsilon|_U \in H_{k\varphi}^{\varepsilon k}(U)$. The almost reproducing property of the local PBK B_k^ε gives

$$K_k^\varepsilon(y, x) = f(x) = \langle K_{k,y}^\varepsilon, \chi B_{k,x}^\varepsilon \rangle_{k\varphi} + O(k^{-N}) \|f\|_{k\varphi}. \quad (3.4)$$

and using Corollary (2.7) this becomes

$$K_k^\varepsilon(y, x) = \langle K_{k,y}^\varepsilon, \chi B_{k,x} \rangle_{k\varphi} + O(k^{n/2-N}). \quad (3.5)$$

Now by definition, $\langle \chi B_{k,x}^\varepsilon, K_{k,y}^\varepsilon \rangle_{k\varphi}$ is the value at y of the L^2 -projection of $\chi B_{k,x}^\varepsilon$ onto the space of holomorphic sections that vanish to order at least εk along Y . That is,

$$\langle \chi B_{k,x}^\varepsilon, K_{k,y}^\varepsilon \rangle_{k\varphi} = \chi B_{k,x}^\varepsilon(y) - v(y)$$

where v is the L^2 -minimal solution of the equation

$$\bar{\partial}v = \bar{\partial}(\chi B_{k,x}^\varepsilon) \quad (3.6)$$

among all such v that vanish to order at least εk along Y (we remark that this makes sense as equation (3.6) in particular implies that v is holomorphic in a neighbourhood of Y as $\chi \equiv 1$ on W). Of course v also depends on x , but we omit that from notation.

Hence we wish to bound $v(y)$ which we do with the Hörmander technique applied with the extremal envelope φ_ε from (2.11). Observe that φ_ε is plurisubharmonic over all of X and moreover is strictly plurisubharmonic in the support of

$$\bar{\partial}(\chi B_{k,x}^\varepsilon) = (\bar{\partial}\chi)B_{k,x}^\varepsilon$$

since by hypothesis $\bar{\partial}\chi$ is supported outside of W and thus within the equilibrium set $\{z \in X : \varphi_\varepsilon(z) = \varphi(z)\}$. Thus we may apply the Hörmander estimate [7, Theorem 4.5, Chapter VIII] to see there exists a v solving (3.6) such that

$$\|v\|_{k\varphi} \leq \|v\|_{k\varphi_\varepsilon} \leq C\|(\bar{\partial}\chi)B_{k,x}^\varepsilon\|_{k\varphi_\varepsilon} = C\|(\bar{\partial}\chi)B_{k,x}^\varepsilon\|_{k\varphi}$$

for some constant C , where the first inequality uses $\varphi_\varepsilon \leq \varphi$ and the final equality uses the statement about the support of $\bar{\partial}\chi B_{k,x}^\varepsilon$. Finally observe that as B_k^ε decays away from the diagonal we in fact have

$$(\bar{\partial}\chi)B_{k,x}^\varepsilon = O(k^{-\infty}) \quad (3.7)$$

since $x \in W'$ is a bounded distance away from the support of $\bar{\partial}\chi$. Therefore $\|v\|_{k\varphi} = O(k^{-\infty})$ which by Proposition 2.6 implies $|v(y)|_{k\varphi} = O(k^{-\infty})$ as well, and one sees this estimate is even uniform over $x, y \in W'$. Thus we have proved the required statement in the C^0 -topology (i.e. when $r = 0$). The statement for higher r follows from this using the Cauchy-integral formula. \square

4. A LOCAL PARTIAL BERGMAN KERNEL FOR AN S^1 -INVARIANT METRIC ON THE DISC

In this section we construct a local Bergman kernel for a potential on the unit disc that is circle invariant. The method here is less general than the construction of Berman-Berndtsson-Sjöstrand but is better suited to dealing with partial Bergman kernels. Strictly speaking the content of this section is not needed for the proof of our main theorem, since it will be repeated when we generalize in Section 5; we have included it here as an illustration of the main ideas of our approach.

4.1. The Legendre transform. Let D be the unit disc $\{|z| < 1\}$ in \mathbb{C} . Suppose φ is a strictly plurisubharmonic function that depends only on $|z|$ that is smooth on the closure \bar{D} . We shall write $z = e^{t+i\theta}$ and $\varphi = \varphi(t)$. Then consider the Legendre transform

$$u(x) + \varphi(t) = xt, \quad x = \varphi'(t), \quad t = u'(x). \quad (4.1)$$

This is well defined since the assumption that the potential is strictly plurisubharmonic means φ is a strictly convex function of t and hence u is also strictly convex. We shall refer to $x =: \mu(z)$ as the *dual variable* (or *momentum variable*) and to $u(x)$ as the *symplectic potential*. This x is defined up to an additive constant, which we choose so $x = 0$ corresponds to $z = 0$ and suppose that $|z| = 1$ corresponds to $x = a > 0$ (the reader may wish to recall the model case in which $\varphi = |z|^2 = e^{2t}$ has $u(x) = \frac{1}{2}(x \ln x - x)$ as the symplectic potential).

Now define

$$\nu := \frac{n}{k},$$

and let $s_{n,k}$ be the monomial

$$s_{n,k}(z) := e^{-ku(\nu)} z^n = e^{-ku(\nu)} e^{n(t+i\theta)} = e^{-k(u(\nu) - \nu(t+i\theta))}. \quad (4.2)$$

Then

$$|s_{n,k}(z)|_{k\varphi} = e^{-k(\varphi(t)+u(\nu)-\nu t)} \quad (4.3)$$

so if we define

$$U(\nu, x) := u(\nu) - u(x) + (x - \nu)u'(x) \quad (4.4)$$

and replace $\varphi(t)$ and t by the dual variables $u(x)$ and x in (4.3) we have

$$|s_{n,k}(z)|_{k\varphi} = e^{-kU(\nu, x)}. \quad (4.5)$$

In particular, the normalization of $s_{n,k}$ is such that the maximum value of $|s_{n,k}(z)|_{k\varphi}$ is 1 and is attained for those z for which $x = \nu$ (this follows easily from the convexity of u). In fact,

$$U_x(\nu, x) = (x - \nu)u''(x)$$

so the unique turning point of $x \mapsto U(\nu, x)$ is at $x = \nu$.

4.2. The Local Partial Bergman Kernel. Fix $\sigma \in (0, a)$ and let $\chi = \chi(x)$ be a smooth positive cutoff function that is identically equal to 1 for $x \leq \sigma$ and has compact support in D . For $\varepsilon < \sigma$ define

$$A_k^\varepsilon(z, z') := \sum_{\varepsilon \leq \nu \leq \sigma} G_{n,k} \overline{s_{n,k}(z)} s_{n,k}(z') \text{ for } (z, z') \in D \times D \quad (4.6)$$

where

$$\frac{1}{G_{n,k}} := \int_D \chi(x) e^{-2kU(\nu, x)} dx. \quad (4.7)$$

Finally let $Y = \{0\}$ be the origin in D .

Theorem 4.1. Fix $\eta < \sigma$. Then A_k^ε has the almost reproducing property on $\mu^{-1}[0, \eta)$; that is

$$\left| \int_D \chi(x') f(z') \overline{A_k^\varepsilon(z, z')} e^{-2k\varphi(t')} dx' d\theta' - f(z) \right|_{k\varphi} = O(k^{-\infty}) \|f\|_{k\varphi} \quad (4.8)$$

for all $f \in H_{k\varphi}^{\varepsilon k}(D)$ and $z \in \mu^{-1}[0, \eta)$.

Proof. Let $f \in H_{k\varphi}^{\varepsilon k}(D)$ which can be written as $f = f_1 + f_2$ where

$$f_1(z) = \sum_{\varepsilon \leq \nu \leq \sigma} a_n z^n \text{ and } f_2(z) = \sum_{\nu > \sigma} a_n z^n \quad (4.9)$$

and $\nu = n/k$ as before. Obviously f_1 and f_2 are L^2 -orthogonal, so

$$\|f\|_{k\varphi}^2 = \|f_1\|_{k\varphi}^2 + \|f_2\|_{k\varphi}^2. \quad (4.10)$$

Now by symmetry,

$$\int_D \chi(z') f(z') \overline{A_k^\varepsilon(z, z')} e^{-2k\varphi(t')} dz' = f_1(z), \quad (4.11)$$

so all that we have to do is prove

$$|f_2(z)|_{k\varphi} \leq O(k^{-\infty}) \|f_2\|_{k\varphi} \text{ for } x = \mu(z) < \eta.$$

To this end, fix some τ with $\varphi'(\tau) \in (\eta, \sigma)$. If $z = e^{t+i\theta}$ has $\mu(z) < \eta$ then $\varphi'(t) < \eta < \varphi'(\tau)$ and so by strict convexity of φ this implies $t \leq \tau - \delta$ for some $\delta > 0$ uniformly over all such z . Then Cauchy's inequalities give

$$|a_n e^{n\tau}| \leq \sup_{\theta} |f_2(e^{\tau+i\theta})| = e^{k\varphi(\tau)} \sup_{\theta} |f_2(e^{\tau+i\theta})|_{k\varphi}. \quad (4.12)$$

Therefore

$$|f_2(z)|_{k\varphi} \leq \sum_{n > \sigma k} |a_n z^n| e^{-k\varphi(t)} \leq e^{-k\varphi(t)} \sup_{\theta} |f_2(e^{\tau+i\theta})| \sum_{n > \sigma k} e^{n(t-\tau)}. \quad (4.13)$$

Now, by Proposition 2.6, there is a constant C such that if $t < \tau$ then

$$|f_2(e^{t+i\theta})|_{k\varphi} \leq Ck \|f_2\|_{k\varphi}, \quad (4.14)$$


and hence

$$|f_2(e^{t+i\theta})|_{k\varphi} \leq Ck \|f_2\|_{k\varphi} e^{-k[\varphi(t)-\varphi(\tau)]} e^{k\sigma[t-\tau]}. \quad (4.15)$$

Thus we will get an exponentially small multiple of $\|f_2\|_{k,\varphi}$ provided that

$$\varphi(t) - \varphi(\tau) - \sigma(t - \tau) > \delta' \quad (4.16)$$

for some positive δ' . But this is clear from the following picture:



That is, by convexity of φ the quotient

$$\frac{\varphi(\tau) - \varphi(t)}{\tau - t} \quad (4.17)$$

is bounded above by the slope $\varphi'(\tau)$ which assumed to be strictly less than σ . \square

5. A LOCAL S^1 -INVARIANT BERGMAN KERNEL ON X

5.1. Outline of the construction. We return to the case in which Y is a divisor in a compact complex manifold X , and L is a complex line bundle with hermitian metric $e^{-\varphi}$ that has strictly positive curvature. We assume from now on that there exists a neighbourhood U of Y that admits a holomorphic S^1 -action so that all the data are invariant when restricted to U (that is, the action lifts to $L|_U$ preserving the hermitian metric, and Y is fixed pointwise by the action). In this section we shall construct a local PBK on U whose asymptotics we can understand as k tends to infinity.

There are several ingredients to this construction. First we exploit the relation between the eigenspaces of the S^1 -action on $H^0(U, L^k)$ and the order of vanishing along Y . By hypothesis, S^1 -acts on the fibre L_p for any $p \in Y$ with some weight w (which is the same for every $p \in Y$) and by renormalising the action we may assume without loss of generality that $w = 0$. Write

$$H^0(U, \mathcal{L}^k) = \bigoplus_n V_k(n)$$

where $V_k(n)$ is the subspace of elements of weight n . Then as Y is fixed pointwise

$$H^0(Y, L^k \otimes \mathcal{S}_Y^n) = \bigoplus_{j \geq n} V_k(j).$$

Using the presence of the S^1 -action, we then have an identification

$$V_k(n) = H^0(U, L^k \otimes \mathcal{S}_Y^{n+1}) / H^0(U, L^k \otimes \mathcal{S}_Y^n) = H^0(Y, L^k \otimes \mathcal{O}(-nY)|_Y). \quad (5.1)$$

This is made explicit by a choice $\sigma \in H^0(U, \mathcal{O}(Y))$ of defining section for Y . Thinking of U as a disc bundle $\pi: U \rightarrow Y$, the isomorphism (5.1) is given by pulling back to U and then multiplying by σ^n . To put it another way, if f is a holomorphic section of $L^k|_U$ we have an expansion

$$f = \sum_n f_n \sigma^n$$

where $f_n \in H^0(Y, L(-\nu Y)|_Y^k)$ and $\nu := n/k$. When U is a disc centered at the origin with the standard S^1 -action and $Y = \{0\}$, this just reduces to power series expansion of holomorphic functions on the disc.

Next we define a hermitian inner product on $H^0(Y, L(-\nu Y)^k|_Y)$ as follows. Fix an S^1 -invariant cut-off function χ which is identically 1 in a neighbourhood of Y and supported in U . Then, with σ as above, we set

$$\|f_n\|_{\nu, k, \chi}^2 = \int_X \chi \pi_* (|f_n|^2 |\sigma|^{2n}) \omega^{[n]} \text{ for } f_n \in H^0(Y, L(-\nu Y)^k|_Y). \quad (5.2)$$

It turns out that this inner product is equal to the L^2 -inner-product with respect to certain hermitian metrics $e^{-2\eta_\nu}$ on $L(-\nu Y)^k|_Y$ and certain volume forms on Y :

Proposition 5.1. *(Proposition 5.12) Fix ε' slightly larger than ε . Then for $\nu \in [0, \varepsilon']$ there is a strictly plurisubharmonic potential η_ν on $L(-\nu Y)|_Y$ and a volume element $dV_{\nu, \hbar}$ on Y depending smoothly on ν and $\hbar := 1/\sqrt{k}$, such that*

$$\|f_n\|_{\nu, k, \chi}^2 = \int_X |f_n|_{k\eta_\nu}^2 dV_{\nu, \hbar}.$$

The proof of this proposition is essentially a standard stationary phase argument for the integral (5.2). To give a rough idea of the argument, consider the local situation. In standard local coordinates $(z = e^{t+i\theta}, w_1, \dots, w_n)$ so that Y is given locally by $z = 0$, our potential $\varphi = \varphi(t, w)$ and is convex in t for fixed w . Introduce (locally) the parametrized Legendre transform u characterised by

$$u(x, w) + \varphi(t, w) = xt \text{ where } x = \varphi_t \text{ and } t = u_x.$$

Here x is the moment map of the S^1 -action. On the other hand, the integrand in (5.2) then takes the form

$$|f_n(w)|^2 \exp(2k(u(x, w) - (x - \nu)u_x(x, w))) \frac{dx \wedge d\theta}{2\pi}$$

multiplied by a smooth volume element in the w variables, where $\nu = n/k$. So standard asymptotic methods give a leading term of the form

$$\frac{1}{\sqrt{u_{xx}(\nu, w)}k} e^{2ku(\nu, w)} |f_n(w)|^2$$

on integrating with respect to x and θ , since the exponential is stationary at $x = \nu$. Thus we see the negative of the Legendre transform $-u(\nu, w)$ appear as a local potential for $L(-\nu Y)|_Y$. We shall show that $w \mapsto -u(\nu, w)$ is strictly psh in w and that these local potentials indeed patch together to give a potential η_ν on $L(-\nu Y)|_Y$. The existence, and properties, of the required volume $dV_{\nu, \hbar}$ then comes from an application of Laplace's method.

Now the projection from the L^2 -sections of $L(-\nu Y)|_Y^k$ to the holomorphic ones is given by an integral kernel on $Y \times Y$ that we denote by $G_{n, k}$. Putting all of this together we shall prove:

Theorem 5.2. *(Theorem 5.22). Fix ε' slightly larger than ε . Then the quantity*

$$B_k^\varepsilon := \sum_{n=\varepsilon k}^{\varepsilon' k} G_{n, k} \sigma^n \boxtimes \bar{\sigma}^n$$

is an S^1 -local PBK for Y .

We refer the reader to Section 5.6 for the precise definition of a S^1 -local PBK (which merely modifies slightly the decay away from the diagonal property). Of course this theorem is to be understood as holding with respect to the chosen cut-off function χ . This is a rather precise formula for the local PBK, which we shall exploit to understand its asymptotics as k tends to infinity.

We now go through the details of these two results.

5.2. Circle-invariant set-up. Let X, Y, L and φ be as before and set $\omega = dd^c\varphi$. We also fix a defining section $\sigma \in H^0(Y, \mathcal{O}(Y))$ for Y . We suppose that there is an open neighbourhood U of Y that admits a holomorphic S^1 -action with Y as the fixed point set, so that U is covered by charts that admit “standard coordinates” (z, w) so that

$$\lambda \cdot (z, w) = (\lambda z, w) \quad (5.3)$$

and so that $\sigma = z$ in these coordinates. We also assume that this action lifts to $L|_U$ preserving φ in such a way that the S^1 -action is trivial over points in Y . By abuse of notation we let $\mathcal{O}(Y)$ denote both the line bundle associated to the divisor Y as well as its sheaf of sections (so is equal to the normal bundle of Y in X). We let $\mu: X \rightarrow \mathbb{R}$ denote the Hamiltonian of this S^1 -action normalised so $Y = \mu^{-1}(0)$ and always assume that ε is sufficiently small so that $\mu^{-1}[0, \varepsilon) \Subset U$.

As we have a holomorphic S^1 -action, we can identify our tubular neighbourhood U of Y holomorphically with a disc-subbundle of $\mathcal{O}(Y)$.

Lemma 5.3. *The neighbourhood U of Y is biholomorphic to a disc subbundle $\pi: D \rightarrow Y$ of $\mathcal{O}(Y)$. Furthermore any S^1 -invariant line bundle L' on U is canonically isomorphic to $\pi^*L'|_Y$.*

Proof. Let (z_α, w_α) and (z_β, w_β) be two sets of standard coordinates. The transition functions between them are necessarily of the form

$$z_\beta = \lambda_{\alpha\beta}(z_\alpha, w_\alpha)z_\alpha \text{ and } w_\beta = \tau_{\alpha\beta}(z_\alpha, w_\alpha) \quad (5.4)$$

where $\lambda_{\alpha\beta}$ is holomorphic and takes values in \mathbb{C}^* . Then both $\lambda_{\alpha\beta}$ and $\tau_{\alpha\beta}$ must be S^1 -invariant as well, which means their dependence upon z_α must be trivial. Hence we have

$$z_\beta = \lambda_{\alpha\beta}(w_\alpha)z_\alpha \quad (5.5)$$

is actually linear in the z coordinate. Since locally Y is given by $z_\alpha = 0$ (resp. $z_\beta = 0$) the $\lambda_{\alpha\beta}$ are the transition functions for the line bundle $\mathcal{O}(Y)$, so we have the desired biholomorphism. The argument for the second statement is the same, as the transition functions for an S^1 -invariant bundle L' must be independent of the normal variables z_α in standard coordinates. \square

Now we recall the eigenspace decomposition we have denoted by

$$H^0(U, L^k) = \bigoplus_n V_k(n)$$

where $V_k(n)$ denotes the subspace of weight n and that here and henceforth we are setting

$$\nu := \frac{n}{k}.$$

Lemma 5.4. *The map*

$$H^0(Y, L(-\nu Y)|_Y^k) \rightarrow V_k(n) \text{ given by } f_n \mapsto \pi^* f_n \sigma^n$$

is an isomorphism.

We omit the proof.

Lemma 5.5. *Given $f \in H^0(U, L^k)$ there is a unique sequence*

$$f_n \in H^0(Y, L(-\nu Y)|_Y^k), \text{ for } n \geq 0$$

such that

$$f = \sum_{n \geq 0} (\pi^* f_n) \sigma^n.$$

Proof. We have that $(\pi^* f_n) \sigma^n$ is just the weight- n component of f with respect to the S^1 -action. We use here the fact that a weight-zero section on U is canonically the same as the pull-back of a section over Y . That f_n lives where claimed is immediate from the definitions. \square

From now on we regard U as a disc-subbundle of $\mathcal{O}(Y)$ without further comment, and drop the π^* from notation so we identify $L^k \otimes \mathcal{O}(-nY)$ with $\pi^*(L(-\nu Y)^k|_Y)$ and also identify sections f_n of $L(-\nu Y)|_Y^k$ with their pull-back π^*f_n to U .

Our next order of business is to define hermitian inner products on $H^0(Y, L(-\nu Y)|_Y^k)$. Pick ε' slightly larger than ε and a domain Ω' arranged as follows:

$$\mu^{-1}[0, \varepsilon] \Subset \Omega' \Subset \mu^{-1}[0, \varepsilon'] \Subset U.$$

Definition 5.6. Fix once and for all an S^1 -invariant cut-off function

$$\chi \in C_0^\infty(U) \text{ with } \chi \equiv 1 \text{ on } \mu^{-1}[0, \varepsilon'].$$

Then for $f_n, g_n \in C^\infty(Y, L(-\nu Y)|_Y^k)$ define

$$\langle f_n, g_n \rangle_{\nu, k, \chi} := \int_U \chi(f_n, g_n)_{k\varphi} |\sigma|^{2\nu k} \omega^{[n]}. \quad (5.6)$$

(We emphasise our abuse of notation in that on the right hand side f_n and g_n are being identified with their pullback, so as to be defined on U).

So by construction if $f = \sum_n f_n \sigma^n$ and $g = \sum_n g_n \sigma^n$ are the expansions of two functions in $H^0(U, L^k)$ as in Lemma 5.5 we have

$$\int_X \chi(f, g)_{k\varphi} \omega^{[n]} = \int_U \chi(f, g)_{k\varphi} \omega^{[n]} = \sum_n \langle f_n, g_n \rangle_{\nu, k, \chi}.$$

5.3. The local Legendre Transformation. Our next goal is better to understand the hermitian inner products on $H^0(Y, L(-\nu Y)|_Y^k)$ defined in (5.6). In this section we shall do so locally over a small chart in Y , and in the next we will see how this globalises over Y . So assume we have standard coordinates z, w_1, \dots, w_{n-1} as in (5.3) on some patch of the form $U_\alpha = \pi^{-1}(W_\alpha)$ for some open subset W_α of Y and write

$$z := e^{t+i\theta}.$$

Recall that this includes the assumption that the defining section σ for Y is locally given by $\sigma = z$ in this coordinate system, and on this chart the hermitian metric on L is given by $e^{-2\varphi}$ where φ is a function of t and w . So, for f supported in such a coordinate chart

$$\|f\|_{\nu, k, \chi}^2 = \int \chi e^{-2k(\varphi - \nu t)} |f|^2 \omega^{[n]}.$$

We fix $\nu > 0$ and investigate the large k asymptotic behaviour of this integral. By standard principles doing the t -integral first, this will be exponentially small in k unless the critical point $\varphi_t = \nu$ occurs in the support of f . Assuming this to be the case, the main term for large k is

$$e^{-2k(\varphi(t_\nu) - \nu t_\nu)} = e^{2ku(\nu, w)}$$

where u is the parametrized Legendre transform defined as follows. Introduce the dual variables

$$x := \varphi_t \quad (5.7)$$

which is precisely the moment map of the S^1 -action, so Y is given by $x = 0$.

Definition 5.7. The parameterized Legendre transform u is characterized by

$$\varphi(t, w) + u(x, w) = tx. \quad (5.8)$$

Thus dually to (5.7) we have

$$t = u_x.$$

Lemma 5.8. *For each fixed x the function $w \mapsto -u(x, w)$ is strictly plurisubharmonic. Moreover, if we introduce the connection 1-form*

$$\alpha = \frac{1}{\varphi_{tt}} d^c x = \frac{1}{2\pi\varphi_{tt}} J dx = \frac{1}{2\pi} \left(d\theta + \frac{i}{\varphi_{tt}} (\varphi_{t\bar{a}} d\bar{w}^{\bar{a}} - \varphi_{ta} dw^a) \right), \quad (5.9)$$

then

$$\omega = \omega_\varphi = dx \wedge \alpha + (dd^c)_w(-u(x, \cdot)), \quad (5.10)$$

where the notation indicates fixing x and computing dd^c of $w \mapsto -u(x, w)$.

Proof. By elementary computation,

$$\begin{aligned} d\varphi &= \varphi_t dt + \varphi_a dw^a + \varphi_{\bar{a}} d\bar{w}^{\bar{a}}, \\ Jd\varphi &= \varphi_t d\theta - i\varphi_a dw^a + i\varphi_{\bar{a}} d\bar{w}^{\bar{a}}, \\ dJd\varphi &= \varphi_{tt} dt \wedge d\theta + \varphi_{ta} dw^a \wedge d\theta + \varphi_{t\bar{a}} d\bar{w}^{\bar{a}} \wedge d\theta \\ &\quad + i(\varphi_{t\bar{a}} dt \wedge d\bar{w}^{\bar{a}} - \varphi_{ta} dt \wedge dw^a) + 2i\varphi_{a\bar{b}} dw^a \wedge d\bar{w}^{\bar{b}}. \end{aligned}$$

Since $x = \varphi_t$,

$$dx = \varphi_{tt} dt + \varphi_{ta} dw^a + \varphi_{t\bar{a}} d\bar{w}^{\bar{a}} \quad (5.11)$$

so

$$dx \wedge \alpha = \frac{1}{2\pi} \left\{ \varphi_{tt} dt + \varphi_{ta} dw^a + \varphi_{t\bar{a}} d\bar{w}^{\bar{a}} \right\} \wedge \left\{ d\theta + \frac{i}{\varphi_{tt}} (\varphi_{t\bar{a}} d\bar{w}^{\bar{a}} - \varphi_{ta} dw^a) \right\} \quad (5.12)$$

$$= dd^c \varphi + \frac{i}{\pi} \left\{ \frac{1}{\varphi_{tt}} \varphi_{ta} \varphi_{t\bar{b}} - \varphi_{a\bar{b}} \right\} dw^a \wedge d\bar{w}^{\bar{b}}. \quad (5.13)$$

Hence we need to prove that

$$(dJd)_w u = 2i \left\{ \frac{1}{\varphi_{tt}} \varphi_{ta} \varphi_{t\bar{b}} - \varphi_{a\bar{b}} \right\} dw^a \wedge d\bar{w}^{\bar{b}}. \quad (5.14)$$

This follows by careful differentiation of (5.8). First, by differentiation with respect to w^a ,

$$\varphi_a(t, w) + u_a(x, w) = 0 \quad (5.15)$$

where in the first term we are holding t fixed and in the second we are holding x fixed. Differentiating again with respect to \bar{w}^b , holding x fixed,

$$\varphi_{a\bar{b}} + \varphi_{ta} t_{\bar{b}} = -u_{a\bar{b}} \quad (5.16)$$

Next, differentiation of $x = \varphi_t(x, w)$ with respect to \bar{w}^b (holding x fixed)

$$\varphi_{t\bar{b}} + \varphi_{tt} t_{\bar{b}} = 0. \quad (5.17)$$

If we insert this into (5.16), we obtain

$$-u_{a\bar{b}} = \varphi_{a\bar{b}} - \varphi_{ta} \varphi_{t\bar{b}} / \varphi_{tt} \quad (5.18)$$

This proves (5.14), from which it follows that $-u(x, w)$ is strictly plurisubharmonic in w for fixed x . \square

Lemma 5.9. *For fixed w the map $x \mapsto u(x, w)$ is strictly convex.*

Proof. This follows as $u_{xx} = t_x$ and $x_t = \varphi_{tt} > 0$ as φ is assumed to have strictly positive curvature. \square

5.4. The Global Legendre Transform.

Definition 5.10. Let $u(x, w)$ be the locally defined Legendre transform from (5.8). We set

$$\eta_\nu(w) := -u(\nu, w)$$

Lemma 5.11. *The above locally defined expression for η_ν gives a well-defined potential on $L(-\nu Y)|_Y$ whose curvature $dd^c \eta_\nu$ is strictly positive (and bounded away from 0 as ν ranges in a bounded interval).*

Proof. Consider a cover by standard coordinates (z_α, w_α) on charts U_α as in (5.3). We assume the line bundle L is trivialized over each U_α with transition functions $\mu_{\alpha\beta}$ and that the metric on L has potential φ_α over U_α . As we have already seen (5.5)

$$z_\beta = \lambda_{\alpha\beta} z_\alpha$$

where $\lambda_{\alpha\beta}$ is a holomorphic function of w_α in $U_\alpha \cap U_\beta$ giving the transition functions for $\mathcal{O}(Y)|_Y$. Now $\partial_{t_\alpha} \varphi_\alpha = \partial_{t_\beta} \varphi_\beta$ since they are both equal to the globally defined moment map x . So by definition of the Legendre transform along the set $\{x = \nu\}$ we have

$$\begin{aligned} \varphi_\alpha(t_\alpha, w_\alpha) + u_\alpha(\nu, w_\alpha) &= t_\alpha \nu \\ \varphi_\beta(t_\beta, w_\beta) + u_\beta(\nu, w_\beta) &= t_\beta \nu \end{aligned}$$

Now

$$t_\alpha = \log |z_\alpha| = \log |\lambda_{\alpha\beta}| + \log |z_\beta| = \log |\lambda_{\alpha\beta}| + t_\beta$$

and as φ is a potential on L we have $\varphi_\alpha = \varphi_\beta + \log |\mu_{\alpha\beta}|$. So we obtain

$$u_\beta(\nu, w_\beta) = t_\beta \nu - \varphi_\beta = t_\alpha \nu - \varphi_\alpha + \log(|\mu_{\alpha\beta}|) - \nu \log(|\lambda_{\alpha\beta}|) = u_\alpha(\nu, w_\alpha) + \log(|\mu_{\alpha\beta}| |\lambda_{\alpha\beta}|^{-\nu})$$

which is precisely the statement that $\eta_\nu(\cdot) = -u(\nu, \cdot)$ is a well-defined potential on $L(-\nu Y)|_Y$. The positivity of the curvature of this potential is a local calculation, and is the content of Lemma 5.8 \square

One of our main uses of this potential is the following expression for the hermitian metric we defined on $H^0(Y, L(-\nu Y)|_Y^k)$. Write $\omega_{x,u} = -(\text{dd}^c)_w u(x, w)$, so by Lemma 5.8

$$\omega_\varphi = dx \wedge \alpha + \omega_{x,u}, \quad \text{with } \alpha = \frac{1}{\varphi_{tt}} d^c x$$

as before.

Proposition 5.12. *There exist volume forms $dV_{\nu, \hbar}$ on Y for $\nu \in (0, \varepsilon')$ such that the inner-product on sections of $L(-\nu Y)|_Y^k$ defined in (5.6) is the L^2 -inner-product with respect to the potential η_ν and volume form $dV_{\nu, \hbar}$, i.e.*

$$\|f\|_{\nu, k, \chi}^2 = \int_Y |f|_{\eta_\nu}^2 dV_{\nu, \hbar} \text{ for all } f \in C^\infty(Y, L(-\nu Y)|_Y^k).$$

In fact

$$dV_{\nu, \hbar} = \hbar \sqrt{\frac{2\pi}{u_{xx}(\nu, w)}} \omega_{\nu, u}^{[n-1]} A(\nu, \hbar) \quad (5.19)$$

where $A(\nu, \hbar)$ is a smooth function and $A(\nu, 0) = 1$.

Proof. Let (z, w_1, \dots, w_{n-1}) be standard coordinates on some chart of the form $U_\alpha = \pi^{-1}(W_\alpha)$ for some $W_\alpha \subset Y$. Without loss of generality we may assume that f is supported in W . Observe that

$$\omega_\varphi^{[n]} = dx \wedge \alpha \wedge \omega_{x, \nu}^{[n-1]} = \frac{1}{2\pi} dx \wedge d\theta \wedge \omega_{x, u}^{[n-1]}$$

Then chasing definitions

$$\|f\|_{\nu, k, \chi}^2 = \frac{1}{2\pi} \int_{U_\alpha} \chi |f(w)|_{\eta_\nu}^2 e^{-2k(\varphi - \nu t - \eta_\nu)} dx \wedge d\theta \wedge \omega_{x, u}^{[n-1]}.$$

This can be calculated by performing the x -integral first, to obtain an expression as an integral on W_α . This gives the existence of the volume form $dV_{\nu, \hbar}$ (which is clearly well-defined over all of Y). Now observe that

$$\begin{aligned} \varphi(t, w) - \nu t - \eta_\nu(w) &= \varphi(t, w) - xt - (\nu - x)t + u(\nu, w) \\ &= -u(x, w) - (\nu - x)u_x(x, w) + u(\nu, w) \\ &= u_{xx}(x, w)(x - \nu)^2 + q(x, \nu, w) \end{aligned}$$

where q vanishes to order at least 3 at $x = \nu$. Thus the x -integral can be calculated using Laplace's method, proving that $dV_{\nu, \hbar}$ is smooth in \hbar , and moreover giving the stated leading term for $dV_{\nu, \hbar}$ (this is a simple case of the Laplace method described in Appendix C, see in particular Remark C.4). \square

Remark 5.13. It is worth noting that $dV_{\nu, \hbar} = O(\hbar)$, so $\hbar^{-1}dV_{\nu, \hbar}$ lie in a compact set of smooth volume forms as \hbar tends to 0 and ν ranges in a bounded interval.

5.5. The extremal envelope. The circle-invariant set-up allows us to identify the extremal envelope and the forbidden region explicitly in terms of the moment map, the potential and the Legendre transform.

In the neighbourhood U of Y , note that the locally defined expression $\varepsilon t - u(\varepsilon, \cdot)$ defines a potential on $L|_U$. This is because in U , εt is a potential on $\mathcal{O}(\varepsilon Y)$ and $\eta_\varepsilon = -u(\varepsilon, \cdot)$ is a potential on $L(-\varepsilon Y)$, so their sum is a potential on L . Note further that

$$\mu(t, w) = \varepsilon \Rightarrow \varepsilon t - u(\varepsilon, w) = \varphi(t, w)$$

by definition of the Legendre transform. Define

$$\psi_\varepsilon(t, w) = \begin{cases} \varepsilon t - u(\varepsilon, w) & \text{if } \mu(t, w) \leq \varepsilon, \\ \varphi(t, w) & \text{otherwise} \end{cases} \quad (5.20)$$

This definition makes sense initially only in U , but can clearly be extended to equal φ over $X \setminus U$. So defined, ψ_ε is a continuous potential on L .

Remark 5.14. The definition (5.20) can be motivated as follows. The simplest possible function with correct Lelong number has the form $\varepsilon t + \text{const}$, where $t = \log |z|$ as before. This does not extend globally, so away from Y we try to patch it to $\varphi(t)$. For fixed w , the slope of $\partial_t \varphi(t, w)$ is equal to ε precisely when $\mu(t, w) = \varepsilon$, since $\partial_t \varphi = \mu$. Thus we extend $\varepsilon t + \text{const}$ across $\mu^{-1}(\varepsilon)$ to equal φ . For this to be continuous, we need the constant to be equal to $-u(\varepsilon, w)$, and we have arrived at (5.20).

Then we have

Theorem 5.15. *The function ψ_ε is the extremal envelope for (X, Y, L, ε) (cf. Definition 2.11). Moreover, for $(t, w) \in U$,*

$$\psi_\varepsilon(t, w) < \varphi(t, w) \text{ if and only if } \mu(t, w) < \varepsilon.$$

In particular the forbidden region D_ε is equal to $\mu^{-1}[0, \varepsilon)$.

Proof. We shall show first that $\psi_\varepsilon(t, w) < \varphi(t, w)$ iff $\mu(t, w) < \varepsilon$. This is a variant of the convexity argument used at the end of the proof of Theorem 4.1.

Fix w and let t_ε satisfy $\mu(t_\varepsilon, w) = \varepsilon$. Then, from the definitions,

$$\varepsilon t - u(\varepsilon, w) = \varepsilon t - (\varepsilon t_\varepsilon - \varphi(t_\varepsilon, w)) = \varepsilon(t - t_\varepsilon) + \varphi(t_\varepsilon, w).$$

Then

$$\varepsilon t - u(\varepsilon, w) - \varphi(t, w) = -(t - t_\varepsilon) \left(\varepsilon - \frac{\varphi(t_\varepsilon, w) - \varphi(t, w)}{t_\varepsilon - t} \right).$$

If $t < t_\varepsilon$ then the difference quotient is strictly less than the derivative at the upper end point $\varphi'(t_\varepsilon) = \varepsilon$. Hence the quantity in the large brackets is positive and so

$$\varepsilon t - u(\varepsilon, w) < \varphi(t, w) \text{ for } t < t_\varepsilon$$

as required.

We shall show next that ψ_ε is C^1 . The only issue is what happens near $\mu^{-1}(\varepsilon)$. It is easier to use (x, w) as local coordinates. Then

$$\psi_\varepsilon = \varepsilon u_x(x, w) - u(\varepsilon, w) \text{ for } x < \varepsilon \quad (5.21)$$

and

$$\psi_\varepsilon = x u_x(x, w) - u(x, w) \text{ for } x > \varepsilon \quad (5.22)$$

Since both expressions are equal for $x = \varepsilon$ it follows that all tangential derivatives agree on this hypersurface, whereas

$$\lim_{x \rightarrow \varepsilon^-} \partial_x \psi_\varepsilon(\varepsilon, w) = \varepsilon u_{xx}(\varepsilon, w)$$

and

$$\lim_{x \rightarrow \varepsilon^+} \partial_x \psi_\varepsilon(\varepsilon, w) = \lim_{x \rightarrow \varepsilon} x u_{xx}(x, w) = \varepsilon u_{xx}(\varepsilon, w).$$

Thus ψ_ε is C^1 as claimed.

A similar calculation, which we leave to the reader, shows that ψ_ε is psh: from the regularity just proved, this follows by showing that ψ_ε is psh on each side of the hypersurface $\mu^{-1}(\varepsilon)$.

Hence ψ_ε is a candidate for the extremal envelope φ_ε in the sense of Definition 2.11. We have to check that there is no better candidate. By definition, any other candidate must also equal φ on the set $X \setminus \mu^{-1}[0, \varepsilon)$.

Suppose for contradiction $\psi_\varepsilon < \varphi_\varepsilon$ at some point $(t_0, w_0) \in U$. Then there is a plurisubharmonic potential γ on L bounded above by φ with Lelong number at least ε along Y such that $\gamma(t_0, w_0) > \psi_\varepsilon(t_0, w_0)$. Since $\gamma \leq \varphi$ we must then have $\mu(t_0, w_0) < \varepsilon$ and so $\psi_\varepsilon = \varepsilon t - u(\varepsilon, w)$ near (t_0, w_0) . Suppose that $\gamma'(t_0, w_0) \geq \psi'_\varepsilon = \varepsilon$. Along the image of $t \mapsto (t, w_0)$ we have $\gamma'(t, w_0)$ is non-decreasing, so at the point (t, w) on $\mu^{-1}(\varepsilon)$ we have $\gamma > \psi_\varepsilon(t, w) = \varphi(t, w)$ which is absurd. Hence $\gamma'(t_0, w_0) < \varepsilon$. Again by the monotonicity of γ' along this image we see that $\gamma' < \varepsilon$ for all $t < t_0$ and so the Lelong number of u is strictly less than ε which is also absurd. Hence such a γ cannot exist, and we conclude $\psi_\varepsilon = \varphi_\varepsilon$ as desired. \square

Corollary 5.16. *With notation as above, the equilibrium set for Y with respect to ε is the complement of $\mu^{-1}[0, \varepsilon)$ and the forbidden region is $\mu^{-1}[0, \varepsilon)$*

Proof. Follows directly from the Theorem. \square

Remark 5.17. Berman has proved that the extremal envelope φ_ε is generally no better than $C^{1,1}$, and our explicit formula (5.20) displays precisely this regularity. On the other hand, we have seen in the course of the proof that $\psi_\varepsilon = \varphi_\varepsilon$ is conormal with respect to the hypersurface $\mu^{-1}(\varepsilon)$: that is to say

$$V_1 \dots V_N \psi_\varepsilon \in C^{1,1}$$

for any number of vector fields V_j , provided that these are all *tangential* to $\mu^{-1}(\varepsilon)$. It would be interesting to investigate the conormal regularity of the extremal envelope in other situations.

As an application of this explicit identification of the extremal envelope, we prove the following technical result.

Lemma 5.18. *Suppose that $f \in H_{k\varphi}^{\varepsilon', k}(U)$ and*

$$\Omega' \Subset \mu^{-1}[0, \varepsilon').$$

Then there are constants C and c such that

$$\sup_{\Omega'} |f|_{k\varphi} \leq C e^{-ck} \|f\|_{k\varphi, U} \quad (5.23)$$

Proof. The proof is similar to that of Proposition 2.13. Let $N_k = \sup_{\Omega'} |f|_{k\varphi}$ and set

$$v := \frac{1}{k} (\log |f|_{k\varphi} - \log N_k).$$

Then $v \leq 0$ and $\varphi + v$ is a competitor to be the envelope for νY on Ω' . Hence $\varphi + v \leq \psi_{\varepsilon'}$. Rearranging this gives that over Ω'

$$|f|_{k\varphi} \leq N_k \exp(-k(\varphi - \psi_{\varepsilon'})) \leq C k^n \exp(-ck) \|f\|_{k\varphi, U}$$

where we have used the L^2 implies L^∞ bound from Proposition 2.6 and set $c : \inf_{\Omega'}(\varphi - \psi_\nu) = c$ which is strictly positive as $\Omega' \Subset \mu^{-1}[0, \varepsilon')$. \square

5.6. A modified glueing result. We will need a slight modification of our glueing result that relaxes the decay away from the diagonal condition in the presence of a holomorphic S^1 -action.

Definition 5.19. We say that B_k^ε is an S^1 -local partial Bergman kernel if it has the almost reproducing property as in Definition 3.1 and the following decay away from the diagonal in standard coordinates:

$$|B_k^\varepsilon(z, w, z', w')|_{k\varphi} \leq Ck^n e^{-c(|w-w'|^2 + |x-x'|^2)} \text{ for all } (z, w), (z', w') \in U$$

where, we recall, $x = \mu(z, w)$ and $x' = \mu(z', w')$ is the value of the moment map at these points.

Theorem 5.20. *With the setup as above, suppose B_k^ε is an S^1 -local PBK of order N on $W \subset U$, and suppose that ε is sufficiently small so that the forbidden region D_ε lies in W . Then if W' is an open, relatively compact subset of W then we have for all $r \geq 0$ that*

$$K_k^\varepsilon(x, y) = B_k^\varepsilon(x, y) + O_{C^r}(k^{n/2+r/2-N}) \text{ for all } x, y \in W'.$$

Proof. The proof is the same as that of Theorem 3.3. The only place in which we used the decay away from the diagonal was in (3.7). But if $(z, w) \in W'$ and $\bar{\partial}\chi(z') \neq 0$ at a point (z', w') then x and x' are a bounded distance apart (as $\chi \equiv 1$ on W) and so (3.7) still holds. \square

5.7. The Local Partial Bergman Kernel. We are now ready to define our local PBK. For $\nu \in [0, \varepsilon')$ let $G_{n,k}$ denote the reproducing kernel on $Y \times Y$ for $L^k \otimes \mathcal{O}(-nY)|_Y$ with respect to the inner-product defined in (5.6).

Definition 5.21. Define

$$B_k^\varepsilon := \sum_{n=\varepsilon k}^{\varepsilon' k} G_{n,k} \sigma^n \boxtimes \bar{\sigma}^n. \quad (5.24)$$

So by our conventions made following Lemma 5.5, B_k^ε is a holomorphic section of $L^k|_U \boxtimes \bar{L}^k|_U$.

Theorem 5.22. *B_k^ε is an S^1 -local PBK for (ε, Y) on Ω' (with respect to the chosen cutoff function χ used in (5.6)).*

Before the proof we make a convenient definition:

Definition 5.23. In standard local coordinates let

$$U(\nu, x, w) := u(\nu, w) - u(x, w) - u_x(x, w)(\nu - x) \text{ for } \nu \in [0, \varepsilon']$$

Lemma 5.24. (1) *There exists a constant $c > 0$ such that $U(\nu, x, w) \geq c(x - \nu)^2$.*

(2) *For any $f \in H^0(L(-\nu Y)^k|_Y)$ we have*

$$|f\sigma^n|_{k\varphi} = |f|_{k\eta_\nu} e^{-kU(\nu, x, w)}.$$

Proof. The first statement follows as $U(\nu, x, w) = u_{xx}(w, x')(\nu - x)^2$ for some x' which is strictly bounded from below as $x \mapsto u(x, w)$ is strictly convex. The second statement is a simple calculation using the definitions and is left to the reader. \square

Proof of Theorem 5.22. We first show that B_k^ε has the almost reproducing property, i.e. for some $c, C > 0$ we have

$$|f(z) - (B_{k,z}^\varepsilon, \chi f)_{k\varphi}|_{k\varphi} \leq C e^{-ck} \|f\|_{k\varphi, U} \quad (5.25)$$

for all $f \in H_{k\varphi}^{\varepsilon k}(U)$ and $z \in \Omega'$. For this we know from Lemma 5.5 that we can write such an f as $f = g + s$ where

$$s = \sum_{n > \varepsilon' k} s_n \sigma^n \text{ and } g = \sum_{n = \varepsilon k}^{\varepsilon' k} g_n \sigma^n$$

and $s_n, g_n \in H^0(L(-\nu Y)^k|_Y)$. By integrating in the normal direction to Y first, we see by construction that if $z \in \Omega'$ we have

$$(\chi g, B_{k,z}^\varepsilon)_{k\varphi} = g(z) \text{ and } (\chi s, B_{k,z}^\varepsilon)_{k\varphi} = 0.$$

On the other hand as s vanishes to order at least $\varepsilon'k$ along Y we have from Lemma 5.18 that $\sup_{\Omega'} |s|_{k\varphi} \leq C e^{-ck} \|s\|_{k\varphi} \leq C e^{-ck} \|f\|_{k\varphi}$ for some $c, C > 0$. Putting this together gives (5.25).

Finally we prove that B_k^ε has the desired decay away from the diagonal property, i.e.

$$|B_k^\varepsilon(z, w, z', w')|_{k\varphi} \leq C k^n e^{-c(|w-w'|^2 + |x-x'|^2)} \text{ for all } (z, w), (z', w') \in U.$$

Fix (z, w) . Then by Cauchy-Schwarz and the definition of the function U (i.e. Lemma 5.24)

$$\begin{aligned} |B_k^\varepsilon(z, w, z', w')|_{k\varphi}^2 &\leq O(k) \sum_{\varepsilon k \leq n \leq \varepsilon' k} |G_{n,k}(w, w')|_{k\eta_\nu}^2 e^{-k(U(\nu, x, w) + U(\nu, x', w'))} \\ &\leq O(k) \sum_{\varepsilon k \leq n \leq \varepsilon' k} |G_{n,k}(w, w')|_{k\eta_\nu}^2 e^{-ck((x-\nu)^2 + (x'-\nu)^2)} \end{aligned}$$

for some $c > 0$. Recall that each $G_{n,k}$ decays exponentially fast away from the diagonal,

$$|G_{n,k}(w, w')|_{k\eta_\nu}^2 \leq O(k^{2n-1}) e^{-ck|w-w'|^2}. \quad (5.26)$$

We have given a proof of this classical fact in Theorem 2.9. Moreover this estimate is uniform as ν ranges in a bounded interval, as follows directly from our proof. (We remark that for a fixed volume form the Bergman kernel has decay at rate $O(k^{2n-2}) e^{-ck|w-w'|^2}$ since $\dim Y = n - 1$, and this decay still holds for volume forms that lie in a compact set; but in the above $G_{n,k}$ is taken with respect to the volume form $dV_{\nu, \hbar} = O(\hbar) = O(k^{-1/2})$ which explains the missing factor of k in (5.26).)

Hence

$$|B_k^\varepsilon(z, w, z', w')|_{k\varphi}^2 = O(k^{2n}) O(e^{-ck|w-w'|^2}) e^{-ck((x-\nu)^2 + (x'-\nu)^2)}$$

for some $c > 0$. On the other hand, by completing the square

$$(x - \nu)^2 + (x' - \nu)^2 = \frac{1}{2}(x - x')^2 + 2 \left(\nu - \frac{x + x'}{2} \right)^2 \geq \frac{1}{2}(x - x')^2$$

which gives the desired result. \square

6. PROOF OF MAIN THEOREMS

We now put what we have done together to prove the main theorems. The main task is to understand the asymptotics of our S^1 -local PBK

$$B_k^\varepsilon := \sum_{n=\varepsilon k}^{\varepsilon' k} G_{n,k} \sigma^n \boxtimes \bar{\sigma}^n. \quad (6.1)$$

The idea is to use the standard asymptotic expansion of the density functions on Y to expand the functions $G_{n,k}$ in powers of k , and then use the Euler-Maclaurin formula to evaluate the sum in terms on an integral, which by Laplace's method can also be expanded in powers of k . To display the main ideas and keep the proof short, we include an account of each of these standard techniques in the Appendix.

We start by recalling the well-known asymptotic expansion of the Bergman kernel. Let L' be an ample line bundle on a compact complex manifold Y of dimension d with positive hermitian metric $e^{-2\varphi}$ and let dV be a smooth volume form on Y . These define an L^2 -inner product on sections of L'^k and we let K'_k denote the reproducing kernel for the projection to the holomorphic sections, and let $\rho'_k(y) = K'_k(y, y)$ be the corresponding density function.

Theorem 6.1. *There exist smooth functions a_0, a_1, \dots on Y such that for any $p, r \geq 0$ there is an asymptotic expansion*

$$\rho'_k = a_0 k^d + a_1 k^{d-1} + \dots + a_p k^{d-p} + O_{C^r}(k^{d-p-1}).$$

Furthermore the a_i are universal quantities that depend smoothly on dV and the curvature of φ , in particular

$$a_0 = \frac{(dd^c\varphi)^d}{dV}. \quad (6.2)$$

Moreover given any $c > 0$ and background Kähler form ω_0 on Y , the $O_{C^r}(k^{d-p-1})$ error term may be taken uniformly over all $(L', e^{-\varphi})$ such that $dd^c\varphi \geq c\omega_0$ as well as uniformly over all volume forms that lie within a compact set.

Proof. The first statement is the famous asymptotic expansion of the Bergman kernel, due to Fefferman [9], Catlin [5], Tian [30] and Zelditch [31]. The statement that the error term may be taken uniformly follows from the same proofs (e.g. it is clear that this is the case for the local Bergman Kernel of [3], and the glueing theorem of [3, Thm 3.1] uses the Hörmander estimate which gives a uniform error given the assumed bound on $dd^c\varphi$, and for the variation with respect to dV see [3, Sec 2.4] which is also clearly uniform as dV varies in a compact set. \square

Proof. (Proof of Theorem 1.1 and Theorem 1.2) First we note that B_k^ε is an S^1 -local PBK which by Theorem 5.20 approximates the globally defined PBK, so in standard coordinates (z, w) we have

$$\rho_k^\varepsilon(z, w) = B_k^\varepsilon(z, w, z, w)e^{-2k\varphi(z, w)} + O(k^{-\infty}).$$

Hence the goal becomes to understand the asymptotics of the quantity

$$B := B_k^\varepsilon(z, w, z, w)e^{-2k\varphi(z, w)}.$$

From now on we will work with the variable $\hbar = k^{-1/2}$. By Lemma 5.11 the η_ν have positive curvature bounded from below uniformly over $\nu \in [0, \varepsilon']$. Observe that η_ν are smooth in ν and recall that $\hbar^{-1}dV_{\nu, \hbar}$ are volume forms that lie in a compact set (and moreover are smooth in \hbar and ν). Thus Theorem 6.1 implies there are smooth functions $a_i(\nu, w)$ on $[0, \varepsilon'] \times Y$ such that for any $r, p \geq 0$ we have

$$\hbar G_{n, k}(w, w)e^{-2\hbar^{-2}\eta_\nu(w)} = a_0(\nu, w)\hbar^{2-2n} + a_1(\nu, w)\hbar^{4-2n} + \dots + a_p(\nu, w)\hbar^{2p+2-2n} + O_{C^r}(\hbar^{2p+4-2n})$$

where, as usual, $\nu = n/k = \hbar^2 n$. To capture this information we define

$$\alpha(\nu, \hbar^2, w) := a_0(\nu, w) + a_1(\nu, w)\hbar^2 + \dots + a_p(\nu, w)\hbar^{2p}$$

which is clearly smooth in all variables.

Remark 6.2. For later use, observe that from the leading order term of $dV_{\nu, \hbar}$ given in (5.19) and the leading order term of the asymptotic expansion of the Bergman function given in (6.2) we have that

$$\alpha(0, 0, w) = a_0(0, 0) = \sqrt{\frac{u_{xx}(0, w)}{2\pi}}.$$

Now using Lemma 5.24 (i.e. the properties of our function U) gives

$$\begin{aligned} B &= \sum_{n=\varepsilon k}^{\varepsilon' k} G_{n, k}(w, w)|z|^{2n} e^{-2k\varphi(z, w)} \\ &= \sum_{n=\varepsilon k}^{\varepsilon' k} G_{n, k}(w, w)e^{-2k\eta_\nu(w)} e^{-2kU(\nu, x, w)} \\ &= \hbar^{1-2n} \left(\sum_{n=\varepsilon k}^{\varepsilon' k} e^{-2kU(x, \nu, w)} \alpha(\hbar^2 n, \hbar^2, w) + O_{C^r}(\hbar^{2p+2}) \right). \end{aligned}$$

Now this sum can be calculated using the Euler-Maclaurin formula (see B.2). To state this let

$$q_\hbar(s) := q(s, \hbar, x, w) = e^{-\hbar^{-2}U(x-\hbar s, x, w)} \alpha(x-\hbar s, \hbar^2, w)$$

and set

$$\xi = \frac{x - \varepsilon}{\hbar}.$$

Then

$$\hbar^{2n}B = \int_{-\infty}^{\xi} q_{\hbar}(s)ds + \sum_{j=0}^{m-1} A_j \hbar^j + O(\hbar^m) \quad (6.3)$$

where

$$A_0 = \frac{1}{2}q_{\hbar}(\xi) \text{ and } A_j = (-1)^{j-1} \frac{\beta_{j+1}}{(j+1)!} q_{\hbar}^{(j)}(\xi)$$

where β_j are the Bernoulli numbers. Moreover one sees directly that the coefficients A_j lift to the real blowup (see Appendix A) as in the statement of the Theorem.

Thus it remains only to analyse the integral in the right hand side of (6.3) which can be done by Laplace's method. In slightly more detail observe that $q_{\hbar}(s)$ has its unique critical point at $s = 0$. We rewrite the integral as

$$I := \int_{-\infty}^{\zeta} q_{\hbar}(s)ds = \hbar^{-1} \int_{-\infty}^{x-\varepsilon} q_{\hbar}\left(\frac{s}{\hbar}\right) ds$$

So this is precisely the setup of Laplace's method as discussed in Appendix C which shows that this integral has an asymptotic expansion in powers of \hbar , in fact

$$I = \frac{\sqrt{2\pi}\alpha(0,0,w)}{\sqrt{u_{xx}(0,w)}} \Phi(\sqrt{u_{xx}(0,w)}\zeta) + O(\hbar)$$

where as usual

$$\Phi(x) := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt.$$

Recall that $u_{xx}(0,w) = |v|$ where v is the generator of the S^1 -action. Thus plugging in (Remark 6.2)

$$\alpha(0,0,w) = \sqrt{\frac{|v|}{2\pi}},$$

as simple change of variables gives

$$I = \frac{1}{\sqrt{2\pi|v|}} \int_{-\infty}^{\frac{x-\varepsilon}{\hbar}} e^{-\frac{t^2}{2|v|^2}} dt + O(\hbar)$$

as required in the statement of Theorem 1.2. Finally Theorem 1.1 follows immediately from this by the behaviour of $\Phi(x)$ on the sets $\{x < \varepsilon\}$ and $\{x > \varepsilon\}$. \square

7. AN APPLICATION

We end with an application of our main theorem to the study of a certain natural function introduced by Ross-Witt Nyström [23] that one can associate to a divisor on a Kähler manifold.

Fix a line bundle L on a compact complex manifold X with hermitian metric $e^{-\varphi}$. Then the order of vanishing of sections along a divisor Y determines a finite length filtration

$$H^0(L^k) \supset H^0(L^k \otimes \mathcal{I}_Y) \supset H^0(L^k \otimes \mathcal{I}_Y^2) \supset \dots \supset \{0\}.$$

Let $\text{ord}_Y(s)$ denote the order of vanishing of a section s along Y , and suppose that $\{s_{\alpha,k}\}$ is an L^2 -orthonormal basis for $H^0(L^k)$ that is compatible with this filtration, i.e. for each j the set

$$\{s_{\alpha,k} : \text{ord}_Y(s_{\alpha,k}) \geq j\}$$

is a basis for $H^0(L^k \otimes \mathcal{I}_Y^j)$,

Definition 7.1. Define $M_k : X \rightarrow \mathbb{R}$ by

$$M_k(z) = \frac{\sum_{\alpha} \text{ord}_Y(s_{\alpha,k}) |s_{\alpha,k}|_{k\varphi}^2}{k \sum_{\alpha} |s_{\alpha,k}|_{k\varphi}^2}.$$

One can check directly that this definition does not depend on choice of compatible orthonormal basis, and thus defines a natural smooth function on X associated to Y and the hermitian metric on L .

Theorem 7.2. *Suppose φ and Y are invariant under an S^1 action on (X, L) . Then there is a neighbourhood of Y and an asymptotic expansion*

$$M_k = c_0 + c_{1/2}k^{-1/2} + c_1k^{-1} + \cdots + c_rk^{-N} + O_{Cr}(k^{-N-1/2}) \quad (7.1)$$

where c_i are smooth functions defined on this neighbourhood. Moreover c_0 is precisely the hamiltonian of the S^1 action normalized so $Y = c_0^{-1}(0)$.

Remark 7.3. It is shown in [23, Sec 8], without any assumption of the existence of an S^1 -action, that the limit

$$\mu := \limsup_{k \rightarrow \infty} M_k$$

converges almost everywhere on X . We do not know anything about the regularity of μ in general, but it does have a ‘‘push-forward’’ property analogous to that of the Duistermaat-Heckman formula [23, Thm. 8.1,8.3] with the moment polytope being replaced by the Okounkov body of (X, L) . The previous theorem shows that in the S^1 -invariant case we in fact have that $\mu = c_0$ is the Hamiltonian in a neighbourhood of Y which reaffirms this property.

Proof of Theorem 7.2. For simplicity write s_α for $s_{\alpha,k}$ and set $n_\alpha = \text{ord}_Y(s_\alpha)$. We also write $x: X \rightarrow \mathbb{R}$ for the hamiltonian of the S^1 -action normalized so $Y = \{x = 0\}$. So the partial Bergman kernel for (ε, Y) is given by

$$\rho_k^\varepsilon = \sum_{n_\alpha \geq \varepsilon k} |s_\alpha|_{k\varphi}^2 \quad \text{for } \varepsilon k \in \mathbb{N}.$$

By Theorem 1.2 there is a neighbourhood of the divisor and an expansion of the partial density in powers of $k^{1/2}$:

$$\rho_k^\varepsilon = b_0(\varepsilon)k^n + b_{1/2}(\varepsilon)k^{n-1/2} + \cdots + b_N(\varepsilon)k^{n-N} + O(k^{n-N-1})$$

where the $b_i(\varepsilon)$ are smooth functions on X with

$$b_0(\varepsilon) = \frac{1}{\sqrt{2\pi x}} \int_{-\infty}^{\sqrt{k}(x-\varepsilon)} e^{-\frac{t^2}{2x}} dt.$$

Now

$$\sum_{j=1}^{k\sigma} \rho_k^{\frac{j}{k}Y} = \sum_{j=1}^{k\sigma} \sum_{n_\alpha \geq j} |s_\alpha|_{k\varphi}^2 = \sum_{n_\alpha \leq k\sigma} n_\alpha |s_\alpha|_{k\varphi}^2 + \sum_{n_\alpha > k\sigma} k\sigma |s_\alpha|_{k\varphi}^2.$$

Thus on a smaller neighbourhood of Y (say where $x \leq \sigma/2$) we have

$$M_k = \frac{1}{k\rho_k} \sum_{j=1}^{k\sigma} \rho_k^{\frac{j}{k}Y} + O(k^{-\infty})$$

where ρ_k is the usual Bergman function on X . Now $\rho_k = k^n + O(k^{n-1})$ has a global asymptotic expansion in powers of k , and thus the Euler-Maclaurin formula gives the required expansion (7.1) for M_k with

$$c_0 = \int_0^\sigma b_0(s) ds.$$

Roughly speaking, for k large $b_0(s)$ is approximately 1 for $x > s$ and 0 for $x < s$, and so $c_0 = \int_0^\sigma b_0(s) ds \sim \int_0^x ds = x$. We claim that in fact for $x < \sigma/2$ we have

$$\int_0^\sigma b_0(s) ds = x + O(k^{-\infty})$$

To see this, integrate by parts to get

$$I := \int_0^\sigma b_0(s) ds = [sb_0(s)]_{s=0}^\sigma + \sqrt{k} \int_0^\sigma \frac{s}{\sqrt{2\pi x}} e^{-\frac{k(x-s)^2}{2x}} ds$$

Since $x < \sigma/2$ the boundary term is $O(k^{-\infty})$, and by direct calculation

$$\begin{aligned} k^{-1/2}I &= x \int_0^\sigma \frac{1}{\sqrt{2\pi x}} e^{-\frac{k(x-s)^2}{2x}} ds - \int_0^\sigma (x-s) \frac{1}{\sqrt{2\pi x}} e^{-\frac{k(x-s)^2}{2x}} ds + O(k^{-\infty}) \\ &= k^{-1/2}x + O(k^{-\infty}) \end{aligned}$$

as claimed. \square

APPENDIX A. THE REAL BLOW-UP

For the reader's convenience we collect here some elementary properties of a special case of the real blow-up. More details and constructions of greater generality can be found in Melrose's book [17, Chapter 5], as well as in [11] and [10, Section 2.3].


Definition A.1. Let $S_+^1 = \{e^{i\theta} : \theta \in [0, \pi]\} \subset \mathbb{R}^2$ be the upper-semicircle. The *real blow-up* of the upper-half space $\mathbb{R} \times \mathbb{R}_{\geq 0}$ at the point $p = (0, 0)$ is defined to be

$$[\mathbb{R} \times \mathbb{R}_{\geq 0}; p] := \mathbb{R}_{\geq 0} \times S_+^1$$

along with the *blow-down* map

$$\beta: [\mathbb{R} \times \mathbb{R}_{\geq 0}; p] \rightarrow \mathbb{R} \times \mathbb{R}_{\geq 0} \quad \beta(r, e^{i\theta}) = re^{i\theta}.$$

Thus $[\mathbb{R} \times \mathbb{R}_{\geq 0}; p]$ is the upper half-space with a copy of the semicircle S_+^1 inserted at the origin, which we consider as a manifold-with-corners. Then β is smooth, and moreover is a diffeomorphism away from the *exceptional set* $E = \beta^{-1}(p) = \{(0, e^{i\theta}) : \theta \in [0, \pi]\}$. The boundary of $[\mathbb{R} \times \mathbb{R}_{\geq 0}; p]$ has three pieces, namely E and the two axes $H_\pm = \{(r, \pm 1) : r \in \mathbb{R}_{\geq 0}\}$, and two corners $C_\pm = H_\pm \cap E$.



For higher dimensions, we let the real blow-up of $\mathbb{R}^n \times \mathbb{R}_{\geq 0}$ along $Y := \mathbb{R}^{n-1} \times \{p\}$ be $[\mathbb{R}^n \times \mathbb{R}_{\geq 0}; Y] := \mathbb{R}^{n-1} \times [\mathbb{R} \times \mathbb{R}_{\geq 0}; p]$, and by patching in the obvious way, this extends to define the real blow-up $[X \times \mathbb{R}_{\geq 0}; D \times \{0\}]$ for any smooth real manifold X and codimension 1 smooth divisor D .

Let x be the standard coordinate on \mathbb{R} and \hbar the standard coordinate on $\mathbb{R}_{\geq 0}$. Then the functions

$$\xi := \frac{x}{\hbar} \quad \text{and} \quad \hbar$$

give coordinates on the blow-up $B := [\mathbb{R} \times \mathbb{R}_{\geq 0}; p]$ on an open set containing the interior of E (precisely, $\xi = \cotan \theta$ and $\hbar = r \sin \theta$ for $(r, \theta) \in B$ with $\theta \neq 0$). Moreover in these coordinates $\beta(\xi, \hbar) = (\xi \hbar, \hbar)$. Around each of the corners C_\pm we have coordinates given by

$$x \quad \text{and} \quad \eta_\pm = \pm \frac{\hbar}{x}$$

(so $x = r \cos \theta$ and $\eta_\pm = \pm \tan \theta$) and in these coordinate $\beta(x, \eta_\pm) = (x, \pm x \eta_\pm)$.

Our interest will be in functions on $\mathbb{R} \times \mathbb{R}_{>0}$ that lift to smooth functions on the blow-up (i.e. that extend to a smooth function right up to the boundary). As a model example (which will be considered in more generality below) let

$$q(x, \hbar) = e^{-\hbar^{-2}x^2}$$

which is clearly smooth for $\hbar > 0$, and extends smoothly to the boundary away from $p := (0, 0)$. We claim the lift $\beta^*g := q \circ \beta$ extends to a smooth function on all of $[\mathbb{R} \times \mathbb{R}_{\geq 0}; p]$. To see this observe that in the coordinates (ξ, \hbar) we have

$$\beta^*q(\xi, \hbar) = q(\xi\hbar, \hbar) = e^{-\xi^2}$$

which is clearly smooth, and in coordinates (x, η_{\pm}) ,

$$\beta^*q(x, \eta_{\pm}) = q(x, \pm x\eta_{\pm}) = e^{\eta_{\pm}^{-2}}$$

which extends to a smooth function for all $\eta_{\pm} \geq 0$ by declaring it to be zero when $\eta_{\pm} = 0$.

APPENDIX B. THE EULER-MACLAURIN FORMULA

Let $U(\nu, x, w)$ be a smooth function, where $\nu \in [0, \varepsilon']$, $x \in \mathbb{R}_{\geq 0}$ and w ranges in an open relatively compact subset of \mathbb{C}^m . Suppose that U is such that $x \mapsto U(\nu, x, w)$ is strictly convex and

$$U(x, x, w) = 0 \quad \text{and} \quad U_x(x, x, w) = 0 \quad \text{for all } x, w. \quad (\text{B.1})$$

Also let $\alpha(\nu, \hbar^2, w)$ be a bounded smooth function where $\hbar^2 \in \mathbb{R}_{\geq 0}$ and fix also $\varepsilon \in [0, \varepsilon']$. For $\hbar > 0$ let

$$p_{\hbar}(t) := p(t, \hbar, x, w) := e^{-\hbar^{-2}U(\hbar^2 t, x, w)} \alpha(\hbar^2 t, \hbar^2, w)$$

Then our interest is in the behaviour of the sum

$$S := S(\hbar, x, w) := \hbar \sum_{n=\varepsilon k}^{\varepsilon' k} e^{-\hbar^{-2}U(\hbar^2 n, x, w)} \alpha(\hbar^2 n, \hbar^2, w) = \hbar \sum_{n=\hbar^{-2}\varepsilon}^{\hbar^{-2}\varepsilon'} p_{\hbar}(n) \quad (\text{B.2})$$

as \hbar tends to zero.

We recall the Bernoulli numbers are defined by $\beta_j = \beta_j(0)$ where

$$\frac{ze^{zx}}{e^z - 1} = \sum_{j \geq 0} \beta_j(x) \frac{z^j}{j!}.$$

Theorem B.1 (Euler-Maclaurin Formula). *Suppose $p: \mathbb{R} \rightarrow \mathbb{R}$ is smooth and $p^{(i)}(t)$ tends to zero as t tends to infinity for all $i \geq 0$. Then for any integers a and $m \geq 1$ we have*

$$\sum_{n=a}^{\infty} p(n) = \int_a^{\infty} p(t) dt + \frac{1}{2}p(a) - \sum_{j=1}^{m-1} \frac{\beta_{j+1}}{(j+1)!} p^{(j)}(a) - \int_a^{\infty} \frac{\beta_m(\{1-t\})}{m!} p^{(m)}(t) dt,$$

where $\{t\}$ denotes the fractional part of t .

Proof. This is proved, for example, in [28, Theorem 5]. □

Since $x \mapsto U(\nu, x, w)$ has its (unique) critical point at $x = \nu$ it is convenient to make a further change of variables by setting

$$q_{\hbar}(s) = p_{\hbar} \left(\frac{x - \hbar s}{\hbar^2} \right).$$

Observe then that from (B.1)

$$q_{\hbar}(s) = e^{-\hbar^{-2}U(x - \hbar s, x, w)} \alpha(x - \hbar s, x, w) = e^{-U_{xx}(x, x, w)s^2 + O(\hbar)} \alpha(x - \hbar s, x, w)$$

which has bounded derivatives with respect to s uniformly as \hbar tends to 0.

Proposition B.2. *Set*

$$\xi = \frac{x - \varepsilon}{\hbar}.$$

Then for any $m \geq 1$ and all $x < \varepsilon'$ we have

$$\hbar \sum_{n=\varepsilon k}^{\varepsilon' k} e^{-\hbar^{-2}U(\hbar^2 n, x, w)} \alpha(\hbar^2 n, \hbar^2, w) = \int_{-\infty}^{\xi} q_{\hbar}(s) ds + \sum_{j=0}^{m-1} A_j \hbar^j + O(\hbar^m) \quad (\text{B.3})$$

where

$$A_0 = \frac{1}{2} q_{\hbar}(\xi) \quad (\text{B.4})$$

$$A_j = (-1)^{j-1} \frac{\beta_{j+1}}{(j+1)!} q_{\hbar}^{(j)}(\xi) \quad (\text{B.5})$$

which are bounded independent of \hbar . Moreover this is uniform as x varies in a compact subset of $[0, \varepsilon']$.

Proof. We can assume that U is the restriction of a function defined for all $\nu \geq 0$ (which for simplicity we also denote by U) with the same convexity properties. Similarly we assume α extends to a bounded function defined for all $\nu \geq 0$. Now using the convexity of U one sees that since $x \leq \varepsilon < \varepsilon'$ we have

$$p_{\hbar}(n) \leq O(e^{-c\hbar^{-2}} e^{-c\hbar^{-2}(\varepsilon' - \hbar^{-2}n)})$$

for some $c > 0$. Hence $\sum_{n > \hbar^{-2}\varepsilon} p_{\hbar}(n) = O(\hbar^{\infty})$ giving

$$S = \hbar \sum_{n \geq \hbar^{-2}\varepsilon} e^{-\hbar^{-2}U(\hbar^2 n, x, w)} \alpha(\hbar^2 n, \hbar^2, w) + O(\hbar^{\infty}).$$

Similarly one verifies that for $x \leq \varepsilon$ the derivatives of p_{\hbar} tend to zero as t tends to infinity. Hence the above Euler-Maclaurin formula (B.3) (applied with $a = \hbar^{-2}\varepsilon$) gives

$$\hbar^{-1} S = \int_{\hbar^{-2}\varepsilon}^{\infty} p_{\hbar}(t) dt + \frac{p_{\hbar}(\hbar^{-2}\varepsilon)}{2} - \sum_{j=2}^{m-1} \frac{\beta_{j+1}}{(j+1)!} p_{\hbar}^{(j)}(\hbar^{-2}\varepsilon) - \int_{\hbar^{-2}\varepsilon}^{\infty} \frac{\beta_m(\{1-t\})}{m!} p_{\hbar}^{(m)}(t) dt + O(\hbar^{\infty}).$$

Now by instant computation

$$q_{\hbar}^{(j)}(\xi) = (-\hbar)^{-j} p_{\hbar}^{(j)}(\hbar^{-2}\varepsilon).$$

On the other hand by a change of variables

$$\int_{\hbar^{-2}\varepsilon}^{\infty} p_{\hbar}(t) dt = \hbar^{-1} \int_{-\infty}^{\xi} q_{\hbar}(s) ds$$

and a similar change of variables shows that

$$\int_{\hbar^{-2}\varepsilon}^{\infty} \frac{\beta_m(\{1-t\})}{m!} p_{\hbar}^{(m)}(t) dt = O(\hbar^{m-1}).$$

Putting this together gives the statement, and the fact that the A_j are bounded as \hbar tends to zero follows as the derivatives of q_{\hbar} are bounded. \square

APPENDIX C. LAPLACE'S METHOD

We now give an account of Laplace's method [15] tailored to our requirements (see also [29]). We are concerned with obtaining the large- k asymptotic behaviour of integrals of the form

$$F_k(x) = \sqrt{\frac{k}{2\pi}} \int_{-\infty}^x e^{-kf(t)} \alpha(t) dt \quad (\text{C.1})$$

where α is smooth and has compact support, and f has the properties

- $f(t) \geq 0$;
- $f(0) = 0$, $f'(0) = 0$, $f''(0) = c > 0$;
- $f(x) \geq f_1 > 0$ for all $|x| > \delta$,

where $\delta > 0$ is some small number.

These properties ensure that if the support of α does not contain 0, then $F_k(x)$ is uniformly exponentially small in k , independent of x . So the interesting case is that $\text{supp}(\alpha) \ni 0$ and one expects the asymptotic expansion in negative powers of k to see only the jets of α and f at 0. The model problem is the integral

$$Z_k(x) = \sqrt{\frac{k}{2\pi}} \int_{-\infty}^x e^{-kt^2/2} dt. \quad (\text{C.2})$$

This is the case $f(t) = t^2/2$, which clearly verifies the above properties (even if the function ‘1’ replacing u in (C.1) does not have compact support).

Make the change of variables

$$y = \frac{x}{\hbar} \text{ and } s = \frac{t}{\hbar} \text{ and } \hbar = \frac{1}{\sqrt{k}} \quad (\text{C.3})$$

in (C.2) so

$$Z_k(\hbar y) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^y e^{-s^2/2} ds = \Phi(y) \quad (\text{C.4})$$

where Φ is the usual normal distribution function. In other words,

$$Z_k(x) = \Phi(x/\hbar). \quad (\text{C.5})$$

Note that if $x > 0$ is fixed, $x/\hbar \rightarrow +\infty$ as $\hbar \rightarrow 0$, so $Z_k(x) \rightarrow 1$ and in fact $Z_k(x) - 1$ is exponentially small in k for fixed x . Similarly, if $x < 0$ is fixed, $Z_k(x)$ is exponentially small in k . But (C.4) shows that this apparently discontinuous behaviour in x near 0 can be ‘smoothed’ by introducing a new variable x/\hbar , which corresponds to a real radial blow-up, as discussed in Appendix A.

Life is more interesting for general exponents f satisfying the above conditions (and for functions α not identically equal to 1). In that case, if $x > 0$, for example, $F_k(x)$ is exponentially close to

$$F_k(\infty) := \sqrt{\frac{k}{2\pi}} \int_{-\infty}^{\infty} e^{-kf(t)} \alpha(t) dt \quad (\text{C.6})$$

and the asymptotic expansion of this is given by Laplace’s method.

C.1. Incomplete Gaussian integrals. We change notation slightly and set

$$Z(x, \hbar; \alpha) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^x e^{-t^2/2\hbar^2} \alpha(t) dt. \quad (\text{C.7})$$

We assume that α is C^∞ and that derivatives of α are bounded,

$$\|\alpha\|_{C^r} \leq A_r \quad (\text{C.8})$$

for each $r \geq 0$.

For such a function, define

$$\delta_0 \alpha(t) = \frac{\alpha(t) - \alpha(0)}{t} \text{ for } t \neq 0, \quad \delta_0 \alpha(0) = \alpha'(0). \quad (\text{C.9})$$

Then $\delta_0 \alpha$ is smooth and all derivatives are bounded. We have the formula

$$\delta_0 \alpha(t) = \int_0^1 \alpha'(\lambda t) d\lambda. \quad (\text{C.10})$$

It follows that $\delta_0 \alpha$ is smooth for $|t| < 1$, say, with

$$\partial_t^n (\delta_0 \alpha) = \int_0^1 \lambda^n \alpha^{(n+1)}(\lambda t) d\lambda. \quad (\text{C.11})$$

In particular if $|t| < 1$, $\partial_t^n \delta_0 \alpha$ is bounded by $\sup |\alpha^{(n+1)}|$. On the other hand, (C.9) shows that $\delta_0 \alpha(t)$ decays as $|t| \rightarrow \infty$, and by differentiating this formula, the same is true of all derivatives.

Inserting the formula

$$\alpha(t) = \alpha(0) + t\delta_0 \alpha(t) \quad (\text{C.12})$$

into (C.7), getting

$$\begin{aligned} Z(x, \hbar; \alpha) &= \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^x e^{-t^2/2\hbar^2} (\alpha(0) + t\delta_0\alpha(t)) dt \\ &= \frac{1}{\sqrt{2\pi\hbar}} \alpha(0) \int_{-\infty}^x e^{-t^2/2\hbar^2} dt + \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^x t e^{-t^2/2\hbar^2} \delta_0\alpha(t) dt \end{aligned} \quad (\text{C.13})$$

$$= \alpha(0)\Phi(x/\hbar) - \frac{\hbar}{\sqrt{2\pi}} e^{-x^2/2\hbar^2} \delta_0\alpha(x) + \frac{\hbar^2}{\sqrt{2\pi\hbar}} \int_{-\infty}^x e^{-kt^2/2} [\delta_0\alpha]'(t) dt. \quad (\text{C.14})$$

Here we have used the notation

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt \text{ so } \Phi'(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2} \quad (\text{C.15})$$

respectively for the normal distribution function and the normal density function. The final line (C.14) was obtained by integration by parts and the formula:

$$t e^{-t^2/2\hbar^2} = -\hbar^2 \frac{d}{dt} e^{-t^2/2\hbar^2}. \quad (\text{C.16})$$

Now define

$$\mathcal{D}\alpha = \frac{d}{dt} \delta_0\alpha. \quad (\text{C.17})$$

Then (C.14) can be written

$$Z(x, \hbar; \alpha) = \alpha(0)\Phi(x/\hbar) + \hbar\delta_0\alpha(x)\Phi'(x/\hbar) + \hbar^2 Z(x, \hbar; \mathcal{D}\alpha) \quad (\text{C.18})$$

This forms the basis of the inductive step in obtaining a full asymptotic expansion of $Z(x, \hbar; \alpha)$:

Theorem C.1. *Suppose that $u(t)$ is smooth with all derivatives uniformly bounded on \mathbb{R} , and let*

$$Z(x, \hbar; \alpha) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^x e^{-kt^2/2} \alpha(t) dt. \quad (\text{C.19})$$

Then for each N , we have an expansion

$$Z(x, \hbar; \alpha) = \left\{ \sum_{j=0}^N \hbar^{2j} \mathcal{D}^j \alpha(0) \right\} \Phi(x/\hbar) + \left\{ \sum_{j=0}^N \hbar^{2j+1} \delta_0 \mathcal{D}^j \alpha(x) \right\} \Phi'(x/\hbar) + \hbar^{2N+2} \mathcal{R}_{N+1}(x, \hbar; \alpha), \quad (\text{C.20})$$

where the error term is given by

$$\mathcal{R}_{N+1}(x, \hbar; \alpha) = Z(x, \hbar, \mathcal{D}^{N+1}\alpha) \quad (\text{C.21})$$

which satisfies

$$|\mathcal{R}_{N+1}(x, \hbar; \alpha)| \leq B_N \|\alpha\|_{C^{2N+2}}. \quad (\text{C.22})$$

Proof. It is clear that (C.18) yields the case $N = 0$ of (C.20). It also gives the inductive step: if (C.20) holds for N , then replacing u by $\mathcal{D}^N \alpha$ implies (C.20) for $N + 1$. So it remains only to prove the estimate (C.22). For this, note that δ_0 behaves as a first-order differential operator: we have estimates of the form

$$\|\delta_0\alpha\|_{C^r} \leq B_r \|\alpha\|_{C^{r+1}} \quad (\text{C.23})$$

for each r , B_r being a constant independent of α . It follows that \mathcal{D} satisfies estimates of the form

$$\|\mathcal{D}\alpha\|_{C^r} \leq B'_r \|\alpha\|_{C^{r+2}} \quad (\text{C.24})$$

for each r .

To obtain the estimate of \mathcal{R}_{N+1} from this, use the change of variables $s = t/\hbar$ in the integral

$$\mathcal{R}_{N+1}(x, \hbar; \alpha) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^x e^{-t^2/2\hbar^2} \mathcal{D}^{N+1}\alpha(t) dt = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x/\hbar} e^{-s^2/2} \mathcal{D}^{N+1}\alpha(\hbar s) ds \quad (\text{C.25})$$

and the bound (C.24), iterated, to give

$$\sup |\mathcal{D}^{N+1}\alpha| \leq B''_{N+1} \|\alpha\|_{C^{2N+2}} \quad (\text{C.26})$$

for some constant B''_{N+1} . \square

C.2. More general exponentials. We now consider the changes needed to adapt this method to the more general exponents appearing in

$$F(x, \hbar; u) := \sqrt{\frac{k}{2\pi}} \int_{-\infty}^x e^{-kf(t)} \alpha(t) dt$$

as in (C.1). Our approach is quite standard: we deform f to its quadratic part and work out what additional terms this introduces. So consider the 1-parameter family of exponents

$$f_\lambda(t) = \frac{ct^2}{2} + \lambda q(t) \quad (\text{C.27})$$

where $q(t)$ is the third-order part of $f(t)$. Then $f_0(t)$ is a standard quadratic, to which the above analysis can be applied, and $f_1(t) = f(t)$ is the function we're really interested in. Set

$$H(\lambda) = H(\lambda; x, \hbar, \alpha) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^x e^{-kf_\lambda(t)} \alpha(t) dt. \quad (\text{C.28})$$

With all other parameters fixed, Taylor's theorem for H gives

$$\left| H(1) - \sum_{j=0}^p \frac{1}{j!} H^{(j)}(0) \right| \leq C \sup_{0 \leq \lambda \leq 1} H^{(p+1)}(\lambda). \quad (\text{C.29})$$

For simplicity take $c = 1$. Then for any j ,

$$H^{(j)}(0) = \frac{(-1)^j}{\sqrt{2\pi\hbar}^{2j+1}} \int_{-\infty}^x e^{-t^2/2\hbar^2} q(t)^j \alpha(t) dt \quad (\text{C.30})$$

and this integral, for fixed j , can be analyzed by the techniques of the previous section. We stress that the the third-order vanishing of q implies that the integrand in (C.30) vanishes to order $3j$ and so contributes terms of order \hbar^{3j} , leaving $H^{(j)}(0)$ of order \hbar^j . In the statements that follow it is clearest to distinguish cases according to the parity of j .

Theorem C.2. *Suppose that $j = 2\nu + 1$ is odd. Then there exist a sequence of smooth functions $\alpha_{i,j}$, $i = 0, \dots, 3\nu + 1$, such that*

$$H^{(2\nu+1)}(0) = -\frac{\hbar^{2\nu+1}}{\sqrt{2\pi}} \sum_{i=0}^{3\nu+1} \alpha_{i,j}(\hbar y) y^{2i} e^{-y^2/2} - \hbar^{2\nu+2} Z(x, \hbar; \mathcal{D}^{3\nu+2}(q^j \alpha)). \quad (\text{C.31})$$

Similarly, if $j = 2\nu \geq 2$ is even, then there exists a sequence of smooth functions $\alpha_{i,j}$, $i = 0, \dots, 3\nu - 1$, such that

$$H^{(2\nu)}(0) = \frac{1}{\sqrt{2\pi}} \left(\hbar^{2\nu} \sum_{i=0}^{3\nu-1} \alpha_{i,2\nu}(\hbar y) y^{2i+1} e^{-y^2/2} + \hbar^{2\nu+1} \delta_0 \mathcal{D}^{3\nu}(q^{2\nu} \alpha)(\hbar y) e^{-y^2/2} \right) + \hbar^{2\nu+2} Z(x, \hbar; \mathcal{D}^{3\nu+1}(q^{2\nu} \alpha)). \quad (\text{C.32})$$

More precisely, the $\alpha_{i,j}$ are defined as follows:

$$\alpha_{i,2\nu+1}(x) x^{2i} = \delta_0 \mathcal{D}^{3\nu+1-i}(q^{2\nu+1} \alpha) \quad (\text{C.33})$$

and

$$\alpha_{i,2\nu}(x) x^{2i+1} = \delta_0 \mathcal{D}^{3\nu-i}(q^{2\nu} \alpha). \quad (\text{C.34})$$

Remark C.3. Implicit in the definitions of $\alpha_{i,j}$ in (C.33) and (C.34) is that the right-hand sides of those equations are indeed divisible by x^{2i} or x^{2i+1} respectively so that $u_{i,j}$ is smooth.

Remark C.4. As a final generalization, observe that the above expansion still holds if f and u are allowed to depend smoothly on \hbar and also on auxiliary parameters. That is, suppose that W is a relatively compact subset of \mathbb{C}^m and $f = f(t, \hbar, w)$ is smooth satisfies the above assumptions uniformly for $\hbar \geq 0$ and $w \in W$ and $\alpha = \alpha(t, \hbar, w)$ is smooth. In fact by expanding $\alpha = \alpha_0 + \alpha_1 \hbar + \dots$ in powers of \hbar we may as well consider the case of $\alpha = \alpha_0$. Furthermore by convexity of f , we see that the integral is changed only by a factor of $O(k^{-\infty})$ if α_0 is replaced by $\chi \alpha_0$ where χ is a cut-off function that is identically 1 on an interval that is larger than the range of x of interest. Thus we may as well assume α_0 has compact support.

Say

$$f(t, \hbar, w) = ct^2/2 + q(t, \hbar, w)$$

where $c = c(\hbar, w) > 0$ and q has order 3 in t . Then Theorem C.2 gives an expansion of

$$F(x, \hbar, w) := \sqrt{\frac{k}{2\pi}} \int_{-\infty}^x e^{-\hbar^{-2}f(t, \hbar, w)} \alpha(t, \hbar, w) dt \quad (\text{C.35})$$

in powers of \hbar whose terms can be deduced from (C.20, C.28, C.31-C.34). Moreover it is clear that the error term is uniform over $w \in W$. Looking back at these expressions (e.g. (C.18)) one sees its leading order term is given by

$$F(x, \hbar, w) = \frac{\alpha(0, 0, w)}{\sqrt{c(0, w)}} \Phi\left(\sqrt{c(0, w)}y\right) + O(\hbar). \quad (\text{C.36})$$

C.3. Lifting to the Real Blow-up. The above formulae can be given a geometric flavour through the introduction of the real blow-up $[\mathbb{R}_x \times [0, \infty)_{\hbar}, (0, 0)]$. Letting $y = x/\hbar$ as before, we have coordinates (y, \hbar) on this blow-up that cover the interior of the exceptional divisor E .

Theorem C.5. *The function $F(x, \hbar, w)$ from (C.35) extends smoothly from the interior of B to all boundary faces.*

Proof. This is almost obvious from the formulae we've obtained. Indeed the form of Theorem C.1 and equations (C.31) and (C.32) show at once (setting $x = \hbar y$) that F extends smoothly to a neighbourhood of the interior of the exceptional divisor E .

The blow-up X has two corners C_{\pm} . Here C_+ is the intersection of E with the lift of the positive real axis. We can take $\eta := \hbar/x = 1/y$ and $x \geq 0$ as coordinates near C_+ and in these coordinates the expansion in Theorem C.1 has the form

$$\begin{aligned} Z(x, \hbar; \alpha) = & \left\{ \sum_{j=0}^N (\eta x)^{2j} \mathcal{D}^j \alpha(0) \right\} \Phi(1/\eta) + \left\{ \sum_{j=0}^N (\eta x)^{2j+1} \delta_0 \mathcal{D}^j \alpha(x) \right\} \Phi'(1/\eta) \\ & + (\eta x)^{2N+2} \mathcal{R}_{N+1}(x, \hbar; \alpha). \end{aligned} \quad (\text{C.37})$$

and because $\Phi(1/\eta)$ and $\Phi'(1/\eta)$ are smooth down to $\eta = 0$, this expression is clearly smooth for (η, x) small and non-negative.

Similarly, in these coordinates,

$$H^{(2l+1)}(0) = -\frac{(\eta x)^{2l}}{\sqrt{2\pi}} \sum_{i=0}^{3l+1} \alpha_{i,j}(x) \eta^{-2i} e^{-1/2\eta^2} + \hbar^{2l+1} Z(x, \hbar; \mathcal{D}^{3l+2} \alpha) \quad (\text{C.38})$$

and each term in the sum is of the form $x^{2l} \eta^{2l-2i} e^{-1/2\eta^2}$ ($i = 0, \dots, 3l+1$) which again is smooth for (η, x) small and non-negative. \square

C.4. Proofs. To prove Theorem C.2 we start with a lemma:

Lemma C.6. *Suppose that $g(t)$ is smooth, with all derivatives bounded, and vanishes to order m at 0: in other words,*

$$g^{(j)}(0) = 0 \text{ for all } j < m. \quad (\text{C.39})$$

Then $\delta_0 g$ vanishes to order $m-1$ and $\mathcal{D}g$ vanishes to order $m-2$ at 0.

Proof. The question is local to 0, so we can use the representation

$$\delta_0 g(t) = \int_0^1 \partial_t g(\lambda t, x) d\lambda \quad (\text{C.40})$$

Then by (C.11), we see at once that $(\delta_0 g)^{(j)}(0) = 0$ for all $j = 0, \dots, m-1$. This implies the corresponding vanishing for $\mathcal{D}g$. \square

Now we can move on to the proof of the Theorem. The proof is little more than using Theorem C.1 to expand $Z(x, \hbar; q^j \alpha)$, in combination with the information from the previous lemma.

More precisely, suppose that $j = 2\nu + 1$. Then the order of vanishing of $q^j \alpha$ is $m = 6\nu + 3 = 2(3\nu + 1) + 1$. It follows from the lemma that $\mathcal{D}^{3\nu+1-i}(q^j \alpha)$ vanishes to order $2i$ for $i = 0, \dots, 3\nu + 1$. In particular the $\alpha_{i,j}$ of (C.33) are well-defined smooth functions. Now apply Theorem C.1 with $N = 3\nu + 1$. Because $\mathcal{D}^{3\nu+1-i}(q^j \alpha)(0) = 0$, the coefficient of Φ in (C.20) is zero. Thus we are left with

$$Z(x, \hbar; q^j \alpha) = \frac{e^{-y^2/2}}{\sqrt{2\pi}} \sum_{i=0}^{3\nu+1} \hbar^{2i+1} \delta_0 \mathcal{D}^i u(q^j x) + \hbar^{6\nu+4} Z(x, \hbar; \mathcal{D}^{3\nu+2}(q^{2\nu+1} \alpha)). \quad (\text{C.41})$$

Now substitute (C.33) into (C.41) and write $x = \hbar y$ to get

$$Z(x, \hbar; q^j \alpha) = \frac{e^{-y^2/2}}{\sqrt{2\pi}} \sum_{i=0}^{3\nu+1} \hbar^{2i+1} \alpha_{3\nu+1-i, 2\nu+1}(\hbar y) (\hbar y)^{2(3\nu+1-i)} + \hbar^{6\nu+4} Z(x, \hbar; \mathcal{D}^{3\nu+2}(q^{2\nu+1} \alpha)). \quad (\text{C.42})$$

Since $H^{(2\nu+1)}(0) = -Z(x, \hbar; q^{2\nu+1} \alpha) / \hbar^{4\nu+2}$, (C.31) follows immediately from (C.42).

If $j = 2\nu$ is even, then the proof follows precisely the same lines. We apply Theorem C.1 with $N = 3\nu$ to expand $Z(x, \hbar; q^{2\nu} \alpha)$ and use the fact that $q^{2\nu} \alpha$ has a zero of order 6ν at 0. This means that the $\alpha_{i, 2\nu}$ of (C.34) are well-defined for $i = 0, \dots, 3\nu - 1$. We proceed as before, using $H^{(2\nu)}(0) = Z(x, \hbar; q^{2\nu} \alpha) / \hbar^{4\nu}$ to obtain (C.32).

C.5. Remainder term. It remains only to estimate the error term $H^{(p)}(\lambda)$ in (C.29). Of course

$$H^{(p)}(\lambda) = \frac{(-1)^p}{\sqrt{2\pi} \hbar^{2p+1}} \int_{-\infty}^x e^{-kf_\lambda(t)} q(t)^{p+1} \alpha(t) dt. \quad (\text{C.43})$$

On the assumption that

$$f_\lambda(t) \geq t^2/4 \quad (\text{C.44})$$

for $t \in \text{supp}(\alpha)$ and $0 \leq \lambda \leq 1$,

$$|H^{(p)}(\lambda)| \leq \frac{1}{\sqrt{2\pi} \hbar^{2p+2}} \int_{-\infty}^x e^{-t^2/4} |q(t)|^{p+1} |\alpha(t)| dt \leq \frac{1}{\sqrt{2\pi} \hbar^{2p}} \int_{-\infty}^x e^{-kt^2/4} Q^{p+1} |t^{3p+3} \alpha(t)| dt \quad (\text{C.45})$$

Making the change of variables $t = \hbar s$, $x = \hbar y$,

$$|H^{(p)}(\lambda)| \leq \frac{\hbar^{p+1}}{\sqrt{2\pi}} \int_{-\infty}^y e^{-s^2/4} Q^{p+1} |s^{3p+3} \alpha(\hbar s)| ds \quad (\text{C.46})$$

REFERENCES

- [1] R. Berman *Bergman kernels and equilibrium measures for ample line bundles* (2007) Preprint arXiv:0704.1640.
- [2] R. Berman *Bergman kernels for weighted polynomials and weighted equilibrium measures of \mathbb{C}^n* Indiana Univ. Math. J. 58 (2009), no. 4, 1921–1946.
- [3] R. Berman, B. Berndtsson and J. Sjöstrand *A direct approach to Bergman Kernel asymptotics for positive line bundles*, Ark. Mat. 46 no. 2 197–217 (2007).
- [4] B. Berndtsson *A remark on approximation on totally real sets* Complex analysis and digital geometry, 75–80, Acta Univ. Upsaliensis Skr. Uppsala Univ. C Organ. Hist., 86, Uppsala Universitet, Uppsala, 2009.
- [5] D. Catlin, *The Bergman kernel and a theorem of Tian*, Analysis and geometry in several complex variables (Katata, 1997), Trends Math., Birkhäuser Boston, Boston, MA, 1999, 1–23.

- [6] X. Dai, K. Liu and X. Ma *On the Asymptotic Expansion of Bergman Kernel* J. Diff. Geom. 72 (2006) 1–41
- [7] J.-P. Demailly *Complex Analytic and Differential Geometry* Version of June 21, 2012 (www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf)
- [8] H. Delin *Pointwise estimates for the weighted Bergman projection kernel in \mathbf{C}^n , using a weighted L^2 estimate for the $\bar{\partial}$ equation* Ann. Inst. Fourier (Grenoble) 48 (1998), no. 4, 967–997.
- [9] C. Fefferman, *The Bergman kernel and biholomorphic mappings of pseudoconvex domains*, Invent. Math. **26**, 1974, 1–65.
- [10] D. Grieser *Basics of the b-calculus*. Approaches to singular analysis (Berlin, 1999), 30–84.
- [11] A. Hassell, R. Mazzeo and R. Melrose *Analytic surgery and the accumulation of eigenvalues*, Commun. in Anal. and Geom. 3, 115-222, 1995.
- [12] H. Hedenmalm and N. Makarov *Quantum Hele-Shaw flow* Preprint arXiv:math/0411437
- [13] H. Hedenmalm and N. Makarov *Coulomb gas ensembles and Laplacian growth* Proc. Lond. Math. Soc. (3) 106 (2013), no. 4, 859–907.
- [14] C. Kiselman *Plurisubharmonic functions and their singularities* Complex Potential Theory, NATO ASI Series Volume 439, 1994, 273-323
- [15] P-S Laplace *Memoir on the probability of causes of events. Mémoires de Mathématique et de Physique, Tome Sixième.* (English translation by S. M. Stigler 1986. Statist. Sci., 364–378).
- [16] N. Lindholm *Sampling in weighted L^p spaces of entire functions in C^n and estimates of the Bergman kernel* J. Funct. Anal. 182 (2001), no. 2, 390–426.
- [17] R. Melrose *Differential analysis on manifolds with corners* In preparation, partially available at <http://math.mit.edu/~rbm/book.htm>.
- [18] R. Melrose *Star products and local line bundles* Ann. Inst. Fourier (Grenoble) 54 (2004), no. 5, 1581–1600, xvi, xxii.
- [19] X. Ma and G. Marinescu, *Exponential estimate for the asymptotics of Bergman kernels* Math. Ann. 362 (2015), no. 3-4, 1327–1347.
- [20] X. Ma and G. Marinescu, *Holomorphic Morse inequalities and Bergman kernels*, Progress in Mathematics, vol. 254, Birkhäuser Verlag, Basel, 2007.
- [21] F. Pokorny and M. Singer *Toric partial density functions and stability of toric varieties* To appear in Math. Ann. (2013)
- [22] R. Michael Range, *Holomorphic Functions and Integral Representations in Several Complex Variables*, Springer Graduate Texts in Mathematics, vol. 108, 1986.
- [23] J. Ross and D. Witt Nyström *Envelopes of plurisubharmonic metrics with prescribed singularities* arXiv:1210.2220
- [24] J. Ross and D. Witt Nyström *Analytic Test configurations and geodesic rays* To appear in Journal of Symplectic Geometry
- [25] J. Ross and R. P. Thomas *Weighted Bergman kernels on orbifolds* Journal of Differential Geometry. 88 (2011) p.87–108
- [26] Y. Rubinstein *Geometric Quantization and Dynamical Constructions on the Space of Kähler Metrics* PhD Thesis, MIT (2008)
- [27] B. Shiffman and S Zelditch *Random polynomials with prescribed Newton polytope* J. Amer. Math. Soc. 17 (2004), no. 1
- [28] J. Stalker *Complex analysis: Fundamentals of the classical theory of functions* Birkhäuser 1998.
- [29] M. Zworski *Semiclassical Analysis* Graduate Studies in Mathematics, vol. 138, American Mathematical Society, 2012
- [30] Gang Tian, *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differential Geom. **32** no. 1, 1990, 99–130.
- [31] Steve Zelditch, *Szegő kernels and a theorem of Tian*, Internat. Math. Res.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE
E-mail address: j.ross@dpmms.cam.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON
E-mail address: michael.singer@ucl.ac.uk