

On the convexity of some free boundaries

Antonio Greco

Dipartimento di Matematica e Informatica
Università di Cagliari
I 09124 Cagliari
Italy

Bernd Kawohl

Mathematisches Institut
Universität zu Köln
D 50923 Köln
Germany

July 7, 2009

Abstract

We demonstrate that a method of Colesanti and Salani, which compares solutions of elliptic differential equations to their quasiconcave envelopes, can be extended to derive convexity of free boundaries. As examples we present the so-called dam problem, a free boundary problem modelling pollution and a Bernoulli problem. Moreover we prove strict convexity of the wet region in the dam problem in arbitrary dimensions.

1 Introduction

In their paper [7] Colesanti and Salani study boundary value problems on convex rings, that is domains $\Omega = \Omega_0 \setminus \bar{\Omega}_1$ which are the difference of two open convex sets $\Omega_i \subset \mathbb{R}^d$ with $\Omega_1 \subset\subset \Omega_0$. In their simplest case $\Delta u = 0$ in Ω , while $u = 1$ on $\partial\Omega_1$ and $u = 0$ on $\partial\Omega_0$ are imposed as boundary conditions, but other differential equations such as $\Delta_p u = f(u)$ with f non-decreasing and $f(0) = 0$ are also considered. If we set $u \equiv 1$ on $\bar{\Omega}_1$, then u is well-defined everywhere in Ω_0 . It is known that the solutions to such problems have convex level sets $\Omega_t = \{x \in \Omega_0 : u(x) \geq t\}$ for $t \in (0, 1)$.

One method of proof is the Gabriel-Lewis method that has been thoroughly studied in the 1980's in [13], [24], [19] and again more recently in [16]. It amounts to showing that $u((x+y)/2) \geq \min\{u(x), u(y)\}$ for every $(x, y) \in \Omega_0^2$.

Another method, which is conceptually very elegant, was developed by Colesanti and Salani in [7] and further developed in [9]. They define the quasiconcave envelope u^* of u as the smallest quasiconcave function above u . In other words, the level sets of u^* are the convex hulls Ω_t^* of the level sets of u , and so by definition $u^* \geq u$. Then they derive a differential inequality for u^* . This inequality holds only in the sense of viscosity solutions, but it suffices to conclude, using an appropriate comparison principle, that $u^* \leq u$ in Ω , because $u^* = u$ on $\partial\Omega$.

Incidentally, the method of comparing u with a related function u^* to show that $u = u^*$ has been successfully exploited in other geometrical contexts. In [2] and [18] u^* is the convex envelope of u , i.e. the function whose graph is the convex hull of the graph of u , for which an interesting characterization has recently been given in [27]. In [29] u^* represents the starshaped envelope of u .

In the present paper we shall study free boundary problems, in which Ω_0 is chosen by the solution. Therefore its shape is not a-priori known to be convex. In replacing u by u^*

we replace a possibly nonconvex support of a function u by its convex hull and thus modify a free boundary in order to show that it has to be convex from the beginning. Ingredients of the proof are the differential inequalities of Colesanti and Salani and a Hopf-type boundary lemma.

2 Preliminaries, differential inequalities for u^*

In this section we consider a ring-shaped domain Ω as before, but we drop the assumption that Ω_0 is convex (this is what we want to prove). We let u be a function of class $C^1(\Omega) \cap C^0(\bar{\Omega}_0)$ satisfying $u = 0$ on $\Gamma = \partial\Omega_0$, $u \equiv 1$ in Ω_1 and $0 < u < 1$ in Ω . We suppose, further, that u is of class C^2 in a neighborhood of every $x \in \Omega$ such that $\nabla u(x) \neq 0$. This is the case, for instance, if u is a p -harmonic function, i.e., a solution of $\Delta_p u = 0$ in Ω [31]. The results also apply to the case when u is the solution to the dam problem discussed in Section 3 and Ω is its support (instead of being ring-shaped).

In the next sections we deal with solutions u to some free-boundary problems that can be compared with suitable one-dimensional supersolutions. To be more precise, let us consider a point $x \in \Omega$ and define $t = u(x)$. If the level set Ω_t has a supporting hyperplane π passing through x , then we construct a supersolution \bar{u} depending only on the distance from π and attaining the value t on π . Since $\nabla \bar{u} \neq 0$, we prove eventually that $\nabla u(x) \neq 0$. For the moment we take the following property as an assumption:

Nondegeneracy. If the level set Ω_t , for t satisfying $\inf_{\Omega} u < t < \sup_{\Omega} u$, has a supporting hyperplane passing through some $x \in \Omega_t$, then $\nabla u(x) \neq 0$.

Nondegeneracy implies the following minimum principle for u^* .

Proposition 1 (A minimum principle) *For every t as above and for every x interior to Ω_t^* we have $u^*(x) > t$.*

Proof. We have to exclude that $u^*(x) = t$. The argument is by contradiction. Suppose that $u^*(x_0) = t$ at some x_0 interior to Ω_t^* . Since $u^* \geq s$ in Ω_s^* , we have $x_0 \notin \Omega_s^*$ for $s > t$. The union $G_t = \cup_{s>t} \Omega_s^*$ is a convex set not containing x_0 . The difference $\Omega_t^* \setminus G_t$ contains x_0 , and has a nonempty interior because x_0 is interior to Ω_t^* and G_t is convex. By construction we have $u^*(x) = t$ for all $x \in \Omega_t^* \setminus G_t$. Furthermore, there must be a point $P \in \Omega \setminus \bar{G}_t$ such that $u(P) = t$. Indeed, if $u(x) < t$ at every $x \in \Omega \setminus \bar{G}_t$ then Ω_t is a subset of the convex set \bar{G}_t , which implies $\Omega_t^* \subset \bar{G}_t$, but this has been excluded. Since P is a local maximum for u , we have $\nabla u(P) = 0$.

Now if Ω is a ring-shaped domain as in the Introduction, we take P as far as possible from a fixed point $Q \in \Omega_1$, and let π be the hyperplane passing through P and orthogonal to the segment QP . Since π supports Ω_t and $\nabla u(P) = 0$, we reach a contradiction with nondegeneracy and the conclusion follows. If, instead, Ω and u are as in the dam problem, then by looking for the boundary points where $u \geq t$ we see that \bar{G}_t contains the convex hull of the four points $(0, 0)$, $(a, 0)$, $(0, H - \sqrt{2t})$, $(0, h - \sqrt{2t})$. Taking P as far as possible from the segment connecting the last two points, we get that the line π passing through P and parallel to the preceding segment supports Ω_t and by nondegeneracy the conclusion follows again.

We shall need the following useful representation of points $x \in \Omega^*$ (see [7, Proposition 3.1] for a similar result).

Proposition 2 (Representation) Take $x \in \Omega^*$ and define $t = u^*(x)$. If $\inf_{\Omega} u < t < \sup_{\Omega} u$ then x is a convex combination of suitable $x_1, \dots, x_k \in \Omega$, $k \leq d$, such that $u(x_i) = t$ and $\nabla u(x_i) \neq 0$ for $i = 1, \dots, k$. Furthermore, there exists a hyperplane π supporting both Ω_t and Ω_t^* and containing every x_i . The gradients $\nabla u(x_i)$ are orthogonal to π and point towards Ω_t^* .

Proof. By Carathéodory's theorem [23, 28], the point x is a convex combination of suitable x_1, \dots, x_{d+1} such that $u(x_i) \geq t$ for $i = 1, \dots, d+1$. Let us say that x_i is *relevant* if the corresponding coefficient λ_i is positive, *negligible* if $\lambda_i = 0$. By Proposition 1, we have $x \in \partial\Omega_t^*$. This implies that there exists a supporting hyperplane π containing all the relevant x_i , say x_1, \dots, x_k . In particular, we must have $k \leq d$. Since $\Omega_t \subset \Omega_t^*$, every relevant x_i lies on $\partial\Omega_t$, hence $u(x_i) = t$. By nondegeneracy, the gradients $\nabla u(x_i)$ do not vanish, are orthogonal to π and point towards Ω_t^* .

Denoting by n the normal to the hyperplane π pointing towards Ω_t^* , and letting $u_n = \partial u / \partial n$, let us finally prove a key result concerning the gradient and Hessian of u^* in the viscosity sense. By a *test function φ touching u^* from above at x* we mean, as usual, a function φ of class C^2 in a neighborhood of x such that $u^* - \varphi$ has a local maximum at x and $u^*(x) = \varphi(x)$. The last condition justifies the preceding terminology but is not essential.

Proposition 3 Take $x \in \Omega^*$ and define $t = u^*(x)$. If $\inf_{\Omega} u < t < \sup_{\Omega} u$ then for every unit vector v such that $v \cdot n \neq 0$, and for every sufficiently small r , the following estimate holds:

$$u^*(x + rv) \geq u^*(x) + rv \cdot n \left(\sum_{j=1}^k \lambda_j / u_n(x_j) \right)^{-1} + o(r). \quad (2.1)$$

Furthermore, any test function φ touching u^* from above at x satisfies

$$\nabla \varphi(x) = \left(\sum_{j=1}^k \lambda_j / u_n(x_j) \right)^{-1} n \quad (2.2)$$

$$D^2 \varphi(x) \geq \left(\sum_{j=1}^k \lambda_j / u_n(x_j) \right)^{-3} \sum_{j=1}^k \lambda_j D^2 u(x_j) / u_n^3(x_j). \quad (2.3)$$

Inequalities similar to (2.3) are proved in [6, ineq. (8)] and in [25, ineq. (50)]. When we presented our results in Firenze on Dec. 12, 2008, Paolo Salani kindly provided us with a copy of [5] and pointed out that our Proposition 3 could also be derived from Prop. 2.2 in [5] in a way similar to the derivation of their Theorem 3.1. For the reader's convenience we give now our original proof.

Proof. By Proposition 2, the point x is a convex combination $x = \sum_{i=1}^k \lambda_i x_i$ of suitable $x_i \in \Omega$ such that $u(x_i) = t$ for all i . Those points belong to a supporting hyperplane π of Ω_t whose normal, directed towards Ω_t , is denoted by n . Let us fix a unit vector $v \in \mathbb{R}^d$ satisfying $v \cdot n \neq 0$. By the implicit function theorem, there exist smooth functions $\rho_i(r)$, $i = 1, \dots, k$, and $s(r)$, defined in a neighborhood of $r = 0$, such that $\rho_i(0) = 0$ for $i = 1, \dots, k$, $s(0) = t$ and

$$u(x_i + \rho_i(r)v) = s(r) \text{ for } i = 1, \dots, k \quad (2.4)$$

$$\sum_{i=1}^k \lambda_i \rho_i(r) = r. \quad (2.5)$$

Indeed, the Jacobian matrix of the preceding system at $\rho_i = 0$ and $s = t$ is

$$J = \begin{pmatrix} u_v(x_1) & \dots & 0 & -1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & u_v(x_k) & -1 \\ \lambda_1 & \dots & \lambda_k & 0 \end{pmatrix}$$

where $u_v = \partial u / \partial v$. Of course, the first k rows of J are linearly independent. Furthermore, the last row is independent from the preceding ones because in order to write this as a linear combination of those we must multiply the i -th row by $\lambda_i / u_v(x_i)$ and then sum over i . Looking at the last component we should obtain $\sum_i \lambda_i / u_v(x_i) = 0$, but this is impossible because the sign of $u_v(x_i)$ is independent from i and the coefficients λ_i are positive. Hence the implicit function theorem applies, as claimed.

Observe that $u^*(x_i + \rho_i(r)v) \geq u(x_i + \rho_i(r)v) = s(r)$ for each i . Since $x + rv = \sum_i \lambda_i (x_i + \rho_i(r)v)$, and since u^* is quasiconcave, we have $u^*(x + rv) \geq s(r)$. Hence any test function φ touching u^* from above at x satisfies $\varphi(x + rv) \geq s(r)$. Let us compute $s'(0)$ and $s''(0)$. By differentiating (2.4)-(2.5) we get

$$u_v(x_i + \rho_i(r)v) \rho_i'(r) = s'(r) \text{ for } i = 1, \dots, k \quad (2.6)$$

$$\sum_{i=1}^k \lambda_i \rho_i'(r) = 1. \quad (2.7)$$

A further differentiation at $r = 0$ yields:

$$u_{vv}(x_i) (\rho_i'(0))^2 + u_v(x_i) \rho_i''(0) = s''(0) \text{ for } i = 1, \dots, k \quad (2.8)$$

$$\sum_{i=1}^k \lambda_i \rho_i''(0) = 0 \quad (2.9)$$

where $u_{vv} = \partial^2 u / \partial v^2$. From (2.6)-(2.7) we obtain, at $r = 0$

$$s'(0) = \left(\sum_{j=1}^k \lambda_j / u_v(x_j) \right)^{-1} \quad (2.10)$$

$$\rho_i'(0) = \left(u_v(x_i) \sum_{j=1}^k \lambda_j / u_v(x_j) \right)^{-1}. \quad (2.11)$$

Using the last expression to replace $\rho_i'(0)$ in (2.8) we get

$$\rho_i''(0) = (u_v(x_i))^{-1} s''(0) - \frac{u_{vv}(x_i)}{(u_v(x_i))^3} \left(\sum_{j=1}^k \lambda_j / u_v(x_j) \right)^{-2}$$

By plugging this into (2.9), and since $u_v(x_j) = u_n(x_j) n \cdot v$, we finally arrive at

$$s''(0) = \left(\sum_{j=1}^k \lambda_j / u_n(x_j) \right)^{-3} \sum_{i=1}^k \lambda_i u_{vv}(x_i) / u_n^3(x_i). \quad (2.12)$$

From (2.10) rewritten as $s'(0) = (\sum_j \lambda_j / u_n(x_j))^{-1} n \cdot v$ we obtain (2.1), and (2.2) follows. By (2.12), instead, we get (2.3). The fact that (2.10) and (2.12) have been derived for $n \cdot v \neq 0$ makes no problem since φ is regular by definition.

Let us point out some consequences of the preceding result that were found in [7] by means of a different argument. Inequality (2.3) implies

$$\begin{aligned}\varphi_{nn}(x) &\geq \left(\sum_{j=1}^d \lambda_j / u_n(x_j) \right)^{-3} \sum_{j=1}^d \lambda_j u_{nn}(x_j) / u_n^3(x_j) \\ \Delta_\pi \varphi &\geq \left(\sum_{j=1}^d \lambda_j / u_n(x_j) \right)^{-3} \sum_{j=1}^d \lambda_j \Delta_\pi u(x_j) / u_n^3(x_j)\end{aligned}$$

where $\Delta_\pi \varphi = \Delta \varphi - \varphi_{nn}$. Furthermore, since the p -Laplace operator $\Delta_p u$ can be represented as $\Delta_p u = |\nabla u|^{p-2} [(p-1)u_{nn} + \Delta_\pi u]$ provided that $\nabla u \neq 0$, and since u is assumed to be nondegenerate in the sense defined above, the quasiconcave rearrangement u^* satisfies in the viscosity sense the following differential inequality:

$$\Delta_p u^*(x) \geq \left(\sum_{j=1}^d \lambda_j / u_n(x_j) \right)^{-(p+1)} \sum_{j=1}^d \lambda_j \Delta_p u(x_j) / (u_n(x_j))^{p+1}.$$

Consequently, if u is a nondegenerate, weak solution of $\Delta_p u = f(u)$ in Ω with $f \geq 0$ then

$$\Delta_p u^*(x) \geq f(u^*(x)) \quad \text{in the sense of viscosity solutions.} \quad (2.13)$$

Several comparison principles for viscosity solutions are found in the literature: see, for instance, [3, Theorem 1], [8, Theorem 3.3], [26, Theorem 1.1]. However, since we shall compare u and u^* in a subset $G \subset \Omega$ where $\nabla u \neq 0$, and u satisfies $\Delta_p u = f(u)$ in the classical sense, a very specific result is enough to proceed. For completeness, we prove a suitable statement hereafter.

Proposition 4 (Comparison principle) *Let G be a bounded, open subset of \mathbb{R}^d , and let u be a function of class $C^2(\bar{G})$ satisfying $\inf_G |\nabla u| > 0$. Thus, the function $\Delta_p u(x)$ is defined in the classical sense in \bar{G} . If $v \in C^0(\bar{G})$ is a viscosity solution of $\Delta_p v \geq \Delta_p u$ in G , then*

$$\sup_G (v - u) = \max_{\partial G} (v - u).$$

Proof. Suppose, by contradiction, that the function $w = v - u$ attains an interior maximum $M > \max_{\partial G} w$. Define $\phi = u - c e^{\gamma x_1}$, where x_1 stands for the first component of the variable x , and c, γ are real numbers. For every $\gamma \in \mathbb{R}$ we choose a sufficiently small $c = c(\gamma) > 0$ such that (i) the function $w(x) + c(\gamma) e^{\gamma x_1} = v - \phi$ attains its maximum at some $x(\gamma) \in G$; (ii) $\sup_G |\nabla u - \nabla \phi| < \frac{1}{2} \inf_G |\nabla u| =: r$. In particular, $\nabla \phi(x) \notin B(0, r)$ for all x, γ . Since ϕ is a *test function* at the point $x(\gamma)$ we may write $a^{ij}(\nabla u(x(\gamma))) u_{ij}(x(\gamma)) \leq a^{ij}(\nabla \phi(x(\gamma))) \phi_{ij}(x(\gamma))$, where $a^{ij}(\xi) = \delta^{ij} |\xi|^{p-2} + (p-2) |\xi|^{p-4} \xi_i \xi_j$ for $\xi \in \mathbb{R}^d \setminus \{0\}$ and the summation convention is used. By plugging the term $\phi_{ij}(x(\gamma)) = u_{ij}(x(\gamma)) - c(\gamma) \gamma^2 e^{\gamma x_1(\gamma)} \delta^{i1} \delta^{j1}$ into the preceding inequality, we obtain

$$[a^{ij}(\nabla u(x(\gamma))) - a^{ij}(\nabla \phi(x(\gamma)))] u_{ij}(x(\gamma)) + c(\gamma) \gamma^2 e^{\gamma x_1(\gamma)} a^{11}(\nabla \phi(x(\gamma))) \leq 0.$$

By the mean value theorem, we may write

$$a^{ij}(\nabla u(x(\gamma))) - a^{ij}(\nabla \phi(x(\gamma))) = c(\gamma) \gamma a_1^{ij}(\xi(\gamma)) e^{\gamma x_1(\gamma)}$$

where $a_1^{ij} = \partial a^{ij} / \partial \xi_1$ and $\xi(\gamma)$ lies on the segment joining ∇u to $\nabla \phi$ in \mathbb{R}^d . With this replacement, provided that $\gamma \neq 0$, we arrive at

$$a_1^{ij}(\xi(\gamma)) u_{ij}(x(\gamma)) + \gamma a^{11}(\nabla \phi(x(\gamma))) \leq 0. \quad (2.14)$$

Now let $\gamma \rightarrow +\infty$. By construction, $\xi(\gamma)$ and $\nabla\phi(x(\gamma))$ stay bounded and far from the origin. Since $u \in C^2(\bar{\Omega})$, the term $a_1^{ij}(\xi(\gamma))u_{ij}(x(\gamma))$ is also bounded and the (positive) coefficient $a^{11}(\nabla\phi(x(\gamma)))$ keeps far from zero. Hence, the left-hand side in (2.14) tends to $+\infty$. This contradiction shows that the comparison principle holds.

In order to compare u and u^* on the free boundary Γ we use the following statement, which is also needed to prove the existence of the set G of Proposition 4.

Proposition 5 *Define $w(x) = u^*(x) - u(x)$ for $x \in \bar{\Omega}$. The function w cannot have a positive maximum M at $x_0 \in \Omega \cup \Gamma$ if $\nabla u(x_0) = 0$.*

Proof. Assume that a positive maximum M is attained at such an x_0 . Then, $u^*(x_0) = u(x_0) + M > 0$. Furthermore, since $u < 1$ in $\Omega \cup \Gamma$, we also get $u^*(x_0) < 1$ and Proposition 3 applies. By (2.1) we have $u^*(x_0 + rn) \geq u^*(x_0) + \varepsilon r$ for a convenient $\varepsilon > 0$ and for all sufficiently small $r > 0$. Since $\nabla u(x_0) = 0$, we also have $u(x_0 + rn) = u(x_0) + o(r)$. By subtracting the last equality from the preceding inequality we obtain $w(x_0 + rn) \geq M + \varepsilon r - o(r)$, a contradiction.

3 The dam problem

The flow of groundwater through a rectangular dam, which is impervious at the bottom, leads after the Baiocchi transformation to the following free boundary problem. Let $D = (0, a) \times (0, b)$ be a rectangular domain, the cross section of the dam, and let H and h denote the height of water left and right of the dam, $0 < h < H < b$. We look for a function u that satisfies Dirichlet conditions $u = g$ on ∂D and the differential equation $\Delta u = 1$ on its support $\Omega = \{x \in D : u(x) > 0\}$. Physically, Ω represents the wet region in the dam. The datum g is specified as follows: $g(x_1, 0) = \frac{1}{2a}[aH^2 - x_1(H^2 - h^2)]$ and $g(x_1, b) = 0$ for $x_1 \in [0, a]$; $g(0, x_2) = \frac{1}{2}(H - x_2)(H - x_2)^+$ and $g(a, x_2) = \frac{1}{2}(h - x_2)(h - x_2)^+$ for $x_2 \in (0, b)$. Another way to characterize the solution is the obstacle problem of minimizing

$$J_1(v) := \int_D \left\{ \frac{|\nabla v|^2}{2} + v \right\} dx$$

on $K_1 := \{v \in W^{1,2}(D) : v \geq 0 \text{ in } D \text{ and } v = g \text{ on } \partial D\}$. It is easy to see that this problem has a unique solution u with the required properties. Moreover, u_x and u_y are negative in Ω and the free boundary $\partial\Omega \cap D$ can be represented as a smooth function of x .

Theorem 6 *The solution u of the dam problem is quasiconcave. Therefore all of its level sets are convex.*

For the proof we compare u^* , the quasiconcave envelope of u , with u on Ω . Since $u_x, u_y < 0$ in Ω , the nondegeneracy assumption of Section 2 is satisfied. By Proposition 3 we know that $\Delta u^* \geq \Delta u$ in Ω in the sense of viscosity solutions, so that $u^* - u$ attains any positive maximum on $\partial\Omega$. However, by definition $u^* = u$ on ∂D , so that a positive maximum can only occur at a point x_0 on the free boundary $\Gamma = \partial\Omega \cap D$. But this is impossible by Proposition 5, because $\nabla u = 0$ on Γ . Hence $u^* > u$ cannot hold anywhere in Ω . Since $u^* \geq u$, the equality follows.

Note that this method of proof can carry over to higher dimensions, and that it proves the convexity of all level sets, not just the support. In three dimensions one can think of the dam as a wall that encloses a pool.

In [12] and [10] one can find proofs that the support is even strictly convex in the sense that Γ has positive curvature. However, the arguments there work only in two dimensions. The (less sharp) fact that *the free boundary Γ contains no segments* can be easily proved as follows. Assume, contrary to the claim, that the free boundary Γ contains a segment S . After a suitable change of coordinates we get that S lies on $\{x_1 = 0\}$ and $\Omega \subset \{x_1 > 0\}$. The proof aims to show that the solution u coincides with $v = x_1^2/2$ in Ω , which is impossible because the boundary condition on the impervious bottom of the dam is not satisfied. Indeed, u should be linear along the bottom. However, since $h < H$, we see that v is quadratic there. To prove that $u \equiv v$, consider the harmonic function $w = u - v$ in Ω . Since $w = |\nabla w| = 0$ on S , it follows that w is identically zero [14, Problem 2.2] and the proof is complete.

4 The pollution problem

Imagine a chemical seeping through the wall of a full tank that occupies Ω_1 into an ambient medium, where it is consumed by a reactant. Stationary problems of this nature can be modelled as variational problems. Minimize

$$J_2(v) := \int_{\mathbb{R}^d} \left\{ \frac{|\nabla v|^p}{p} + F(v) \right\} dx$$

on $K_2 := \{v \in W^{1,p}(\mathbb{R}^d) : v \equiv 1 \text{ on } \overline{\Omega}_1\}$. Here $F : \mathbb{R} \rightarrow \mathbb{R}_0^+$ is a convex, monotone nondecreasing function of class $C^{1,\alpha}$ with $F(s) = 0$ for $s \leq 0$. The minimizing function u represents the dimensionless concentration of the chemical. If

$$\int_0^1 F(t)^{-1/p} dt < \infty$$

then the absorption term in the Euler equation $-\Delta_p u = -f(u)$, where $f = F'$, becomes so strong for small u that a minimizer has finite support. Let us call this the pollution problem.

Theorem 7 *If Ω_1 is convex then u is quasiconcave. Therefore the support of u and all of its level sets are convex.*

This theorem was derived in [21] for $p = 2$, but here we give a different proof that applies to any $p \in (1, \infty)$. The existence of a unique weak solution follows from the direct methods in the calculus of variations. By standard comparison results with a radially symmetric situation one can show that solutions have compact support and that $u(x) \in [0, 1)$ outside Ω_1 . Moreover, since $f(u)$ is bounded, u is of class $C^{1,\alpha}$ according to standard regularity results from [31] and [11], but then $f(u(x))$ is Hölder continuous everywhere outside Ω_1 . Only for $p = 2$ does this imply $u \in C^{2,\beta}(\mathbb{R} \setminus \overline{\Omega}_1)$. In fact, for $p \neq 2$ the p -Laplacian operator degenerates in points where $\nabla u = 0$, and we expect this to happen on the free boundary $\partial\Omega_0$, but not in $\Omega = \{x : u(x) \in (0, 1)\}$. How can we prove that $|\nabla u| > 0$ in Ω ?

That the level sets of u are starshaped with respect to any point in Ω_1 follows from [20] or [29]. Without loss of generality suppose that $0 \in \Omega_1$. Then we know that radial derivatives of u are nonpositive, but not negative as desired. A way out of this difficulty is the observation that we do not need ∇u to be nonzero everywhere, but only in special points x_i that span nontrivial (and convex at most $(d - 1)$ -dimensional) components of $\partial\Omega_t^* \setminus \partial\Omega_t$. This is where the inequalities of Proposition 3 are exploited, and this goal motivates the following step in the proof.

Lemma 8 (Nondegeneracy) *Let $x_0 \in \Omega$, and define $t = u(x_0)$. If there exists a hyperplane π passing through x_0 and supporting Ω_t , then $\nabla u(x_0)$ does not vanish, is orthogonal to π and points towards Ω_t .*

Proof. The claim follows from the weak comparison principle, after having constructed a suitable supersolution. Without loss of generality, we may assume $x_0 = 0$, $\pi = \{x_d = 0\}$, and $\Omega_t \subset \{x_d \geq 0\}$. Since Ω_0 is bounded, we have $\Omega_0 \subset \{x_d > \mu\}$ for some negative μ . The function $\bar{u}(x_1, \dots, x_d) = t(1 - x_d/\mu)$ satisfies $\Delta_p \bar{u} = 0$ in \mathbb{R}^d , $\bar{u} = t$ on π , $\bar{u} \geq 0$ on $\partial\Omega_0$. By the weak comparison principle we get from $f(u) \geq 0$ that $u \leq \bar{u}$ in $\Omega \cap \{x_d < 0\}$ and the conclusion follows.

Now we are in a position to give a **proof of Theorem 7**. By definition $u^* \geq u$, and since f is nondecreasing, by (2.13) we have $\Delta_p u^* \geq \Delta_p u$. Let us first realize that any positive maximum M of $w(x) = u^*(x) - u(x)$ over $\bar{\Omega}$ cannot occur on the free boundary $\partial\Omega_0$. It is well known that $u \in C^{1,\alpha}$ through the free boundary (see, for instance, [31]). Hence, $\nabla u = 0$ on $\partial\Omega_0$ and we may apply Proposition 5. This prevents w from having a positive maximum on $\partial\Omega_0$, and therefore any positive maximum of $w(x) = u^*(x) - u(x)$ over $\bar{\Omega}$ must occur at a point $x_0 \in \Omega$. Now we can no longer argue as in the proof of Theorem 6, because the differential equation for u is nonlinear. Hence, we proceed as follows. If $M = \sup_{\Omega} w > 0$, then we set $m := \max_{\partial\Omega} \bar{w}$ and observe that there exists $c \in (m, M)$ such that ∇u does not vanish in the closure \bar{G} of the set $G = \{x \in \Omega : w(x) > c\}$. For, if this were not the case, we would have a sequence $c_n \nearrow M^-$ and a sequence $x_n \in \Omega$ such that $w(x_n) > c_n$ and $\nabla u(x_n) = 0$. Passing to the limit we would find $x_0 \in \Omega$ such that $w(x_0) = M$ and $\nabla u(x_0) = 0$, against Proposition 5. Hence, a set G having the mentioned properties does exist. Of course, the supremum of w in G equals M , and is attained in the interior and not on the boundary (where $w = c$). But this contradicts Proposition 4 where we take $v = u^*$. Hence Ω_0 is convex and $u^* \equiv u$ in Ω , i.e. u is quasiconcave.

Remark 9 *Since $u = u^*$, a further application of (2.1) shows that $x \cdot \nabla u(x) < 0$ in all of Ω . This sharpens the conclusion in [20] and [29, Theorem 5.1], asserting (in case $u \in C^{1,\alpha}$) that $x \cdot \nabla u(x) \leq 0$ in Ω .*

5 Bernoulli's problem

Bernoulli's problem describes the outcome of a galvanization process. A body Ω_1 in a galvanization-bath is covered by another substance. The galvanized body occupies a domain Ω_0 . Mathematically this is modelled as follows. Minimize

$$J_3(v) := \int_{\mathbb{R}^d} \left\{ \frac{|\nabla v|^p}{p} + \lambda \chi_{\{v>0\}} \right\} dx$$

on $K_2 := \{v \in W^{1,p}(\mathbb{R}^d) : v \equiv 1 \text{ on } \bar{\Omega}_1\}$. Here λ is a positive constant and χ_A the characteristic function of a set A .

Theorem 10 *The solution w of Bernoulli's problem is quasiconcave. Therefore its support and all of its level sets are convex.*

Several proofs of this result are known for various p and n , see e.g. [4], [22], [1], [15], [17]. The following proof is based on a strategy similar to the one used in Section 4. Define $\Omega = \Omega_0 \setminus \bar{\Omega}_1$ and $\Omega_t = \{x \in \mathbb{R}^d : u(x) \geq t\}$, as before. Recall that if $\partial\Omega_0$ is sufficiently smooth, then $|\nabla u|$ attains the constant value $a = (p\lambda/(p-1))^{1/p}$ there. Let us start with a key observation:

Lemma 11 (Nondegeneracy) *Let x_0 be a point of Ω such that $u(x_0) \in (0, 1)$, and define $t = u(x_0)$. If there exists a hyperplane π passing through x_0 and supporting Ω_t , then $|\nabla u(x_0)| > a$.*

Proof. Let n be the normal to π directed towards Ω_t . Consider the affine (and p -harmonic) function $z(x) = t + a(x - x_0) \cdot n(x_0)$. The following comparison argument shows that $\Omega_0 \subset \{x \in \mathbb{R}^d : z(x) > 0\}$. Denote by $u_{R,t}$ the (radial) minimizer of the same functional under the constraint $u_{R,t} = t$ in the closed ball \overline{B}_R . If $\Omega_t \subset \overline{B}_R$, then we have $u \leq u_{R,t}$ outside Ω_t . Now let $R \rightarrow +\infty$ while the center of B_R also goes to infinity, in such a way that $\partial B_R \rightarrow \pi$. Since the function $u_{R,t}$ is easily found by integration, we see that $u_{R,t} \rightarrow z$. Hence we have $\Omega_0 \subset \{x \in \mathbb{R}^d : z(x) > 0\}$, as claimed. Furthermore, we have $u \leq z$ in $D = \{x \in \Omega : 0 < z(x) < t\}$ and $u(x_0) = z(x_0)$, which in turn implies $|\nabla u(x_0)| \geq a$. Consequently, u is a classical solution of the nondegenerate elliptic quasilinear equation $\Delta_p u = 0$ in a neighborhood of x_0 . The difference $w = u - z$ satisfies therefore a linear elliptic inequality of the form $Lw \geq 0$ near x_0 (see [14, Theorem 10.1]). Observe that the domain D satisfies an interior sphere condition at x_0 , hence we may apply Hopf's lemma and deduce the strict inequality $|\nabla u(x_0)| > a$.

Combining Proposition 2 and Lemma 11 we obtain:

Lemma 12 (Bound from below for u^*) *Let x_0 be a point of Ω^* such that $u^*(x_0) = t \in (0, 1)$. There exist positive numbers ε, ρ and a unit vector n such that the ball $\mathcal{B} = B(x_0 + \rho n, \rho)$ centered at $x_0 + \rho n$ and of radius ρ is included in Ω_t^* , and the restriction of u^* to \mathcal{B} is bounded from below as follows: $u^*(x) > t + (a + \varepsilon)n \cdot (x - x_0)$ for $x \in \mathcal{B}$.*

Proof. Let us apply Proposition 2 at the point $x = x_0$. Let π and $x_i, i = 1, \dots, k$ have the meaning given there, and let n be the normal to π pointing towards Ω_t^* . By Lemma 11 there exists $\varepsilon > 0$ so small that $a + \varepsilon < |\nabla u(x_i)|$ for all i . Denoting by η a variable in \mathbb{R}^d , and by the implicit function theorem, for each i there exists a neighborhood U_i of the origin such that the function $\varphi_i(\eta) = u(x_i + \eta) - t - (a + \varepsilon)n \cdot \eta$ vanishes along a regular hypersurface $\Gamma_i \subset U_i$ dividing U_i into two parts. Note that the hypersurfaces Γ_i are orthogonal to n at the origin. Since the index i ranges in a finite set, there exists a sufficiently small radius $\rho > 0$ such that the ball $B(\rho n, \rho)$ is contained in the set $U_i \cap \{\varphi_i(\eta) > 0\}$ for every i . This may be rewritten as follows:

$$u(x_i + \eta) > t + (a + \varepsilon)n \cdot \eta \quad \text{for } \eta \in B(\rho n, \rho). \quad (5.1)$$

Since $u^* \geq u$ by definition, we obtain $u^*(x_i + \eta) > t + (a + \varepsilon)n \cdot \eta$. Since $x_0 + \eta$ is a convex combination of the points $x_i + \eta$, and since u^* is quasiconcave, it follows that $u^*(x_0 + \eta) > t + (a + \varepsilon)n \cdot \eta$ in $B(\rho n, \rho) + \rho n$, which is equivalent to the claim.

As a consequence of Lemma 12, the difference $w = u^* - u$ cannot have a positive maximum at any $x_0 \in \partial\Omega_0$ provided that $|\nabla u(x_0)| = a$ in the classical sense. The argument is the same as in the proof of Proposition 5. Since the last equality does not hold, in general, we proceed as follows.

Lemma 13 *Define $w(x) = u^*(x) - u(x)$ for $x \in \overline{\Omega}$. The function w cannot have a positive maximum at any $x_0 \in \partial\Omega_0$.*

Proof. Suppose, contrary to the claim, that w attains a positive maximum at some $x_0 \in \partial\Omega_0$. The first part of the proof aims to deduce from this assumption an estimate of u

near x_0 , namely (5.3) below. Since $u(x_0) = 0$ and $w(x_0) > 0$, it follows that $u^*(x_0) > 0$. Clearly $u^*(x_0) < 1$ and therefore we may apply Lemma 12. Define $t = u^*(x_0)$ as in that lemma. Before proceeding further, note that the interior of Ω_t^* is included in Ω_0 . Indeed, by assumption we have

$$u^*(x) - u(x) \leq t \text{ in } \Omega_0, \quad (5.2)$$

hence $u^*(x) \leq t$ on $\partial\Omega_0$; furthermore, by Proposition 1 we see that $u^*(x) > t$ for x interior to Ω_t^* . Hence the interior of Ω_t^* does not intersect $\partial\Omega_0$, but since it does intersect Ω_0 , it must be included in Ω_0 , as claimed. Consequently, the ball $\mathcal{B} = B(x_0 + \rho n, \rho)$ of Lemma 12 is also included in Ω_0 , and by (5.2) we may write

$$u(x) > (a + \varepsilon)n \cdot (x - x_0) \text{ in } \mathcal{B}. \quad (5.3)$$

This inequality tells us, in a somewhat weak form, that $|\nabla u(x_0)| > a$, although $\nabla u(x_0)$ may not exist in the classical sense. In order to reach a contradiction, let us construct a radial minimizer $u_{R,s}$ having \mathcal{B} as its support and attaining a suitable constant value s in a concentric ball $B_R = B(x_0 + \rho n, R) \subset \mathcal{B}$ so that $u(x) > s$ in the closure $\overline{B_R}$. Of course, $u_{R,s}$ can be seen as a function of $r = |x - x_0 - \rho n|$. By applying the divergence theorem to the equation $\Delta_p u_{R,s} = 0$ it follows that $u_{R,s}(r)$ can be constructed as the solution to the first-order initial-value problem

$$\begin{aligned} r^{d-1} (-u'_{R,s}(r))^{p-1} &= \rho^{d-1} a^{p-1}, \quad r \in (R, \rho) \\ u_{R,s}(\rho) &= 0 \end{aligned}$$

where $u'_{R,s} = du_{R,s}/dr$. Note that $u'_{R,s}(\rho) = -a$. Moreover, the value of s is related to R by $s = u_{R,s}(R)$. By choosing R close enough to ρ (or equivalently by choosing s small enough) and taking (5.3) into account, we get that $u(x) > s$ in $\overline{B_R}$, as desired. Now we translate the function $u_{R,s}$ by some small $\sigma > 0$ in the direction of $-n$, i.e., we pass from $u_{R,s}(x)$ to $v(x) = u_{R,s}(x + \sigma n)$. The support of v is clearly $\mathcal{B} - \sigma n$, and the value s is attained in the ball $B_R - \sigma n$. By choosing σ small enough, we get that $u(x) > s$ on the last ball, and x_0 lies in the annulus $(\mathcal{B} \setminus \overline{B_R}) - \sigma n$. Let us see why this leads to a contradiction. Denote by S the set of all $x \in \mathcal{B} - \sigma n$ such that $v(x) > u(x)$. The set S is open, and nonempty because it contains x_0 . The set $S_0 = S \cap \Omega_0$ is also nonempty because the point x_0 is interior to S and belongs to $\partial\Omega_0$. Since $u \equiv 0$ outside Ω_0 , we have $S_0 \neq S$. We may decompose $J_3(v)$ as follows:

$$J_3(v) = \int_{(\mathcal{B} - \sigma n) \setminus S} \left\{ \frac{|\nabla v|^p}{p} + \lambda \right\} dx + \int_S \left\{ \frac{|\nabla v|^p}{p} + \lambda \right\} dx. \quad (5.4)$$

Now we compare the last integral with the one obtained by replacing the domain of integration S with S_0 and the function v with u . More precisely, we claim that

$$\int_S \left\{ \frac{|\nabla v|^p}{p} + \lambda \right\} dx < \int_{S_0} \left\{ \frac{|\nabla u|^p}{p} + \lambda \right\} dx. \quad (5.5)$$

To see this, define $\underline{u}(x) = \min\{u(x), v(x)\}$ and observe that

$$J_3(\underline{u}) = \int_{(\mathcal{B} - \sigma n) \setminus S} \left\{ \frac{|\nabla v|^p}{p} + \lambda \right\} dx + \int_{S_0} \left\{ \frac{|\nabla u|^p}{p} + \lambda \right\} dx. \quad (5.6)$$

If inequality (5.5) were violated, we would have $J_3(\underline{u}) \leq J_3(v)$. Since $\underline{u} \equiv s$ on the ball $B_R - \sigma n$, and since $J_3(v)$ is the minimum of J_3 under that constraint, we would arrive at $J_3(\underline{u}) = J_3(v)$ contradicting the fact that the minimizer under that constraint is unique

and radial. Hence inequality (5.5) must hold, as claimed. To complete the proof, define $\bar{u}(x) = \max\{u(x), v(x)\}$. We have

$$J_3(\bar{u}) = \int_{\Omega_0 \setminus S} \left\{ \frac{|\nabla u|^p}{p} + \lambda \right\} dx + \int_S \left\{ \frac{|\nabla v|^p}{p} + \lambda \right\} dx \quad (5.7)$$

and also

$$J_3(u) = \int_{\Omega_0 \setminus S} \left\{ \frac{|\nabla u|^p}{p} + \lambda \right\} dx + \int_{S_0} \left\{ \frac{|\nabla u|^p}{p} + \lambda \right\} dx. \quad (5.8)$$

By (5.5), we get $J_3(\bar{u}) < J_3(u)$, but this is impossible because $\bar{u} \equiv 1$ on Ω_1 , and u minimizes J_3 under that constraint by assumption. The lemma follows.

Now we can give a **proof of Theorem 10**. Lemma 13 prevents the maximum in $\bar{\Omega}$ of the difference $w = u^* - u$ from being attained on $\partial\Omega_0$ unless $w \equiv 0$. Furthermore, if an interior, positive maximum M is attained at some $x_0 \in \Omega$, then Lemma 12 implies that

$$|\nabla u(x_0)| > a, \quad (5.9)$$

because the variation of u must coincide with the one of u^* . Arguing as in the proof of Theorem 7, and using (5.9) in place of Proposition 5, we still find a convenient neighborhood G of x_0 such that $\nabla u \neq 0$ in G and $w < M$ on ∂G . Since this contradicts Proposition 4, we must have $u^* = u$, as claimed.

Acknowledgement: This research was financially supported by Università di Cagliari. Its subject is part of the ESF Programme “Global and geometrical aspects of nonlinear partial differential equations (GLOBAL)”.

References

- [1] Acker, A. & Meyer, R., *A free boundary problem for the p -Laplacian: uniqueness, convexity and successive approximation of solutions*. Electron. J. Differential Equations **1995**(8) (1995) pp. 1–20.
- [2] Alvarez, O., Lasry, J.-M. & Lions, P.-L., *Convex viscosity solutions and state constraints*. J. Math. Pures Appl. **76** (1997) pp. 265–288.
- [3] Barles, G. & Busca, J., *Existence and comparison results for fully nonlinear degenerate elliptic equations without zeroth-order term*. Comm. Partial Differential Equations **26** (2001) pp. 2323–2337.
- [4] Beurling, A., *On free boundary problems for the Laplace equation*. Sem. on Analytic Functions, Inst. f. Adv. Studies, Princeton **1** (1957) pp. 248–263.
- [5] Bianchini, C., Longinetti, M. & Salani, P., *Quasiconcave solutions to elliptic problems in convex rings*. Indiana Univ. Math. J., to appear
- [6] Cardaliaguet, P. & Tahraoui, R., *On the strict concavity of the harmonic radius in dimension $N \geq 3$* . J. Math. Pures Appl. **81** (2002) pp. 223–240.
- [7] Colesanti, A. & Salani, P., *Quasi-concave envelope of a function and convexity of level sets of solutions to elliptic equations*. Math. Nachr. **258** (2003) pp. 3–15.

- [8] Crandall M.G., Ishii H. & Lions P.-L., *User's guide to viscosity solutions of second order partial differential equations*. Bull. Amer. Math. Soc. **27** (1992) pp. 1–67.
- [9] Cuoghi, P. & Salani, P., *Convexity of level sets for solutions to nonlinear elliptic problems in convex rings*. Electron. J. Differential Equations **2006**(124) (2006) pp. 1–12.
- [10] Cryer, C.W., *A proof of the convexity of the free boundary for porous flow through a rectangular dam using the maximum principle*. J. Inst. Math. Appl. **25** (1980) pp. 111–120.
- [11] DiBenedetto, E., *$C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations*. Nonlinear Anal. **7** (1983) pp. 827–850.
- [12] Friedman, A. & Jensen, R., *Convexity of the free boundary in the Stefan Problem and in the dam problem*. Arch. Rational Mech. Anal. **67** (1977) pp. 1–24.
- [13] Gabriel, R.M., *A result concerning convex level surfaces of 3-dimensional harmonic functions*. J. London Math. Soc. **32** (1957) pp. 286–294.
- [14] Gilbarg, D. & Trudinger, N.S., *Elliptic Differential Equations of Second Order*, second ed., Springer Grundlehren **224**, Springer-Verlag, Heidelberg, 1998.
- [15] Gustafsson, B. & Shahgholian, H., *Existence and geometric properties of solutions of a free boundary problem in potential theory*. J. reine angew. Math. **473** (1996) pp. 137–179.
- [16] Greco, A., *Quasi-concavity for semilinear elliptic equations with non-monotone and anisotropic nonlinearities*. Bound. Value Probl. **2006** (2006) Article ID 80347, pp. 1–15.
- [17] Henrot, A. & Shahgholian, H., *Existence of classical solutions to a free boundary problem for the p -Laplace operator: (I) the exterior convex case*. J. reine angew. Math. **521** (2000) pp. 85–97.
- [18] Imbert, C., *Convexity of solutions and $C^{1,1}$ estimates for fully nonlinear elliptic equations*. J. Math. Pures Appl. **85** (2006) pp. 791–807.
- [19] Kawohl, B., *Rearrangements and convexity of level sets in PDE*. Springer Lecture Notes in Mathematics **1150**. Springer-Verlag, Heidelberg, 1985.
- [20] Kawohl, B., *On starshaped rearrangement and applications*. Trans. Amer. Math. Soc. **296** (1986) pp. 377–386.
- [21] Kawohl, B., *On the convexity and symmetry of solutions to a free boundary problem*. J. Austr. Math. Soc. Ser. A **42** (1987) pp. 57–68.
- [22] Kawohl, B., *On the convexity of level sets for elliptic and parabolic exterior boundary value problems*. In: *Potential Theory*, Ed.: J.Kral, J.Lukes, I.Netuka, J.Vesely. Plenum Publishers, New York (1988) pp. 153–159.
- [23] Lay, S.R., *Convex sets and their applications*. John Wiley & Sons, New York, 1982.
- [24] Lewis, J.L., *Capacitary functions in convex rings*. Arch. Rational Mech. Anal. **66** (1977) pp. 201–224.
- [25] Longinetti M. & Salani P., *On the Hessian matrix and Minkowski addition of quasi-convex functions*. J. Math. Pures Appl. **88** (2007) pp. 276–292.

- [26] Lu, G. & Wang, P., *A uniqueness theorem for degenerate elliptic equations*. Manuscript available in pdf at http://www.math.wayne.edu/~pywang/Degen_EQ_Italy2008.pdf
- [27] Oberman, A.M., *The convex envelope is the solution of a nonlinear obstacle problem*. Proc. Amer. Math. Soc. **135** (2007) pp. 1689–1694.
- [28] Rockafellar, R.T., *Convex Analysis*. Princeton University Press, Princeton, N.J., 1970.
- [29] Salani, P., *Starshapedness of level sets of solutions to elliptic PDEs*. Appl. Anal. **84** (2005) pp. 1185–1197.
- [30] Sakaguchi, S., *Concavity properties of solutions to some degenerate quasilinear elliptic Dirichlet problems*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. Ser. 4, **14** (1987) pp. 403–421.
- [31] Tolksdorf, P., *Regularity for a more general class of quasilinear elliptic equations*. J. Differential Equations **51** (1984) pp. 126–150.