



Universität zu Köln
Mathematisches Institut
Prof. Dr. F. Vallentin
Dr. A. Gundert
Dr. F. von Heymann

Convex Optimization

Winter Term 2018/19

— Exercise Sheet 12 —

Exercise 12.1. Determine the Löwner-John ellipsoid $\mathcal{E}_{\text{in}}(C_n)$ of the regular n -gon C_n in the plane

$$C_n = \text{conv}\{(\cos(2\pi k/n), \sin(2\pi k/n)) \in \mathbb{R}^2 : k = 0, 1, \dots, n-1\}.$$

Exercise 12.2.

(a) Let $P \subseteq \mathbb{R}^n$ be an n -dimensional polytope and let $A \in \mathbb{R}^{n \times n}$ be an invertible matrix. Show:

$$\mathcal{E}_{\text{out}}(AP) = A\mathcal{E}_{\text{out}}(P).$$

(b) Let $T \subseteq \mathbb{R}^3$ be a regular tetrahedron with inradius 1. Show that $\mathcal{E}_{\text{in}}(T) = B_3$ and that

$$B_3 \subseteq T \subseteq 3B_3.$$

holds.

Exercise 12.3. (Hand-in) Let $C \in \mathcal{S}^n$ be a symmetric matrix and let $G = (V, E)$ be a graph with vertex set $V = [n]$. The solution of the following MAXDET problem

$$\begin{aligned} \max \quad & \det \left(C + \sum_{\{i,j\} \in E} x_{ij} E_{ij} \right)^{1/n} \\ & C + \sum_{\{i,j\} \in E} x_{ij} E_{ij} \in \mathcal{S}_{\geq 0}^n, \end{aligned}$$

is said to be a G -modification of C with maximal entropy. Show: If a G -modification of C with maximal entropy $A^* = C + \sum_{\{i,j\} \in E} x_{ij}^* E_{ij}$ exists, then

$$\forall \{i, j\} \in E : ((A^*)^{-1})_{ij} = 0.$$

Exercise 12.4. (Hand-in) Let $P \subseteq \mathbb{R}^n$ be a centrally symmetric polytope ($P = -P$). Find a conic program (with possibly infinitely many constraints) which determines the minimal value $\rho \in \mathbb{R}$ such that there exists an ellipsoid E for which $E \subseteq P \subseteq \rho E$ holds.

Hand-in: Until Wednesday January 23, 12:00 (noon).

Exercises 12.3 and 12.4 to be submitted to the “Convex optimization” mailbox in room 3.01 (Studierendenarbeitsraum) of the Mathematical Institute.