


Introduction to Stochastic Analysis



Andreas Eberle

February 23, 2016

Contents

Contents	2
I Stochastic Processes	13
1 Brownian Motion	14
1.1 From Random Walks to Brownian Motion	15
Central Limit Theorem	16
Brownian motion as a Lévy process.	20
Brownian motion as a Markov process.	21
Wiener Measure	24
1.2 Brownian Motion as a Gaussian Process	27
Multivariate normals	27
Gaussian processes	31
1.3 The Wiener-Lévy Construction	38
A first attempt	39
The Wiener-Lévy representation of Brownian motion	41
Lévy's construction of Brownian motion	47
1.4 The Brownian Sample Paths	52
Typical Brownian sample paths are nowhere differentiable	52
Hölder continuity	54
Law of the iterated logarithm	56
Passage times	57
1.5 Strong Markov property and reflection principle	60

Maximum of Brownian motion	60
Strong Markov property for Brownian motion	63
A rigorous reflection principle	65
II Introduction to Stochastic Analysis	67
2 Martingales in discrete time	68
2.1 Definitions and examples	68
Martingales and supermartingales	69
Some fundamental examples	71
2.2 Doob Decomposition and Martingale Problem	78
Doob Decomposition	78
Conditional Variance Process	79
Martingale problem	83
2.3 Gambling strategies and stopping times	85
Martingale transforms	85
Stopped Martingales	90
Optional Stopping Theorems	95
Wald's identity for random sums	99
2.4 Maximal inequalities	99
Doob's inequality	100
L^p inequalities	102
Hoeffding's inequality	103
3 Martingales in continuous time	107
3.1 Some fundamental martingales of Brownian Motion	108
Filtrations generated by Brownian motion	108
Brownian Martingales	109
3.2 Optional Sampling and Optional Stopping	112
The Optional Sampling Theorem	112
Ruin probabilities and passage times revisited	115
Exit laws and Dirichlet problem	116

3.3	Maximal inequalities and the LIL	118
	Maximal inequalities in continuous time	118
	Application to LIL	120
4	Martingale Convergence Theorems	125
4.1	Convergence in L^2	125
	Martingales in L^2	125
	The Convergence Theorem	126
	Summability of sequences with random signs	128
	L^2 convergence in continuous time	129
4.2	Almost sure convergence of supermartingales	130
	Doob's upcrossing inequality	131
	Proof of Doob's Convergence Theorem	133
	Examples and first applications	134
	Generalized Borel-Cantelli Lemma	138
	Upcrossing inequality and convergence theorem in continuous time	140
4.3	Uniform integrability and L^1 convergence	141
	Uniform integrability	142
	Definitive version of Lebesgue's Dominated Convergence Theorem	145
	L^1 convergence of martingales	147
	Backward Martingale Convergence	150
5	Stochastic Integration w.r.t. Continuous Martingales	152
5.1	Defining stochastic integrals: A first attempt	154
	Riemann sum approximations	154
	Itô integrals for continuous bounded integrands	156
	The Hilbert space M_c^2	158
	Definition of Itô integral in M_c^2	161
5.2	Itô's isometry	162
	Predictable step functions	162
	Itô's isometry for Brownian motion	165
	Itô's isometry for martingales	167

	Definition of Itô integrals for square-integrable integrands	169
	Identification of admissible integrands	171
5.3	Localization	174
	Local dependence on integrand and integrator	175
	Itô integrals for locally square-integrable integrands	176
	Stochastic integrals as local martingales	178
	Approximation by Riemann-Itô sums	181
6	Itô's formula and pathwise integrals	183
6.1	Stieltjes integrals and chain rule	185
	Lebesgue-Stieltjes integrals	185
	The chain rule in Stieltjes calculus	188
6.2	Quadratic variation and Itô's formula	191
	Quadratic variation	191
	Itô's formula and pathwise integrals in \mathbb{R}^1	194
	The chain rule for anticipative integrals	197
6.3	Itô's formula for Brownian motion and martingales	198
	Quadratic variation of Brownian motion	199
	Itô's formula for Brownian motion	201
	Quadratic variation of continuous martingales	204
	From continuous martingales to Brownian motion	207
6.4	Multivariate and time-dependent Itô formula	209
	Covariation	210
	Itô to Stratonovich conversion	211
	Itô's formula in \mathbb{R}^d	212
	Product rule, integration by parts	214
	Time-dependent Itô formula	216
7	Brownian Motion and PDE	221
7.1	Dirichlet problem, recurrence and transience	222
	The Dirichlet problem revisited	222
	Recurrence and transience of Brownian motion in \mathbb{R}^d	223

7.2	Boundary value problems, exit and occupation times	227
	The stationary Feynman-Kac-Poisson formula	227
	Poisson problem and mean exit time	230
	Occupation time density and Green function	231
	Stationary Feynman-Kac formula and exit time distributions	232
	Boundary value problems in \mathbb{R}^d and total occupation time	234
7.3	Heat equation and time-dependent FK formula	236
	Brownian Motion with Absorption	237
	Time-dependent Feynman-Kac formula	240
	Occupation times and arc-sine law	242
8	SDE: Explicit Computations	245
8.1	Stochastic Calculus for Itô processes	248
	Stochastic integrals w.r.t. Itô processes	249
	Calculus for Itô processes	252
	The Itô-Doeblin formula in \mathbb{R}^1	255
	Martingale problem for solutions of SDE	257
8.2	Stochastic growth	257
	Scale functions and exit distributions	258
	Recurrence and asymptotics	259
	Geometric Brownian motion	261
	Feller's branching diffusion	262
	Cox-Ingersoll-Ross model	264
8.3	Linear SDE with additive noise	265
	Variation of constants	266
	Solutions as Gaussian processes	267
	The Ornstein-Uhlenbeck process	269
	Change of time-scale	272
8.4	Brownian bridge	274
	Wiener-Lévy construction	275
	Finite-dimensional distributions	276
	SDE for the Brownian bridge	278

8.5	Stochastic differential equations in \mathbb{R}^n	280
	Existence, uniqueness and stability	280
	Itô processes driven by several Brownian motions	282
	Multivariate Itô-Doebelin formula	283
	General Ornstein-Uhlenbeck processes	285
	Examples	285
9	Change of measure	286
9.1	Local and global densities of probability measures	286
	Absolute Continuity	286
	From local to global densities	289
	Derivatives of monotone functions	293
	Absolute continuity of infinite product measures	294
9.2	Translations of Wiener measure	298
	The Cameron-Martin Theorem	299
	Passage times for Brownian motion with constant drift	304
9.3	Girsanov transform	305
	Change of measure on filtered probability spaces	306
	Girsanov's Theorem	307
	Novikov's condition	310
9.4	Itô's Representation Theorem and Option Pricing	312
	Representation theorems for functions and martingales	313
	Application to option pricing	316
	Application to stochastic filtering	317
III	Stochastic Analysis	318
10	Lévy processes and Poisson point processes	313
10.1	Lévy processes	314
	Characteristic exponents	315
	Basic examples	316
	Compound Poisson processes	318

Examples with infinite jump intensity	321
10.2 Martingales and Markov property	325
Martingales of Lévy processes	325
Lévy processes as Markov processes	326
10.3 Poisson random measures and Poisson point processes	330
The jump times of a Poisson process	330
The jumps of a Lévy process	333
Poisson point processes	336
Construction of compound Poisson processes from PPP	338
10.4 Stochastic integrals w.r.t. Poisson point processes	340
Elementary integrands	341
Lebesgue integrals	344
Itô integrals w.r.t. compensated Poisson point processes	345
10.5 Lévy processes with infinite jump intensity	347
Construction from Poisson point processes	347
The Lévy-Itô decomposition	351
Subordinators	354
Stable processes	357
11 Transformations of SDE	360
11.1 Lévy characterizations and martingale problems	362
Lévy's characterization of Brownian motion	364
Martingale problem for Itô diffusions	367
Lévy characterization of weak solutions	371
11.2 Random time change	373
Continuous local martingales as time-changed Brownian motions	373
Time-change representations of stochastic integrals	376
Time substitution in stochastic differential equations	376
One-dimensional SDE	379
11.3 Change of measure	382
Change of measure for Brownian motion	384
Applications to SDE	386

Doob's h -transform	388
11.4 Path integrals and bridges	389
Path integral representation	390
The Markov property	392
Bridges and heat kernels	393
SDE for diffusion bridges	395
11.5 Large deviations on path spaces	397
Support of Wiener measure	397
Schilder's Theorem	399
Random perturbations of dynamical systems	403
12 Extensions of Itô calculus	406
12.1 SDE with jumps	407
L^p Stability	409
Existence of strong solutions	411
Non-explosion criteria	414
12.2 Stratonovich differential equations	416
Itô-Stratonovich formula	416
Stratonovich SDE	418
Brownian motion on hypersurfaces	419
Doss-Sussmann method	422
Wong Zakai approximations of SDE	425
12.3 Stochastic Taylor expansions	426
Itô-Taylor expansions	426
12.4 Numerical schemes for SDE	431
Strong convergence order	432
Weak convergence order	437
12.5 Local time	439
Local time of continuous semimartingales	440
Itô-Tanaka formula	444
12.6 Continuous modifications and stochastic flows	446
Continuous modifications of deterministic functions	447

Continuous modifications of random fields	449
Existence of a continuous flow	451
Markov property	453
Continuity of local time	454
13 Variations of parameters in SDE	457
13.1 Variations of parameters in SDE	458
Differentiation of solutions w.r.t. a parameter	459
Derivative flow and stability of SDE	460
Consequences for the transition semigroup	464
13.2 Malliavin gradient and Bismut integration by parts formula	467
Gradient and integration by parts for smooth functions	468
Skorokhod integral	472
Definition of Malliavin gradient II	473
Product and chain rule	475
Clark-Ocone formula	476
13.3 First applications to SDE	477
13.4 Existence and smoothness of densities	477
14 Stochastic calculus for semimartingales with jumps	478
Semimartingales in discrete time	479
Semimartingales in continuous time	480
14.1 Finite variation calculus	483
Lebesgue-Stieltjes integrals revisited	484
Product rule	485
Chain rule	488
Exponentials of finite variation functions	490
14.2 Stochastic integration for semimartingales	498
Integrals with respect to bounded martingales	498
Localization	504
Integration w.r.t. semimartingales	507
14.3 Quadratic variation and covariation	508

Covariation and integration by parts	509
Quadratic variation and covariation of local martingales	511
Covariation of stochastic integrals	517
The Itô isometry for stochastic integrals w.r.t. martingales	520
14.4 Itô calculus for semimartingales	521
Integration w.r.t. stochastic integrals	521
Itô's formula	522
Application to Lévy processes	525
Burkholder's inequality	528
14.5 Stochastic exponentials and change of measure	530
Exponentials of semimartingales	530
Change of measure for Poisson point processes	533
Change of measure for Lévy processes	536
Change of measure for general semimartingales	539
14.6 General predictable integrands	539
Definition of stochastic integrals w.r.t. semimartingales	541
Localization	543
Properties of the stochastic integral	544
A Conditional expectations	549
A.1 Conditioning on discrete random variables	549
Conditional expectations as random variables	549
Characteristic properties of conditional expectations	550
A.2 General conditional expectations	551
The factorization lemma	551
Conditional expectations given σ -algebras	553
Properties of conditional expectations	555
A.3 Conditional expectation as best L^2 -approximation	557
Jensen's inequality	558
Conditional expectation as best L^2 -prediction value	559
Existence of conditional expectations	561

Index	564
Bibliography	567

Part I

Stochastic Processes

Chapter 1

Brownian Motion

This introduction to stochastic analysis starts with an introduction to Brownian motion. Brownian Motion is a diffusion process, i.e. a continuous-time Markov process $(B_t)_{t \geq 0}$ with continuous sample paths $t \mapsto B_t(\omega)$. In fact, it is the only nontrivial continuous-time process that is a Lévy process as well as a martingale and a Gaussian process. A rigorous construction of this process has been carried out first by N. Wiener in 1923. Already about 20 years earlier, related models had been introduced independently for financial markets by L. Bachelier [*Théorie de la spéculation*, Ann. Sci. École Norm. Sup. 17, 1900], and for the velocity of molecular motion by A. Einstein [*Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen*, *Annalen der Physik* 17, 1905].

It has been a groundbreaking approach of K. Itô to construct general diffusion processes from Brownian motion, cf. [...]. In classical analysis, the solution of an ordinary differential equation $x'(t) = f(t, x(t))$ is a function, that can be approximated locally for t close to t_0 by the linear function $x(t_0) + f(t_0, x(t_0)) \cdot (t - t_0)$. Similarly, Itô showed, that a diffusion process behaves locally like a linear function of Brownian motion – the connection being described rigorously by a stochastic differential equation (SDE).

The fundamental rôle played by Brownian motion in stochastic analysis is due to the central limit Theorem. Similarly as the normal distribution arises as a universal scaling limit of standardized sums of independent, identically distributed, square integrable

random variables, Brownian motion shows up as a universal scaling limit of Random Walks with square integrable increments.

1.1 From Random Walks to Brownian Motion

To motivate the definition of Brownian motion below, we first briefly discuss discrete-time stochastic processes and possible continuous-time scaling limits on an informal level.

A standard approach to model stochastic dynamics in discrete time is to start from a sequence of random variables η_1, η_2, \dots defined on a common probability space (Ω, \mathcal{A}, P) . The random variables η_n describe the stochastic influences (*noise*) on the system. Often they are assumed to be *independent and identically distributed (i.i.d.)*. In this case the collection (η_n) is also called a **white noise**, whereas a **colored noise** is given by dependent random variables. A stochastic process $X_n, n = 0, 1, 2, \dots$, taking values in \mathbb{R}^d is then defined recursively on (Ω, \mathcal{A}, P) by

$$X_{n+1} = X_n + \Phi_{n+1}(X_n, \eta_{n+1}), \quad n = 0, 1, 2, \dots \quad (1.1.1)$$

Here the Φ_n are measurable maps describing the *random law of motion*. If X_0 and η_1, η_2, \dots are independent random variables, then the process (X_n) is a Markov chain with respect to P .

Now let us assume that the random variables η_n are independent and identically distributed taking values in \mathbb{R} , or, more generally, \mathbb{R}^d . The easiest type of a nontrivial stochastic dynamics as described above is the Random Walk $S_n = \sum_{i=1}^n \eta_i$ which satisfies

$$S_{n+1} = S_n + \eta_{n+1} \quad \text{for } n = 0, 1, 2, \dots$$

Since the noise random variables η_n are the increments of the Random Walk (S_n) , the law of motion (1.1.1) in the general case can be rewritten as

$$X_{n+1} - X_n = \Phi_{n+1}(X_n, S_{n+1} - S_n), \quad n = 0, 1, 2, \dots \quad (1.1.2)$$

This equation is a difference equation for (X_n) driven by the stochastic process (S_n) .

Our aim is to carry out a similar construction as above for stochastic dynamics in continuous time. The stochastic difference equation (1.1.2) will then eventually be replaced by a *stochastic differential equation (SDE)*. However, before even being able to think about how to write down and make sense of such an equation, we have to identify a continuous-time stochastic process that takes over the rôle of the Random Walk. For this purpose, we first determine possible scaling limits of Random Walks when the time steps tend to 0. It will turn out that if the increments are square integrable and the size of the increments goes to 0 as the length of the time steps tends to 0, then by the Central Limit Theorem there is essentially only one possible limit process in continuous time: Brownian motion.

Central Limit Theorem

Suppose that $Y_{n,i} : \Omega \rightarrow \mathbb{R}^d, 1 \leq i \leq n < \infty$, are identically distributed, square-integrable random variables on a probability space (Ω, \mathcal{A}, P) such that $Y_{n,1}, \dots, Y_{n,n}$ are independent for each $n \in \mathbb{N}$. Then the rescaled sums

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n (Y_{n,i} - E[Y_{n,i}])$$

converge in distribution to a multivariate normal distribution $N(0, C)$ with covariance matrix

$$C_{kl} = \text{Cov}[Y_{n,i}^{(k)}, Y_{n,i}^{(l)}].$$


To see, how the CLT determines the possible scaling limits of Random Walks, let us consider a one-dimensional Random Walk

$$S_n = \sum_{i=1}^n \eta_i, \quad n = 0, 1, 2, \dots,$$

on a probability space (Ω, \mathcal{A}, P) with independent increments $\eta_i \in \mathcal{L}^2(\Omega, \mathcal{A}, P)$ normalized such that

$$E[\eta_i] = 0 \quad \text{and} \quad \text{Var}[\eta_i] = 1. \quad (1.1.3)$$

Plotting many steps of the Random Walk seems to indicate that there is a limit process with continuous sample paths after appropriate rescaling:



To see what appropriate means, we fix a positive integer m , and try to define a rescaled Random Walk $S_t^{(m)}$ ($t = 0, 1/m, 2/m, \dots$) with time steps of size $1/m$ by

$$S_{k/m}^{(m)} = c_m \cdot S_k \quad (k = 0, 1, 2, \dots)$$

for some constants $c_m > 0$. If t is a multiple of $1/m$, then

$$\text{Var}[S_t^{(m)}] = c_m^2 \cdot \text{Var}[S_{mt}] = c_m^2 \cdot m \cdot t.$$

Hence in order to achieve convergence of $S_t^{(m)}$ as $m \rightarrow \infty$, we should choose c_m proportional to $m^{-1/2}$. This leads us to define a continuous time process $(S_t^{(m)})_{t \geq 0}$ by

$$S_t^{(m)}(\omega) := \frac{1}{\sqrt{m}} S_{mt}(\omega) \quad \text{whenever } t = k/m \text{ for some integer } k,$$

and by linear interpolation for $t \in (\frac{k-1}{m}, \frac{k}{m}]$.




Figure 1.1: Rescaling of a Random Walk.

Clearly,

$$E[S_t^{(m)}] = 0 \quad \text{for all } t \geq 0,$$

and

$$\text{Var}[S_t^{(m)}] = \frac{1}{m} \text{Var}[S_{mt}] = t$$

whenever t is a multiple of $1/m$. In particular, the expectation values and variances for a fixed time t do not depend on m . Moreover, if we fix a partition $0 \leq t_0 < t_1 < \dots < t_n$ such that each t_i is a multiple of $1/m$, then the increments

$$S_{t_{i+1}}^{(m)} - S_{t_i}^{(m)} = \frac{1}{\sqrt{m}} (S_{mt_{i+1}} - S_{mt_i}), \quad i = 0, 1, 2, \dots, n-1, \quad (1.1.4)$$

of the rescaled process $(S_t^{(m)})_{t \geq 0}$ are independent centered random variables with variances $t_{i+1} - t_i$. If t_i is not a multiple of $1/m$, then a corresponding statement holds approximately with an error that should be negligible in the limit $m \rightarrow \infty$. Hence, if the rescaled Random Walks $(S_t^{(m)})_{t \geq 0}$ converge in distribution to a limit process $(B_t)_{t \geq 0}$, then $(B_t)_{t \geq 0}$ should have *independent increments* $B_{t_{i+1}} - B_{t_i}$ over *disjoint time intervals with mean 0 and variances* $t_{i+1} - t_i$.

It remains to determine the precise distributions of the increments. Here the Central Limit Theorem applies. In fact, we can observe that by (1.1.4) each increment

$$S_{t_{i+1}}^{(m)} - S_{t_i}^{(m)} = \frac{1}{\sqrt{m}} \sum_{k=mt_i+1}^{mt_{i+1}} \eta_k$$

of the rescaled process is a rescaled sum of $m \cdot (t_{i+1} - t_i)$ i.i.d. random variables with mean 0 and variance 1. Therefore, the CLT implies that the distributions of the increments converge weakly to a normal distribution:

$$S_{t_{i+1}}^{(m)} - S_{t_i}^{(m)} \xrightarrow{\mathcal{D}} N(0, t_{i+1} - t_i).$$

Hence if a limit process (B_t) exists, then it should have *independent, normally distributed increments*.

Our considerations motivate the following definition:

Definition (Brownian Motion).

(1). Let $a \in \mathbb{R}$. A continuous-time stochastic process $B_t : \Omega \rightarrow \mathbb{R}$, $t \geq 0$, defined on a probability space (Ω, \mathcal{A}, P) , is called a **Brownian motion (starting in a)** if and only if

(a) $B_0(\omega) = a$ for each $\omega \in \Omega$.

(b) For any partition $0 \leq t_0 < t_1 < \dots < t_n$, the increments $B_{t_{i+1}} - B_{t_i}$ are independent random variables with distribution

$$B_{t_{i+1}} - B_{t_i} \sim N(0, t_{i+1} - t_i).$$

(c) P -almost every sample path $t \mapsto B_t(\omega)$ is continuous.

(2). An \mathbb{R}^d -valued stochastic process $B_t(\omega) = (B_t^{(1)}(\omega), \dots, B_t^{(d)}(\omega))$ is called a *multi-dimensional Brownian motion* if and only if the component processes $(B_t^{(1)}), \dots, (B_t^{(d)})$ are independent one-dimensional Brownian motions.

Thus the increments of a d -dimensional Brownian motion are independent over disjoint time intervals and have a multivariate normal distribution:

$$B_t - B_s \sim N(0, (t - s) \cdot I_d) \quad \text{for any } 0 \leq s \leq t.$$

Remark. (1). *Continuity:* Continuity of the sample paths has to be assumed separately: If $(B_t)_{t \geq 0}$ is a one-dimensional Brownian motion, then the modified process $(\tilde{B}_t)_{t \geq 0}$ defined by $\tilde{B}_0 = B_0$ and

$$\tilde{B}_t = B_t \cdot I_{\{B_t \in \mathbb{R} \setminus \mathbb{Q}\}} \quad \text{for } t > 0$$

has almost surely discontinuous paths. On the other hand, it satisfies (a) and (b) since the distributions of $(\tilde{B}_{t_1}, \dots, \tilde{B}_{t_n})$ and $(B_{t_1}, \dots, B_{t_n})$ coincide for all $n \in \mathbb{N}$ and $t_1, \dots, t_n \geq 0$.

- (2). *Spatial Homogeneity:* If $(B_t)_{t \geq 0}$ is a Brownian motion starting at 0, then the translated process $(a + B_t)_{t \geq 0}$ is a Brownian motion starting at a .
- (3). *Existence:* There are several constructions and existence proofs for Brownian motion. In Section 1.3 below we will discuss in detail the Wiener-Lévy construction of Brownian motion as a random superposition of infinitely many deterministic paths. This explicit construction is also very useful for numerical approximations. A more general (but less constructive) existence proof is based on Kolmogorov's extension Theorem, cf. e.g. [Klenke].
- (4). *Functional Central Limit Theorem:* The construction of Brownian motion as a scaling limit of Random Walks sketched above can also be made rigorous. *Donsker's invariance principle* is a functional version of the central limit Theorem which states that the rescaled Random Walks $(S_t^{(m)})$ converge in distribution to a Brownian motion. As in the classical CLT the limit is universal, i.e., it does not depend on the distribution of the increments η_i provided (1.1.3) holds, cf. Section ??.

Brownian motion as a Lévy process.

The definition of Brownian motion shows in particular that Brownian motion is a *Lévy process*, i.e., it has stationary independent increments (over disjoint time intervals). In fact, the analogues of Lévy processes in discrete time are Random Walks, and it is rather obvious, that all scaling limits of Random Walks should be Lévy processes. Brownian

motion is the only Lévy process L_t in continuous time with paths such that $E[L_1] = 0$ and $\text{Var}[L_1] = 1$. The normal distribution of the increments follows under these assumptions by an extension of the CLT, cf. e.g. [Breiman: Probability]. A simple example of a Lévy process with non-continuous paths is the Poisson process. Other examples are α -stable processes which arise as scaling limits of Random Walks when the increments are not square-integrable. Stochastic analysis based on general Lévy processes has attracted a lot of interest recently.

Let us now consider a Brownian motion $(B_t)_{t \geq 0}$ starting at a fixed point $a \in \mathbb{R}^d$, defined on a probability space (Ω, \mathcal{A}, P) . The information on the process up to time t is encoded in the σ -algebra

$$\mathcal{F}_t^B = \sigma(B_s \mid 0 \leq s \leq t)$$

generated by the process. The independence of the increments over disjoint intervals immediately implies:

Lemma 1.1. *For any $0 \leq s \leq t$, the increment $B_t - B_s$ is independent of \mathcal{F}_s^B .*

Proof. For any partition $0 = t_0 \leq t_1 \leq \dots \leq t_n = s$ of the interval $[0, s]$, the increment $B_t - B_s$ is independent of the σ -algebra

$$\sigma(B_{t_1} - B_{t_0}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}})$$

generated by the increments up to time s . Since

$$B_{t_k} = B_{t_0} + \sum_{i=1}^k (B_{t_i} - B_{t_{i-1}})$$

and B_{t_0} is constant, this σ -algebra coincides with $\sigma(B_{t_0}, B_{t_1}, \dots, B_{t_n})$. Hence $B_t - B_s$ is independent of all finite subcollections of $(B_u \mid 0 \leq u \leq s)$ and therefore independent of \mathcal{F}_s^B . \square

Brownian motion as a Markov process.

As a process with stationary increments, Brownian motion is in particular a time-homogeneous Markov process. In fact, we have:

Theorem 1.2 (Markov property). *A Brownian motion $(B_t)_{t \geq 0}$ in \mathbb{R}^d is a time-homogeneous Markov process with transition densities*

$$p_t(x, y) = (2\pi t)^{-d/2} \cdot \exp\left(-\frac{|x - y|^2}{2t}\right), \quad t > 0, \quad x, y \in \mathbb{R}^d,$$

i.e., for any Borel set $A \subseteq \mathbb{R}^d$ and $0 \leq s < t$,

$$P[B_t \in a \mid \mathcal{F}_s^B] = \int_A p_{t-s}(B_s, y) dy \quad P\text{-almost surely.}$$

Proof. For $0 \leq s < t$ we have $B_t = B_s + (B_t - B_s)$ where B_s is \mathcal{F}_s^B -measurable, and $B_t - B_s$ is independent of \mathcal{F}_s^B by Lemma 1.1. Hence

$$\begin{aligned} P[B_t \in A \mid \mathcal{F}_s^B](\omega) &= P[B_s(\omega) + B_t - B_s \in A] = N(B_s(\omega), (t - s) \cdot I_d)[A] \\ &= \int_A (2\pi(t - s))^{-d/2} \cdot \exp\left(-\frac{|y - B_s(\omega)|^2}{2(t - s)}\right) dy \quad P\text{-almost surely.} \end{aligned}$$

□

Remark (Heat equation as backward equation and forward equation). The transition function of Brownian motion is the *heat kernel* in \mathbb{R}^d , i.e., it is the fundamental solution of the heat equation

$$\frac{\partial u}{\partial t} = \frac{1}{2} \Delta u.$$

More precisely, $p_t(x, y)$ solves the initial value problem

$$\begin{aligned} \frac{\partial}{\partial t} p_t(x, y) &= \frac{1}{2} \Delta_x p_t(x, y) \quad \text{for any } t > 0, x, y \in \mathbb{R}^d, \\ \lim_{t \searrow 0} \int p_t(x, y) f(y) dy &= f(x) \quad \text{for any } f \in C_b(\mathbb{R}^d), x \in \mathbb{R}^d, \end{aligned} \tag{1.1.5}$$

where $\Delta_x = \sum_{i=1}^d \frac{\partial^2}{\partial x_i^2}$ denotes the action of the Laplace operator on the x -variable. The equation (1.1.5) can be viewed as a version of *Kolmogorov's backward equation* for

Brownian motion as a time-homogeneous Markov process, which states that for each $t > 0$, $y \in \mathbb{R}^d$ and $f \in C_b(\mathbb{R}^d)$, the function

$$v(s, x) = \int p_{t-s}(x, y) f(y) dy$$

solves the terminal value problem

$$\frac{\partial v}{\partial s}(s, x) = -\frac{1}{2} \Delta_x v(s, x) \quad \text{for } s \in [0, t), \quad \lim_{s \nearrow t} v(s, x) = f(x). \quad (1.1.6)$$

Note that by the Markov property, $v(s, x) = (p_{t-s} f)(x)$ is a version of the conditional expectation $E[f(B_t) | B_s = x]$. Therefore, the backward equation describes the dependence of the expectation value on starting point and time.

By symmetry, $p_t(x, y)$ also solves the initial value problem

$$\frac{\partial}{\partial t} p_t(x, y) = \frac{1}{2} \Delta_y p_t(x, y) \quad \text{for any } t > 0, \quad \text{and } x, y \in \mathbb{R}^d, \quad (1.1.7)$$

$$\lim_{t \searrow 0} \int g(x) p_t(x, y) dx = g(y) \quad \text{for any } g \in C_b(\mathbb{R}^d), y \in \mathbb{R}^d.$$

The equation (1.1.7) is a version of *Kolmogorov's forward equation*, stating that for $g \in C_b(\mathbb{R}^d)$, the function $u(t, y) = \int g(x) p_t(x, y) dx$ solves

$$\frac{\partial u}{\partial t}(t, y) = \frac{1}{2} \Delta_y u(t, y) \quad \text{for } t > 0, \quad \lim_{t \searrow 0} u(t, y) = g(y). \quad (1.1.8)$$

The forward equation describes the forward time evolution of the transition densities $p_t(x, y)$ for a given starting point x .

The Markov property enables us to compute the marginal distributions of Brownian motion:

Corollary 1.3 (Finite dimensional marginals). *Suppose that $(B_t)_{t \geq 0}$ is a Brownian motion starting at $x_0 \in \mathbb{R}^d$ defined on a probability space (Ω, \mathcal{A}, P) . Then for any*

$n \in \mathbb{N}$ and $0 = t_0 < t_1 < t_2 < \dots < t_n$, the joint distribution of $B_{t_1}, B_{t_2}, \dots, B_{t_n}$ is absolutely continuous with density

$$\begin{aligned} f_{B_{t_1}, \dots, B_{t_n}}(x_1, \dots, x_n) &= p_{t_1}(x_0, x_1) p_{t_2-t_1}(x_1, x_2) p_{t_3-t_2}(x_2, x_3) \cdots p_{t_n-t_{n-1}}(x_{n-1}, x_n) \\ &= \prod_{i=1}^n (2\pi(t_i - t_{i-1}))^{-d/2} \cdot \exp\left(-\frac{1}{2} \sum_{i=1}^n \frac{|x_i - x_{i-1}|^2}{t_i - t_{i-1}}\right) \end{aligned} \quad (1.1.9)$$

Proof. By the Markov property and induction on n , we obtain

$$\begin{aligned} &P[B_{t_1} \in A_1, \dots, B_{t_n} \in A_n] \\ &= E[P[B_{t_n} \in A_n \mid \mathcal{F}_{t_{n-1}}^B]; B_{t_1} \in A_1, \dots, B_{t_{n-1}} \in A_{n-1}] \\ &= E[p_{t_n-t_{n-1}}(B_{t_{n-1}}, A_n); B_{t_1} \in A_1, \dots, B_{t_{n-1}} \in A_{n-1}] \\ &= \int_{A_1} \cdots \int_{A_{n-1}} p_{t_1}(x_0, x_1) p_{t_2-t_1}(x_1, x_2) \cdots \\ &\quad \cdot p_{t_{n-1}-t_{n-2}}(x_{n-2}, x_{n-1}) p_{t_n-t_{n-1}}(x_{n-1}, A_n) dx_{n-1} \cdots dx_1 \\ &= \int_{A_1} \cdots \int_{A_n} \left(\prod_{i=1}^n p_{t_i-t_{i-1}}(x_{n-1}, x_n) \right) dx_n \cdots dx_1 \end{aligned}$$

for all $n \geq 0$ and $A_1, \dots, A_n \in \mathcal{B}(\mathbb{R}^d)$. \square

Remark (Brownian motion as a Gaussian process). The corollary shows in particular that Brownian motion is a Gaussian process, i.e., all the marginal distributions in (1.1.9) are multivariate normal distributions. We will come back to this important aspect in the next section.

Wiener Measure

The distribution of Brownian motion could be considered as a probability measure on the product space $(\mathbb{R}^d)^{[0, \infty)}$ consisting of all maps $x : [0, \infty) \rightarrow \mathbb{R}^d$. A disadvantage of this approach is that the product space is far too large for our purposes: It contains extremely irregular paths $x(t)$, although at least almost every path of Brownian motion is continuous by definition. Actually, since $[0, \infty)$ is uncountable, the subset of all

continuous paths is not even measurable w.r.t. the product σ -algebra on $(\mathbb{R}^d)^{[0,\infty)}$.

Instead of the product space, we will directly consider the distribution of Brownian motion on the continuous path space $C([0, \infty), \mathbb{R}^d)$. For this purpose, we fix a Brownian motion $(B_t)_{t \geq 0}$ starting at $x_0 \in \mathbb{R}^d$ on a probability space (Ω, \mathcal{A}, P) , and we **assume** that **every** sample path $t \mapsto B_t(\omega)$ is continuous. This assumption can always be fulfilled by modifying a given Brownian motion on a set of measure zero. The full process $(B_t)_{t \geq 0}$ can then be interpreted as a single path-space valued random variable (or a "random path").




Figure 1.2: $B : \Omega \rightarrow C([0, \infty), \mathbb{R}^d)$, $B(\omega) = (B_t(\omega))_{t \geq 0}$.

We endow the space of continuous paths $x : [0, \infty) \rightarrow \mathbb{R}^d$ with the σ -algebra

$$\mathcal{B} = \sigma(X_t \mid t \geq 0)$$

generated by the coordinate maps

$$X_t : C([0, \infty), \mathbb{R}^d) \rightarrow \mathbb{R}^d, \quad X_t(x) = x_t, \quad t \geq 0.$$

Note that we also have

$$\mathcal{B} = \sigma(X_t \mid t \in \mathcal{D})$$

for any dense subset \mathcal{D} of $[0, \infty)$, because $X_t = \lim_{s \rightarrow t} X_s$ for each $t \in [0, \infty)$ by continuity. Furthermore, it can be shown that \mathcal{B} is the Borel σ -algebra on $C([0, \infty), \mathbb{R}^d)$ endowed with the topology of uniform convergence on finite intervals.

Theorem 1.4 (Distribution of Brownian motion on path space). *The map $B : \Omega \rightarrow C([0, \infty), \mathbb{R}^d)$ is measurable w.r.t. the σ -algebras \mathcal{A}/\mathcal{B} . The distribution $P \circ B^{-1}$ of B is the unique probability measure μ_{x_0} on $(C([0, \infty), \mathbb{R}^d), \mathcal{B})$ with marginals*

$$\begin{aligned} & \mu_{x_0} [\{x \in C([0, \infty), \mathbb{R}^d) : x_{t_1} \in A_1, \dots, x_{t_n} \in A_n\}] & (1.1.10) \\ & = \prod_{i=1}^n (2\pi(t_i - t_{i-1}))^{-d/2} \int_{A_1} \dots \int_{A_n} \exp\left(-\frac{1}{2} \sum_{i=1}^n \frac{|x_i - x_{i-1}|^2}{t_i - t_{i-1}}\right) dx_n \dots dx_1 \end{aligned}$$

for any $n \in \mathbb{N}$, $0 < t_1 < \dots < t_n$, and $A_1, \dots, A_n \in \mathcal{B}(\mathbb{R}^d)$.

Definition. *The probability measure μ_{x_0} on the path space $C([0, \infty), \mathbb{R}^d)$ determined by (1.1.10) is called **Wiener measure** (with start in x_0).*

Remark (Uniqueness in distribution). The Theorem asserts that the path space distribution of a Brownian motion starting at a given point x_0 is the corresponding Wiener measure. In particular, it is uniquely determined by the marginal distributions in (1.1.9).

Proof of Theorem 1.4. For $n \in \mathbb{N}$, $0 < t_1 < \dots < t_n$, and $A_1, \dots, A_n \in \mathcal{B}(\mathbb{R}^d)$, we have

$$\begin{aligned} B^{-1}(\{X_{t_1} \in A_1, \dots, X_{t_n} \in A_n\}) &= \{\omega : X_{t_1}(B(\omega)) \in A_1, \dots, X_{t_n}(B(\omega)) \in A_n\} \\ &= \{B_{t_1} \in A_1, \dots, B_{t_n} \in A_n\} \in \mathcal{A}. \end{aligned}$$

Since the cylinder sets of type $\{X_{t_1} \in A_1, \dots, X_{t_n} \in A_n\}$ generate the σ -algebra \mathcal{B} , the map B is \mathcal{A}/\mathcal{B} -measurable. Moreover, by corollary 1.3, the probabilities

$$P[B \in \{X_{t_1} \in A_1, \dots, X_{t_n} \in A_n\}] = P[B_{t_1} \in A_1, \dots, B_{t_n} \in A_n],$$

are given by the right hand side of (1.1.10). Finally, the measure μ_{x_0} is uniquely determined by (1.1.10), since the system of cylinder sets as above is stable under intersections and generates the σ -algebra \mathcal{B} . \square

Definition (Canonical model for Brownian motion.) By (1.1.10), the coordinate process

$$X_t(x) = x_t, \quad t \geq 0,$$

on $C([0, \infty), \mathbb{R}^d)$ is a Brownian motion starting at x_0 w.r.t. Wiener measure μ_{x_0} . We refer to the stochastic process $(C([0, \infty), \mathbb{R}^d), \mathcal{B}, \mu_{x_0}, (X_t)_{t \geq 0})$ as the **canonical model for Brownian motion starting at x_0** .

1.2 Brownian Motion as a Gaussian Process

We have already verified that Brownian motion is a Gaussian process, i.e., the finite dimensional marginals are multivariate normal distributions. We will now exploit this fact more thoroughly.

Multivariate normals

Let us first recall some basics on normal random vectors:

Definition. Suppose that $m \in \mathbb{R}^n$ is a vector and $C \in \mathbb{R}^{n \times n}$ is a symmetric non-negative definite matrix. A random variable $Y : \Omega \rightarrow \mathbb{R}^n$ defined on a probability space (Ω, \mathcal{A}, P) has a **multivariate normal distribution** $\mathbf{N}(m, C)$ with **mean m and covariance matrix C** if and only if its characteristic function is given by

$$E[e^{ip \cdot Y}] = e^{ip \cdot m - \frac{1}{2} p \cdot C p} \quad \text{for any } p \in \mathbb{R}^n. \quad (1.2.1)$$

If C is non-degenerate, then a multivariate normal random variable Y is absolutely continuous with density

$$f_Y(x) = (2\pi \det C)^{-1/2} \exp\left(-\frac{1}{2}(x - m) \cdot C^{-1}(x - m)\right).$$

A degenerate normal distribution with vanishing covariance matrix is a Dirac measure:

$$N(m, 0) = \delta_m.$$

Differentiating (1.2.1) w.r.t. p shows that for a random variable $Y \sim N(m, C)$, the mean vector is m and $C_{i,j}$ is the covariance of the components Y_i and Y_j . Moreover, the following important facts hold:

Theorem 1.5 (Properties of normal random vectors).

- (1). A random variable $Y : \Omega \rightarrow \mathbb{R}^n$ has a multivariate normal distribution if and only if any linear combination

$$p \cdot Y = \sum_{i=1}^n p_i Y_i, \quad p \in \mathbb{R}^n,$$

of the components Y_i has a one dimensional normal distribution.

- (2). Any affine function of a normally distributed random vector Y is again normally distributed:

$$Y \sim N(m, C) \implies AY + b \sim N(Am + b, ACA^\top)$$

for any $d \in \mathbb{N}$, $A \in \mathbb{R}^{d \times n}$ and $b \in \mathbb{R}^d$.

- (3). If $Y = (Y_1, \dots, Y_n)$ has a multivariate normal distribution, and the components Y_1, \dots, Y_n are uncorrelated random variables, then Y_1, \dots, Y_n are independent.

Proof. (1). follows easily from the definition.

- (2). For $Y \sim N(m, C)$, $A \in \mathbb{R}^{d \times n}$ and $b \in \mathbb{R}^d$ we have

$$\begin{aligned} E[e^{ip \cdot (AY+b)}] &= e^{ip \cdot b} E[e^{i(A^\top p) \cdot Y}] \\ &= e^{ip \cdot b} e^{i(A^\top p) \cdot m - \frac{1}{2}(A^\top p) \cdot CA^\top p} \\ &= e^{ip \cdot (Am+b) - \frac{1}{2}p \cdot ACA^\top} \quad \text{for any } p \in \mathbb{R}^d, \end{aligned}$$

i.e., $AY + b \sim N(Am + b, ACA^\top)$.

- (3). If Y_1, \dots, Y_n are uncorrelated, then the covariance matrix $C_{i,j} = \text{Cov}[Y_i, Y_j]$ is a diagonal matrix. Hence the characteristic function

$$E[e^{ip \cdot Y}] = e^{ip \cdot m - \frac{1}{2} p \cdot C p} = \prod_{k=1}^n e^{im_k p_k - \frac{1}{2} C_{k,k} p_k^2}$$

is a product of characteristic functions of one-dimensional normal distributions. Since a probability measure on \mathbb{R}^n is uniquely determined by its characteristic function, it follows that the adjoint distribution of Y_1, \dots, Y_n is a product measure, i.e. Y_1, \dots, Y_n are independent. □

If Y has a multivariate normal distribution $N(m, C)$ then for any $p, q \in \mathbb{R}^n$, the random variables $p \cdot Y$ and $q \cdot Y$ are normally distributed with means $p \cdot m$ and $q \cdot m$, and covariance

$$\text{Cov}[p \cdot Y, q \cdot Y] = \sum_{i,j=1}^n p_i C_{i,j} q_j = p \cdot C q.$$

In particular, let $\{e_1, \dots, e_n\} \subseteq \mathbb{R}^n$ be an orthonormal basis consisting of eigenvectors of the covariance matrix C . Then the components $e_i \cdot Y$ of Y in this basis are uncorrelated and therefore independent, jointly normally distributed random variables with variances given by the corresponding eigenvalues λ_i :

$$\text{Cov}[e_i \cdot Y, e_j \cdot Y] = \lambda_i \delta_{i,j}, \quad 1 \leq i, j \leq n. \quad (1.2.2)$$

Correspondingly, the contour lines of the density of a non-degenerate multivariate normal distribution $N(m, C)$ are ellipsoids with center at m and principal axes of length $\sqrt{\lambda_i}$ given by the eigenvalues λ_i of the covariance matrix C .




Figure 1.3: Level lines of the density of a normal random vector $Y \sim N\left(\begin{pmatrix} 1 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}\right)$.

Conversely, we can generate a random vector Y with distribution $N(m, C)$ from i.i.d. standard normal random variables Z_1, \dots, Z_n by setting

$$Y = m + \sum_{i=1}^n \sqrt{\lambda_i} Z_i e_i. \quad (1.2.3)$$

More generally, we have:

Corollary 1.6 (Generating normal random vectors). *Suppose that $C = U\Lambda U^\top$ with a matrix $U \in \mathbb{R}^{n \times d}$, $d \in \mathbb{N}$, and a diagonal matrix $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_d) \in \mathbb{R}^{d \times d}$ with nonnegative entries λ_i . If $Z = (Z_1, \dots, Z_d)$ is a random vector with i.i.d. standard normal random components Z_1, \dots, Z_d then*

$$Y = U\Lambda^{1/2}Z + m$$

has distribution $N(m, C)$.

Proof. Since $Z \sim N(0, I_d)$, the second assertion of Theorem 1.5 implies

$$Y \sim N(m, U\Lambda U^\top).$$

□

Choosing for U the matrix (e_1, \dots, e_n) consisting of the orthonormal eigenvectors e_1, \dots, e_n of C , we obtain (1.2.3) as a special case of the corollary. For computational purposes it is often more convenient to use the Cholesky decomposition

$$C = LL^\top$$

of the covariance matrix as a product of a lower triangular matrix L and the upper triangular transpose L^\top :

Algorithm 1.7 (Simulation of multivariate normal random variables).

Given: $m \in \mathbb{R}^n, C \in \mathbb{R}^{n \times n}$ symmetric and non-negative definite.

Output: Sample $y \sim N(m, C)$.

- (1). Compute the Cholesky decomposition $C = LL^\top$.
- (2). Generate independent samples $z_1, \dots, z_n \sim N(0, 1)$ (e.g. by the Box-Muller method).
- (3). Set $y := Lz + m$.

Gaussian processes

Let I be an arbitrary index set, e.g. $I = \mathbb{N}, I = [0, \infty)$ or $I = \mathbb{R}^n$.

Definition. A collection $(Y_t)_{t \in I}$ of random variables $Y_t : \Omega \rightarrow \mathbb{R}^d$ defined on a probability space (Ω, \mathcal{A}, P) is called a **Gaussian process** if and only if the joint distribution of any finite subcollection Y_{t_1}, \dots, Y_{t_n} with $n \in \mathbb{N}$ and $t_1, \dots, t_n \in I$ is a multivariate normal distribution.

The distribution of a Gaussian process $(Y_t)_{t \in I}$ on the path space \mathbb{R}^I or $C(I, \mathbb{R})$ endowed with the σ -algebra generated by the maps $x \mapsto x_t$, $t \in I$, is uniquely determined by the multinormal distributions of finite subcollections Y_{t_1}, \dots, Y_{t_n} as above, and hence by the expectation values

$$m(t) = E[Y_t], \quad t \in I,$$

and the covariances

$$c(s, t) = \text{Cov}[Y_s, Y_t], \quad s, t \in I.$$

A Gaussian process is called **centered**, if $m(t) = 0$ for any $t \in I$.

Example (AR(1) process). The autoregressive process $(Y_n)_{n=0,1,2,\dots}$ defined recursively by $Y_0 \sim N(0, v_0)$,

$$Y_n = \alpha Y_{n-1} + \varepsilon \eta_n \quad \text{for } n \in \mathbb{N},$$

with parameters $v_0 > 0$, $\alpha, \varepsilon \in \mathbb{R}$, η_n i.i.d. $\sim N(0, 1)$, is a centered Gaussian process. The covariance function is given by

$$c(n, n+k) = v_0 + \varepsilon^2 n \quad \text{for any } n, k \geq 0 \quad \text{if } \alpha = 1,$$

and

$$c(n, n+k) = \alpha^k \cdot \left(\alpha^{2n} v_0 + (1 - \alpha^{2n}) \cdot \frac{\varepsilon^2}{1 - \alpha^2} \right) \quad \text{for } n, k \geq 0 \quad \text{otherwise.}$$

This is easily verified by induction. We now consider some special cases:

$\alpha = 0$: In this case $Y_n = \varepsilon \eta_n$. Hence (Y_n) is a *white noise*, i.e., a sequence of independent normal random variables, and

$$\text{Cov}[Y_n, Y_m] = \varepsilon^2 \cdot \delta_{n,m} \quad \text{for any } n, m \geq 1.$$

$\alpha = 1$: Here $Y_n = Y_0 + \varepsilon \sum_{i=1}^n \eta_i$, i.e., the process (Y_n) is a *Gaussian Random Walk*, and

$$\text{Cov}[Y_n, Y_m] = v_0 + \varepsilon^2 \cdot \min(n, m) \quad \text{for any } n, m \geq 0.$$

We will see a corresponding expression for the covariances of Brownian motion.

$\alpha < 1$: For $\alpha < 1$, the covariances $\text{Cov}[Y_n, Y_{n+k}]$ decay exponentially fast as $k \rightarrow \infty$. If $v_0 = \frac{\varepsilon^2}{1-\alpha^2}$, then the covariance function is translation invariant:

$$c(n, n+k) = \frac{\varepsilon^2 \alpha^k}{1-\alpha^2} \quad \text{for any } n, k \geq 0.$$

Therefore, in this case the process (Y_n) is *stationary*, i.e., $(Y_{n+k})_{n \geq 0} \sim (Y_n)_{n \geq 0}$ for all $k \geq 0$.

Brownian motion is our first example of a nontrivial Gaussian process in continuous time. In fact, we have:

Theorem 1.8 (Gaussian characterization of Brownian motion). *A real-valued stochastic process $(B_t)_{t \in [0, \infty)}$ with continuous sample paths $t \mapsto B_t(\omega)$ and $B_0 = 0$ is a Brownian motion if and only if (B_t) is a centered Gaussian process with covariances*

$$\text{Cov}[B_s, B_t] = \min(s, t) \quad \text{for any } s, t \geq 0. \quad (1.2.4)$$

Proof. For a Brownian motion (B_t) and $0 = t_0 < t_1 < \dots < t_n$, the increments $B_{t_i} - B_{t_{i-1}}$, $1 \leq i \leq n$, are independent random variables with distribution $N(0, t_i - t_{i-1})$. Hence,

$$(B_{t_1} - B_{t_0}, \dots, B_{t_n} - B_{t_{n-1}}) \sim \bigotimes_{i=1}^n N(0, t_i - t_{i-1}),$$

which is a multinormal distribution. Since $B_{t_0} = B_0 = 0$, we see that

$$\begin{pmatrix} B_{t_1} \\ \vdots \\ B_{t_n} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 \\ & & \ddots & & & \\ & & & \ddots & & \\ 1 & 1 & 1 & \dots & 1 & 0 \\ 1 & 1 & 1 & \dots & 1 & 1 \end{pmatrix} \begin{pmatrix} B_{t_1} - B_{t_0} \\ \vdots \\ B_{t_n} - B_{t_{n-1}} \end{pmatrix}$$

also has a multivariate normal distribution, i.e., (B_t) is a Gaussian process. Moreover, since $B_t = B_t - B_0$, we have $E[B_t] = 0$ and

$$\text{Cov}[B_s, B_t] = \text{Cov}[B_s, B_s] + \text{Cov}[B_s, B_t - B_s] = \text{Var}[B_s] = s$$

for any $0 \leq s \leq t$, i.e., (1.2.4) holds.

Conversely, if (B_t) is a centered Gaussian process satisfying (1.2.4), then for any $0 = t_0 < t_1 < \dots < t_n$, the vector $(B_{t_1} - B_{t_0}, \dots, B_{t_n} - B_{t_{n-1}})$ has a multivariate normal distribution with

$$E[B_{t_i} - B_{t_{i-1}}] = E[B_{t_i}] - E[B_{t_{i-1}}] = 0, \quad \text{and}$$

$$\begin{aligned} \text{Cov}[B_{t_i} - B_{t_{i-1}}, B_{t_j} - B_{t_{j-1}}] &= \min(t_i, t_j) - \min(t_i, t_{j-1}) \\ &\quad - \min(t_{i-1}, t_j) + \min(t_{i-1}, t_{j-1}) \\ &= (t_i - t_{i-1}) \cdot \delta_{i,j} \quad \text{for any } i, j = 1, \dots, n. \end{aligned}$$

Hence by Theorem 1.5 (3), the increments $B_{t_i} - B_{t_{i-1}}, 1 \leq i \leq n$, are independent with distribution $N(0, t_i - t_{i-1})$, i.e., (B_t) is a Brownian motion. \square

Symmetries of Brownian motion

A first important consequence of the Gaussian characterization of Brownian motion are several symmetry properties of Wiener measure:

Theorem 1.9 (Invariance properties of Wiener measure). *Let $(B_t)_{t \geq 0}$ be a Brownian motion starting at 0 defined on a probability space (Ω, \mathcal{A}, P) . Then the following processes are again Brownian motions:*

- (1). $(-B_t)_{t \geq 0}$ (Reflection invariance)
- (2). $(B_{t+h} - B_h)_{t \geq 0}$ for any $h \geq 0$ (Stationarity)
- (3). $(a^{-1/2} B_{at})_{t \geq 0}$ for any $a > 0$ (Scale invariance)
- (4). The time inversion $(\tilde{B}_t)_{t \geq 0}$ defined by

$$\tilde{B}_0 = 0, \quad \tilde{B}_t = t \cdot B_{1/t} \quad \text{for } t > 0.$$

Proof. The proofs of (1), (2) and (3) are left as an exercise to the reader. To show (4), we first note that for each $n \in \mathbb{N}$ and $0 \leq t_1 < \dots < t_n$, the vector $(\tilde{B}_{t_1}, \dots, \tilde{B}_{t_n})$ has a multivariate normal distribution since it is a linear transformation of $(B_{1/t_1}, \dots, B_{1/t_n})$, $(B_0, B_{1/t_2}, \dots, B_{1/t_n})$ respectively. Moreover,

$$\begin{aligned} E[\tilde{B}_t] &= 0 \quad \text{for any } t \geq 0, \\ \text{Cov}[\tilde{B}_s, \tilde{B}_t] &= st \cdot \text{Cov}[B_{1/s}, B_{1/t}] \\ &= st \cdot \min\left(\frac{1}{s}, \frac{1}{t}\right) = \min(t, s) \quad \text{for any } s, t > 0, \quad \text{and} \\ \text{Cov}[\tilde{B}_0, \tilde{B}_t] &= 0 \quad \text{for any } t \geq 0. \end{aligned}$$

Hence $(\tilde{B}_t)_{t \geq 0}$ is a centered Gaussian process with the covariance function of Brownian motion. By Theorem 1.8, it only remains to show that P -almost every sample path $t \mapsto \tilde{B}_t(\omega)$ is continuous. This is obviously true for $t > 0$. Furthermore, since the finite dimensional marginals of the processes $(\tilde{B}_t)_{t \geq 0}$ and $(B_t)_{t \geq 0}$ are multivariate normal distributions with the same means and covariances, the distributions of $(\tilde{B}_t)_{t \geq 0}$ and $(B_t)_{t \geq 0}$ on the product space $\mathbb{R}^{(0, \infty)}$ endowed with the product σ -algebra generated by the cylinder sets agree. To prove continuity at 0 we note that the set

$$\left\{ x : (0, \infty) \rightarrow \mathbb{R} \left| \lim_{\substack{t \searrow 0 \\ t \in \mathbb{Q}}} x_t = 0 \right. \right\}$$

is measurable w.r.t. the product σ -algebra on $\mathbb{R}^{(0, \infty)}$. Therefore,

$$P \left[\lim_{\substack{t \searrow 0 \\ t \in \mathbb{Q}}} \tilde{B}_t = 0 \right] = P \left[\lim_{\substack{t \searrow 0 \\ t \in \mathbb{Q}}} B_t = 0 \right] = 1.$$

Since \tilde{B}_t is almost surely continuous for $t > 0$, we can conclude that outside a set of measure zero,

$$\sup_{s \in (0, t)} |\tilde{B}_s| = \sup_{s \in (0, t) \cap \mathbb{Q}} |\tilde{B}_s| \longrightarrow 0 \quad \text{as } t \searrow 0,$$

i.e., $t \mapsto \tilde{B}_t$ is almost surely continuous at 0 as well. \square

Remark (Long time asymptotics versus local regularity, LLN). The time inversion invariance of Wiener measure enables us to translate results on the long time asymptotics of Brownian motion ($t \nearrow \infty$) into local regularity results for Brownian paths ($t \searrow 0$) and vice versa. For example, the continuity of the process (\tilde{B}_t) at 0 is equivalent to the *law of large numbers*:

$$P \left[\lim_{t \rightarrow \infty} \frac{1}{t} B_t = 0 \right] = P \left[\lim_{s \searrow 0} s B_{1/s} = 0 \right] = 1.$$

At first glance, this looks like a simple proof of the LLN. However, the argument is based on the existence of a continuous Brownian motion, and the existence proof requires similar arguments as a direct proof of the law of large numbers.

Wiener measure as a Gaussian measure, path integral heuristics

Wiener measure (with start at 0) is the unique probability measure μ on the continuous path space $C([0, \infty), \mathbb{R}^d)$ such that the coordinate process

$$X_t : C([0, \infty), \mathbb{R}^d) \rightarrow \mathbb{R}^d, \quad X_t(x) = x_t,$$

is a Brownian motion starting at 0. By Theorem 1.8, Wiener measure is a centered **Gaussian measure** on the infinite dimensional space $C([0, \infty), \mathbb{R}^d)$, i.e., for any $n \in \mathbb{N}$ and $t_1, \dots, t_n \in \mathbb{R}_+$, $(X_{t_1}, \dots, X_{t_n})$ is normally distributed with mean 0. We now "derive" a heuristic representation of Wiener measure that is not mathematically rigorous but nevertheless useful:

Fix a constant $T > 0$. Then for $0 = t_0 < t_1 < \dots < t_n \leq T$, the distribution of $(X_{t_1}, \dots, X_{t_n})$ w.r.t. Wiener measure is

$$\mu_{t_1, \dots, t_n}(dx_{t_1}, \dots, dx_{t_n}) = \frac{1}{Z(t_1, \dots, t_n)} \exp \left(-\frac{1}{2} \sum_{i=1}^n \frac{|x_{t_i} - x_{t_{i-1}}|^2}{t_i - t_{i-1}} \right) \prod_{i=1}^n dx_{t_i}, \quad (1.2.5)$$

where $Z(t_1, \dots, t_n)$ is an appropriate finite normalization constant, and $x_0 := 0$. Now choose a sequence $(\tau_k)_{k \in \mathbb{N}}$ of partitions $0 = t_0^{(k)} < t_1^{(k)} < \dots < t_{n(k)}^{(k)} = T$ of the interval

$[0, T]$ such that the mesh size $\max_i |t_{i+1}^{(k)} - t_i^{(k)}|$ tends to zero. Taking informally the limit in (1.2.5), we obtain the heuristic asymptotic representation

$$\mu(dx) = \frac{1}{Z_\infty} \exp\left(-\frac{1}{2} \int_0^T \left|\frac{dx}{dt}\right|^2 dt\right) \delta_0(dx_0) \prod_{t \in (0, T]} dx_t \quad (1.2.6)$$

for Wiener measure on continuous paths $x : [0, T] \rightarrow \mathbb{R}^d$ with a "normalizing constant" Z_∞ . Trying to make the informal expression (1.2.6) rigorous fails for several reasons:

- The normalizing constant $Z_\infty = \lim_{k \rightarrow \infty} Z(t_1^{(k)}, \dots, t_n^{(k)})$ is infinite.
- The integral $\int_0^T \left|\frac{dx}{dt}\right|^2 dt$ is also infinite for μ -almost every path x , since typical paths of Brownian motion are nowhere differentiable, cf. below.
- The product measure $\prod_{t \in (0, T]} dx_t$ can be defined on cylinder sets but an extension to the σ -algebra generated by the coordinate maps on $C([0, \infty), \mathbb{R}^d)$ does not exist.

Hence there are several infinities involved in the informal expression (1.2.6). These infinities magically balance each other such that the measure μ is well defined in contrast to all of the factors on the right hand side.

In physics, R. Feynman introduced correspondingly integrals w.r.t. "Lebesgue measure on path space", cf. e.g. the famous Feynman Lecture notes [...], or Glimm and Jaffe [...].

Although not mathematically rigorous, the heuristic expression (1.2.5) can be a very useful guide for intuition. Note for example that (1.2.5) takes the form

$$\mu(dx) \propto \exp(-\|x\|_H^2/2) \lambda(dx), \quad (1.2.7)$$

where $\|x\|_H = (x, x)_H^{1/2}$ is the norm induced by the inner product

$$(x, y)_H = \int_0^T \frac{dx}{dt} \frac{dy}{dt} dt \quad (1.2.8)$$

of functions $x, y : [0, T] \rightarrow \mathbb{R}^d$ vanishing at 0, and λ is a corresponding "infinite-dimensional Lebesgue measure" (which does not exist!). The vector space

$$H = \left\{ x : [0, T] \rightarrow \mathbb{R}^d : x(0) = 0, x \text{ is absolutely continuous with } \frac{dx}{dt} \in L^2 \right\}$$

is a Hilbert space w.r.t. the inner product (1.2.8). Therefore, (1.2.7) suggests to consider Wiener measure as a *standard normal distribution on H* . It turns out that this idea can be made rigorous although not as easily as one might think at first glance. The difficulty is that a standard normal distribution on an infinite-dimensional Hilbert space does not exist on the space itself but only on a larger space. In particular, we will see in the next sections that Wiener measure μ can indeed be realized on the continuous path space $C([0, T], \mathbb{R}^d)$, but μ -almost every path is not contained in H !

Remark (Infinite-dimensional standard normal distributions). The fact that a standard normal distribution on an infinite dimensional separable Hilbert space H can not be realized on the space H itself can be easily seen by contradiction: Suppose that μ is a standard normal distribution on H , and $e_n, n \in \mathbb{N}$, are infinitely many orthonormal vectors in H . Then by rotational symmetry, the balls

$$B_n = \left\{ x \in H : \|x - e_n\|_H < \frac{1}{2} \right\}, \quad n \in \mathbb{N},$$

should all have the same measure. On the other hand, the balls are disjoint. Hence by σ -additivity,

$$\sum_{n=1}^{\infty} \mu[B_n] = \mu \left[\bigcup B_n \right] \leq \mu[H] = 1,$$

and therefore $\mu[B_n] = 0$ for all $n \in \mathbb{N}$. A scaling argument now implies

$$\mu[\{x \in H : \|x - h\| \leq \|h\|/2\}] = 0 \quad \text{for all } h \in H,$$

and hence $\mu \equiv 0$.

1.3 The Wiener-Lévy Construction

In this section we discuss how to construct Brownian motion as a random superposition of deterministic paths. The idea already goes back to N. Wiener, who constructed

Brownian motion as a random Fourier series. The approach described here is slightly different and due to P. Lévy: The idea is to approximate the paths of Brownian motion on a finite time interval by their piecewise linear interpolations w.r.t. the sequence of dyadic partitions. This corresponds to a development of the Brownian paths w.r.t. Schauder functions ("wavelets") which turns out to be very useful for many applications including numerical simulations.

Our aim is to construct a one-dimensional Brownian motion B_t starting at 0 for $t \in [0, 1]$. By stationarity and independence of the increments, a Brownian motion defined for all $t \in [0, \infty)$ can then easily be obtained from infinitely many independent copies of Brownian motion on $[0, 1]$. We are hence looking for a random variable

$$B = (B_t)_{t \in [0,1]} : \Omega \longrightarrow C([0, 1])$$

defined on a probability space (Ω, \mathcal{A}, P) such that the distribution $P \circ B^{-1}$ is Wiener measure μ on the continuous path space $C([0, 1])$.

A first attempt

Recall that μ_0 should be a kind of standard normal distribution w.r.t. the inner product

$$(x, y)_H = \int_0^1 \frac{dx}{dt} \frac{dy}{dt} dt \quad (1.3.1)$$

on functions $x, y : [0, 1] \rightarrow \mathbb{R}$. Therefore, we could try to define

$$B_t(\omega) := \sum_{i=1}^{\infty} Z_i(\omega) e_i(t) \quad \text{for } t \in [0, 1] \text{ and } \omega \in \Omega, \quad (1.3.2)$$

where $(Z_i)_{i \in \mathbb{N}}$ is a sequence of independent standard normal random variables, and $(e_i)_{i \in \mathbb{N}}$ is an orthonormal basis in the Hilbert space

$$H = \{x : [0, 1] \rightarrow \mathbb{R} \mid x(0) = 0, x \text{ is absolutely continuous with } (x, x)_H < \infty\}. \quad (1.3.3)$$

However, the resulting series approximation does not converge in H :

Theorem 1.10. *Suppose $(e_i)_{i \in \mathbb{N}}$ is a sequence of orthonormal vectors in a Hilbert space H and $(Z_i)_{i \in \mathbb{N}}$ is a sequence of i.i.d. random variables with $P[Z_i \neq 0] > 0$. Then the series $\sum_{i=1}^{\infty} Z_i(\omega)e_i$ diverges with probability 1 w.r.t. the norm on H .*

Proof. By orthonormality and by the law of large numbers,

$$\left\| \sum_{i=1}^n Z_i(\omega)e_i \right\|_H^2 = \sum_{i=1}^n Z_i(\omega)^2 \longrightarrow \infty$$

P -almost surely as $n \rightarrow \infty$. □

The Theorem again reflects the fact that a standard normal distribution on an infinite-dimensional Hilbert space can not be realized on the space itself.

To obtain a positive result, we will replace the norm

$$\|x\|_H = \left(\int_0^1 \left| \frac{dx}{dt} \right|^2 dt \right)^{\frac{1}{2}}$$

on H by the supremum norm

$$\|x\|_{\text{sup}} = \sup_{t \in [0,1]} |x_t|,$$

and correspondingly the Hilbert space H by the Banach space $C([0, 1])$. Note that the supremum norm is weaker than the H -norm. In fact, for $x \in H$ and $t \in [0, 1]$, the Cauchy-Schwarz inequality implies

$$|x_t|^2 = \left| \int_0^t x'_s ds \right|^2 \leq t \cdot \int_0^t |x'_s|^2 ds \leq \|x\|_H^2,$$

and therefore

$$\|x\|_{\text{sup}} \leq \|x\|_H \quad \text{for any } x \in H.$$


There are two choices for an orthonormal basis of the Hilbert space H that are of particular interest: The first is the Fourier basis given by

$$e_0(t) = t, \quad e_n(t) = \frac{\sqrt{2}}{\pi n} \sin(\pi n t) \quad \text{for } n \geq 1.$$

With respect to this basis, the series in (1.3.2) is a Fourier series with random coefficients. Wiener's original construction of Brownian motion is based on a *random Fourier series*. A second convenient choice is the basis of *Schauder functions* ("wavelets") that has been used by P. Lévy to construct Brownian motion. Below, we will discuss Lévy's construction in detail. In particular, we will prove that for the Schauder functions, the series in (1.3.2) converges almost surely w.r.t. the supremum norm towards a continuous (but not absolutely continuous) random path $(B_t)_{t \in [0,1]}$. It is then not difficult to conclude that $(B_t)_{t \in [0,1]}$ is indeed a Brownian motion.

The Wiener-Lévy representation of Brownian motion

Before carrying out Lévy's construction of Brownian motion, we introduce the Schauder functions, and we show how to expand a given Brownian motion w.r.t. this basis of function space. Suppose we would like to approximate the paths $t \mapsto B_t(\omega)$ of a Brownian motion by their piecewise linear approximations adapted to the sequence of dyadic partitions of the interval $[0, 1]$.





An obvious advantage of this approximation over a Fourier expansion is that the values of the approximating functions at the dyadic points remain fixed once the approximating


partition is fine enough. The piecewise linear approximations of a continuous function on $[0, 1]$ correspond to a series expansion w.r.t. the base functions

$$e(t) = t, \text{ and}$$

$$e_{n,k}(t) = 2^{-n/2} e_{0,0}(2^n t - k), \quad n = 0, 1, 2, \dots, k = 0, 1, 2, \dots, 2^n - 1, \quad , \text{ where}$$

$$e_{0,0}(t) = \min(t, 1-t)^+ = \begin{cases} t & \text{for } t \in [0, 1/2] \\ 1-t & \text{for } t \in (1/2, 1] \\ 0 & \text{for } t \in \mathbb{R} \setminus [0, 1] \end{cases} .$$





The functions $e_{n,k}$ ($n \geq 0, 0 \leq k < 2^n$) are called **Schauder functions**. It is rather obvious that piecewise linear approximation w.r.t. the dyadic partitions corresponds to the expansion of a function $x \in C([0,1])$ with $x(0) = 0$ in the basis given by $e(t)$ and the Schauder functions. The normalization constants in defining the functions $e_{n,k}$ have been chosen in such a way that the $e_{n,k}$ are orthonormal w.r.t. the H -inner product introduced above.

Definition. A sequence $(e_i)_{i \in \mathbb{N}}$ of vectors in an infinite-dimensional Hilbert space H is called an **orthonormal basis** (or **complete orthonormal system**) of H if and only if

(1). *Orthonormality:* $(e_i, e_j) = \delta_{ij}$ for any $i, j \in \mathbb{N}$, and

(2). *Completeness:* Any $h \in H$ can be expressed as

$$h = \sum_{i=1}^{\infty} (h, e_i)_H e_i.$$

Remark (Equivalent characterizations of orthonormal bases). Let $e_i, i \in \mathbb{N}$, be orthonormal vectors in a Hilbert space H . Then the following conditions are equivalent:

(1). $(e_i)_{i \in \mathbb{N}}$ is an orthonormal basis of H .

(2). The linear span

$$\text{span}\{e_i \mid i \in \mathbb{N}\} = \left\{ \sum_{i=1}^k c_i e_i \mid k \in \mathbb{N}, c_1, \dots, c_k \in \mathbb{R} \right\}$$

is a dense subset of H .

(3). There is no element $x \in H, x \neq 0$, such that $(x, e_i)_H = 0$ for every $i \in \mathbb{N}$.

(4). For any element $x \in H$, Parseval's relation

$$\|x\|_H^2 = \sum_{i=1}^{\infty} (x, e_i)_H^2 \quad (1.3.4)$$

holds.

(5). For any $x, y \in H$,

$$(x, y)_H = \sum_{i=1}^{\infty} (x, e_i)_H (y, e_i)_H. \quad (1.3.5)$$

For the proofs we refer to any book on functional analysis, cf. e.g. [Reed and Simon: Methods of modern mathematical physics, Vol. I].


Lemma 1.11. *The Schauder functions e and $e_{n,k}$ ($n \geq 0, 0 \leq k < 2^n$) form an orthonormal basis in the Hilbert space H defined by (1.3.3).*

Proof. By definition of the inner product on H , the linear map d/dt which maps an absolutely continuous function $x \in H$ to its derivative $x' \in L^2(0, 1)$ is an isometry from H onto $L^2(0, 1)$, i.e.,

$$(x, y)_H = (x', y')_{L^2(0,1)} \quad \text{for any } x, y \in H.$$

The derivatives of the Schauder functions are the Haar functions

$$\begin{aligned} e'(t) &\equiv 1, \\ e'_{n,k}(t) &= 2^{n/2} (I_{[k \cdot 2^{-n}, (k+1/2) \cdot 2^{-n})}(t) - I_{[(k+1/2) \cdot 2^{-n}, (k+1) \cdot 2^{-n})}(t)) \quad \text{for a.e. } t. \end{aligned}$$



It is easy to see that these functions form an orthonormal basis in $L^2(0, 1)$. In fact, orthonormality w.r.t. the L^2 inner product can be verified directly. Moreover, the linear span of the functions e' and $e'_{n,k}$ for $n = 0, 1, \dots, m$ and $k = 0, 1, \dots, 2^n - 1$ consists of all step functions that are constant on each dyadic interval $[j \cdot 2^{-(m+1)}, (j+1) \cdot 2^{-(m+1)})$. An arbitrary function in $L^2(0, 1)$ can be approximated by dyadic step functions w.r.t. the L^2 norm. This follows for example directly from the L^2 martingale convergence Theorem, cf. ... below. Hence the linear span of e' and the Haar functions $e'_{n,k}$ is dense in $L^2(0, 1)$, and therefore these functions form an orthonormal basis of the Hilbert space $L^2(0, 1)$. Since $x \mapsto x'$ is an isometry from H onto $L^2(0, 1)$, we can conclude that e and the Schauder functions $e_{n,k}$ form an orthonormal basis of H . \square

The expansion of a function $x : [0, 1] \rightarrow \mathbb{R}$ in the basis of Schauder functions can now be made explicit. The coefficients of a function $x \in H$ in the expansion are

$$(x, e)_H = \int_0^1 x' e' dt = \int_0^1 x' dt = x(1) - x(0) = x(1)$$

$$\begin{aligned}
(x, e_{n,k})_H &= \int_0^1 x' e'_{n,k} dt = 2^{n/2} \int_0^1 x'(t) e'_{0,0}(2^n t - k) dt \\
&= 2^{n/2} \left[(x((k + \frac{1}{2}) \cdot 2^{-n}) - x(k \cdot 2^{-n})) - (x((k + 1) \cdot 2^{-n}) - x((k + \frac{1}{2}) \cdot 2^{-n})) \right].
\end{aligned}$$

Theorem 1.12. *Let $x \in C([0, 1])$. Then the expansion*

$$x(t) = x(1)e(t) - \sum_{n=0}^{\infty} \sum_{k=0}^{2^n-1} 2^{n/2} \Delta_{n,k} x \cdot e_{n,k}(t),$$

$$\Delta_{n,k} x = \left[(x((k + 1) \cdot 2^{-n}) - x((k + \frac{1}{2}) \cdot 2^{-n})) - (x((k + \frac{1}{2}) \cdot 2^{-n}) - x(k \cdot 2^{-n})) \right]$$

holds w.r.t. uniform convergence on $[0, 1]$. For $x \in H$ the series also converges w.r.t. the stronger H -norm.

Proof. It can be easily verified that by definition of the Schauder functions, for each $m \in \mathbb{N}$ the partial sum

$$x^{(m)}(t) := x(1)e(t) - \sum_{n=0}^m \sum_{k=0}^{2^n-1} 2^{n/2} \Delta_{n,k} x \cdot e_{n,k}(t) \quad (1.3.6)$$

is the polygonal interpolation of $x(t)$ w.r.t. the $(m + 1)$ -th dyadic partition of the interval $[0, 1]$. Since the function x is uniformly continuous on $[0, 1]$, the polygonal interpolations converge uniformly to x . This proves the first statement. Moreover, for $x \in H$, the series is the expansion of x in the orthonormal basis of H given by the Schauder functions, and therefore it also converges w.r.t. the H -norm. \square

Applying the expansion to the paths of a Brownian motions, we obtain:

Corollary 1.13 (Wiener-Lévy representation). *For a Brownian motion $(B_t)_{t \in [0,1]}$ the series representation*

$$B_t(\omega) = Z(\omega)e(t) + \sum_{n=0}^{\infty} \sum_{k=0}^{2^n-1} Z_{n,k}(\omega) e_{n,k}(t), \quad t \in [0, 1], \quad (1.3.7)$$

holds w.r.t. uniform convergence on $[0, 1]$ for P -almost every $\omega \in \Omega$, where

$$Z := B_1, \quad \text{and} \quad Z_{n,k} := -2^{n/2} \Delta_{n,k} B \quad (n \geq 0, 0 \leq k \leq 2^n - 1)$$

are independent random variables with standard normal distribution.

Proof. It only remains to verify that the coefficients Z and $Z_{n,k}$ are independent with standard normal distribution. A vector given by finitely many of these random variables has a multivariate normal distribution, since it is a linear transformation of increments of the Brownian motion B_t . Hence it suffices to show that the random variables are uncorrelated with variance 1. This is left as an exercise to the reader. \square

Lévy's construction of Brownian motion

The series representation (1.3.7) can be used to construct Brownian motion starting from independent standard normal random variables. The resulting construction does not only prove existence of Brownian motion but it is also very useful for numerical implementations:

Theorem 1.14 (P. Lévy 1948). *Let Z and $Z_{n,k}$ ($n \geq 0, 0 \leq k \leq 2^n - 1$) be independent standard normally distributed random variables on a probability space (Ω, \mathcal{A}, P) . Then the series in (1.3.7) converges uniformly on $[0, 1]$ with probability 1. The limit process $(B_t)_{t \in [0,1]}$ is a Brownian motion.*

The convergence proof relies on a combination of the Borel-Cantelli Lemma and the Weierstrass criterion for uniform convergence of series of functions. Moreover, we will need the following result to identify the limit process as a Brownian motion:

Lemma 1.15 (Parseval relation for Schauder functions). *For any $s, t \in [0, 1]$,*

$$e(t)e(s) + \sum_{n=0}^{\infty} \sum_{k=0}^{2^n-1} e_{n,k}(t)e_{n,k}(s) = \min(t, s).$$

Proof. Note that for $g \in H$ and $s \in [0, 1]$, we have

$$g(s) = g(s) - g(0) = \int_0^1 g' \cdot I_{(0,s)} = (g, h^{(s)})_H,$$

where $h^{(s)}(t) := \int_0^t I_{(0,s)} = \min(s, t)$. Hence the Parseval relation (1.3.4) applied to the functions $h^{(s)}$ and $h^{(t)}$ yields

$$\begin{aligned} e(t)e(s) + \sum_{n,k} e_{n,k}(t)e_{n,k}(s) &= (e, h^{(t)})(e, h^{(s)}) + \sum_{n,k} (e_{n,k}, h^{(t)})(e_{n,k}, h^{(s)}) \\ &= (h^{(t)}, h^{(s)}) = \int_0^1 I_{(0,t)}I_{(0,s)} = \min(t, s). \end{aligned}$$

□

Proof of Theorem 1.14. We proceed in 4 steps:

- (1). *Uniform convergence for P-a.e. ω :* By the Weierstrass criterion, a series of functions converges uniformly if the sum of the supremum norms of the summands is finite. To apply the criterion, we note that for any fixed $t \in [0, 1]$ and $n \in \mathbb{N}$, only one of the functions $e_{n,k}$, $k = 0, 1, \dots, 2^n - 1$, does not vanish at t . Moreover, $|e_{n,k}(t)| \leq 2^{-n/2}$. Hence

$$\sup_{t \in [0,1]} \left| \sum_{k=0}^{2^n-1} Z_{n,k}(\omega) e_{n,k}(t) \right| \leq 2^{-n/2} \cdot M_n(\omega), \quad (1.3.8)$$

where

$$M_n := \max_{0 \leq k < 2^n} |Z_{n,k}|.$$

We now apply the Borel-Cantelli Lemma to show that with probability 1, M_n grows at most linearly. Let Z denote a standard normal random variable. Then we have

$$\begin{aligned} P[M_n > n] &\leq 2^n \cdot P[|Z| > n] \leq \frac{2^n}{n} \cdot E[|Z|; |Z| > n] \\ &= \frac{2 \cdot 2^n}{n \cdot \sqrt{2\pi}} \int_n^\infty x e^{-x^2/2} dx = \sqrt{\frac{2}{\pi}} \frac{2^n}{n} \cdot e^{-n^2/2} \end{aligned}$$

for any $n \in \mathbb{N}$. Since the sequence on the right hand side is summable, $M_n \leq n$ holds eventually with probability one. Therefore, the sequence on the right hand side of (1.3.8) is also summable for P -almost every ω . Hence, by (1.3.8) and the Weierstrass criterion, the partial sums

$$B_t^{(m)}(\omega) = Z(\omega)e(t) + \sum_{n=0}^m \sum_{k=0}^{2^n-1} Z_{n,k}(\omega)e_{n,k}(t), \quad m \in \mathbb{N},$$

converge almost surely uniformly on $[0, 1]$. Let

$$B_t = \lim_{m \rightarrow \infty} B_t^{(m)}$$

denote the almost surely defined limit.

- (2). *L^2 convergence for fixed t :* We now want to prove that the limit process (B_t) is a Brownian motion, i.e., a continuous Gaussian process with $E[B_t] = 0$ and $\text{Cov}[B_t, B_s] = \min(t, s)$ for any $t, s \in [0, 1]$. To compute the covariances we first show that for a given $t \in [0, 1]$ the series approximation $B_t^{(m)}$ of B_t converges also in L^2 . Let $l, m \in \mathbb{N}$ with $l < m$. Since the $Z_{n,k}$ are independent (and hence uncorrelated) with variance 1, we have

$$E[(B_t^{(m)} - B_t^{(l)})^2] = E \left[\left(\sum_{n=l+1}^m \sum_{k=0}^{2^n-1} Z_{n,k} e_{n,k}(t) \right)^2 \right] = \sum_{n=l+1}^m \sum_k e_{n,k}(t)^2.$$

The right hand side converges to 0 as $l, m \rightarrow \infty$ since $\sum_{n,k} e_{n,k}(t)^2 < \infty$ by Lemma 1.15. Hence $B_t^{(m)}, m \in \mathbb{N}$, is a Cauchy sequence in $L^2(\Omega, \mathcal{A}, P)$. Since $B_t = \lim_{m \rightarrow \infty} B_t^{(m)}$ almost surely, we obtain

$$B_t^{(m)} \xrightarrow{m \rightarrow \infty} B_t \quad \text{in } L^2(\Omega, \mathcal{A}, P).$$

- (3). *Expectations and Covariances:* By the L^2 convergence we obtain for any $s, t \in [0, 1]$:

$$\begin{aligned} E[B_t] &= \lim_{m \rightarrow \infty} E[B_t^{(m)}] = 0, \quad \text{and} \\ \text{Cov}[B_t, B_s] &= E[B_t B_s] = \lim_{m \rightarrow \infty} E[B_t^{(m)} B_s^{(m)}] \\ &= e(t)e(s) + \lim_{m \rightarrow \infty} \sum_{n=0}^m \sum_{k=0}^{2^n-1} e_{n,k}(t)e_{n,k}(s). \end{aligned}$$

Here we have used again that the random variables Z and $Z_{n,k}$ are independent with variance 1. By Parseval's relation (Lemma 1.15), we conclude

$$\text{Cov}[B_t, B_s] = \min(t, s).$$

Since the process $(B_t)_{t \in [0,1]}$ has the right expectations and covariances, and, by construction, almost surely continuous paths, it only remains to show that (B_t) is a Gaussian process in order to complete the proof:

- (4). $(B_t)_{t \in [0,1]}$ is a Gaussian process: We have to show that $(B_{t_1}, \dots, B_{t_l})$ has a multivariate normal distribution for any $0 \leq t_1 < \dots < t_l \leq 1$. By Theorem 1.5, it suffices to verify that any linear combination of the components is normally distributed. This holds by the next Lemma since

$$\sum_{j=1}^l p_j B_{t_j} = \lim_{m \rightarrow \infty} \sum_{j=1}^l p_j B_{t_j}^{(m)} \quad P\text{-a.s.}$$

is an almost sure limit of normally distributed random variables for any $p_1, \dots, p_l \in \mathbb{R}$.

Combining Steps 3, 4 and the continuity of sample paths, we conclude that $(B_t)_{t \in [0,1]}$ is indeed a Brownian motion. \square

Lemma 1.16. *Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of normally distributed random variables defined on a joint probability space (Ω, \mathcal{A}, P) , and X_n converges almost surely to a random variable X . Then X is also normally distributed.*

Proof. Suppose $X_n \sim N(m_n, \sigma_n^2)$ with $m_n \in \mathbb{R}$ and $\sigma_n \in (0, \infty)$. By the Dominated Convergence Theorem,

$$E[e^{ipX}] = \lim_{n \rightarrow \infty} E[e^{ipX_n}] = \lim_{n \rightarrow \infty} e^{ipm_n} e^{-\frac{1}{2}\sigma_n^2 p^2}.$$


The limit on the right hand side only exists for all p , if either $\sigma_n \rightarrow \infty$, or the sequences σ_n and m_n both converge to finite limits $\sigma \in [0, \infty)$ and $m \in \mathbb{R}$. In the first case, the limit would equal 0 for $p \neq 0$ and 1 for $p = 0$. This is a contradiction, since

characteristic functions are always continuous. Hence the second case occurs, and, therefore

$$E[e^{ipX}] = e^{ipm - \frac{1}{2}\sigma^2 p^2} \quad \text{for any } p \in \mathbb{R},$$

i.e., $X \sim N(m, \sigma^2)$. □

So far, we have constructed Brownian motion only for $t \in [0, 1]$. Brownian motion on any finite time interval can easily be obtained from this process by rescaling. Brownian motion defined for all $t \in \mathbb{R}_+$ can be obtained by joining infinitely many Brownian motions on time intervals of length 1:



Theorem 1.17. Suppose that $B_t^{(1)}, B_t^{(2)}, \dots$ are independent Brownian motions starting at 0 defined for $t \in [0, 1]$. Then the process

$$B_t := B_{t - [t]}^{([t] + 1)} + \sum_{i=1}^{[t]} B_1^{(i)}, \quad t \geq 0,$$

is a Brownian motion defined for $t \in [0, \infty)$.

The proof is left as an exercise.

1.4 The Brownian Sample Paths

In this section we study some properties of Brownian sample paths in dimension one. We show that a typical Brownian path is nowhere differentiable, and Hölder-continuous with parameter α if and only if $\alpha < 1/2$. Furthermore, the set $\Lambda_a = \{t \geq 0 : B_t = a\}$ of all passage times of a given point $a \in \mathbb{R}$ is a fractal. We will show that almost surely, Λ_a has Lebesgue measure zero but any point in Λ_a is an accumulation point of Λ_a . We consider a one-dimensional Brownian motion $(B_t)_{t \geq 0}$ with $B_0 = 0$ defined on a probability space (Ω, \mathcal{A}, P) . Then:

Typical Brownian sample paths are nowhere differentiable

For any $t \geq 0$ and $h > 0$, the difference quotient $\frac{B_{t+h} - B_t}{h}$ is normally distributed with mean 0 and standard deviation

$$\sigma[(B_{t+h} - B_t)/h] = \sigma[B_{t+h} - B_t]/h = 1/\sqrt{h}.$$

This suggests that the derivative

$$\frac{d}{dt} B_t = \lim_{h \searrow 0} \frac{B_{t+h} - B_t}{h}$$

does not exist. Indeed, we have the following stronger statement.

Theorem 1.18 (Paley, Wiener, Zygmund 1933). *Almost surely, the Brownian sample path $t \mapsto B_t$ is nowhere differentiable, and*

$$\limsup_{s \searrow t} \left| \frac{B_s - B_t}{s - t} \right| = \infty \quad \text{for any } t \geq 0.$$

Note that, since there are uncountably many $t \geq 0$, the statement is stronger than claiming only the almost sure non-differentiability for any given $t \geq 0$.


Proof. It suffices to show that the set

$$N = \left\{ \omega \in \Omega \mid \exists t \in [0, T], k, L \in \mathbb{N} \forall s \in (t, t + \frac{1}{k}) : |B_s(\omega) - B_t(\omega)| \leq L|s - t| \right\}$$

is a null set for any $T \in \mathbb{N}$. Hence fix $T \in \mathbb{N}$, and consider $\omega \in N$. Then there exist $k, L \in \mathbb{N}$ and $t \in [0, T]$ such that

$$|B_s(\omega) - B_t(\omega)| \leq L \cdot |s - t| \quad \text{holds for } s \in (t, t + \frac{1}{k}). \quad (1.4.1)$$

To make use of the independence of the increments over disjoint intervals, we note that for any $n > 4k$, we can find an $i \in \{1, 2, \dots, nT\}$ such that the intervals $(\frac{i}{n}, \frac{i+1}{n})$, $(\frac{i+1}{n}, \frac{i+2}{n})$, and $(\frac{i+2}{n}, \frac{i+3}{n})$ are all contained in $(t, t + \frac{1}{k})$:



Hence by (1.4.1), the bound

$$\begin{aligned} \left| B_{\frac{i+1}{n}}(\omega) - B_{\frac{i}{n}}(\omega) \right| &\leq \left| B_{\frac{i+1}{n}}(\omega) - B_t(\omega) \right| + \left| B_t(\omega) - B_{\frac{i}{n}}(\omega) \right| \\ &\leq L \cdot \left(\frac{j+1}{n} - t \right) + L \cdot \left(\frac{j}{n} - t \right) \leq \frac{8L}{n} \end{aligned}$$

holds for $j = i, i + 1, i + 2$. Thus we have shown that N is contained in the set

$$\tilde{N} := \bigcup_{k, L \in \mathbb{N}} \bigcap_{n > 4k} \bigcup_{i=1}^{nT} \left\{ \left| B_{\frac{i+1}{n}} - B_{\frac{i}{n}} \right| \leq \frac{8L}{n} \text{ for } j = i, i + 1, i + 2 \right\}.$$

We now prove $P[\tilde{N}] = 0$. By independence and stationarity of the increments we have

$$\begin{aligned} & P \left[\left\{ \left| B_{\frac{i+1}{n}} - B_{\frac{i}{n}} \right| \leq \frac{8L}{n} \quad \text{for } j = i, i+1, i+2 \right\} \right] \\ &= P \left[\left| B_{\frac{1}{n}} \right| \leq \frac{8L}{n} \right]^3 = P \left[|B_1| \leq \frac{8L}{\sqrt{n}} \right]^3 \quad (1.4.2) \\ &\leq \left(\frac{1}{\sqrt{2\pi}} \frac{16L}{\sqrt{n}} \right)^3 = \frac{16^3}{\sqrt{2\pi}^3} \cdot \frac{L^3}{n^{3/2}} \end{aligned}$$

for any i and n . Here we have used that the standard normal density is bounded from above by $1/\sqrt{2\pi}$. By (1.4.2) we obtain

$$\begin{aligned} & P \left[\bigcap_{n>4k} \bigcup_{i=1}^{nT} \left\{ \left| B_{\frac{i+1}{n}} - B_{\frac{i}{n}} \right| \leq \frac{8L}{n} \quad \text{for } j = i, i+1, i+2 \right\} \right] \\ &\leq \frac{16^3}{\sqrt{2\pi}^3} \cdot \inf_{n>4k} nTL^3/n^{3/2} = 0. \end{aligned}$$

Hence, $P[\tilde{N}] = 0$, and therefore N is a null set. \square

Hölder continuity

The statement of Theorem 1.18 says that a typical Brownian path is not Lipschitz continuous on any non-empty open interval. On the other hand, the Wiener-Lévy construction shows that the sample paths are continuous. We can almost close the gap between these two statements by arguing in both cases slightly more carefully:

Theorem 1.19. *The following statements hold almost surely:*

(1). *For any $\alpha > 1/2$,*

$$\limsup_{s \searrow t} \frac{|B_s - B_t|}{|s - t|^\alpha} = \infty \quad \text{for all } t \geq 0.$$

(2). *For any $\alpha < 1/2$,*

$$\sup_{\substack{s, t \in [0, T] \\ s \neq t}} \frac{|B_s - B_t|}{|s - t|^\alpha} < \infty \quad \text{for all } T > 0.$$

Hence a typical Brownian path is nowhere Hölder continuous with parameter $\alpha > 1/2$, but it is Hölder continuous with parameter $\alpha < 1/2$ on any finite interval. The critical case $\alpha = 1/2$ is more delicate, and will be briefly discussed below.

Proof of Theorem 1.19. The first statement can be shown by a similar argument as in the proof of Theorem 1.18. The details are left to the reader.

To prove the second statement for $T = 1$, we use the Wiener-Lévy representation

$$B_t = Z \cdot t + \sum_{n=0}^{\infty} \sum_{k=0}^{2^n-1} Z_{n,k} e_{n,k}(t) \quad \text{for any } t \in [0, 1]$$

with independent standard normal random variables $Z, Z_{n,k}$. For $t, s \in [0, 1]$ we obtain

$$|B_t - B_s| \leq |Z| \cdot |t - s| + \sum_n M_n \sum_k |e_{n,k}(t) - e_{n,k}(s)|,$$

where $M_n := \max_k |Z_{n,k}|$ as in the proof of Theorem 1.14. We have shown above that by the Borel-Cantelli Lemma, $M_n \leq n$ eventually with probability one, and hence

$$M_n(\omega) \leq C(\omega) \cdot n$$

for some almost surely finite constant $C(\omega)$. Moreover, note that for each s, t and n , at most two summands in $\sum_k |e_{n,k}(t) - e_{n,k}(s)|$ do not vanish. Since $|e_{n,k}(t)| \leq \frac{1}{2} \cdot 2^{-n/2}$ and $|e'_{n,k}(t)| \leq 2^{n/2}$, we obtain the estimates

$$|e_{n,k}(t) - e_{n,k}(s)| \leq 2^{-n/2}, \quad \text{and} \quad (1.4.3)$$

$$|e_{n,k}(t) - e_{n,k}(s)| \leq 2^{n/2} \cdot |t - s|. \quad (1.4.4)$$

For given $s, t \in [0, 1]$, we now choose $N \in \mathbb{N}$ such that

$$2^{-N} \leq |t - s| < 2^{1-N}. \quad (1.4.5)$$

By applying (1.4.3) for $n > N$ and (1.4.4) for $n \leq N$, we obtain

$$|B_t - B_s| \leq |Z| \cdot |t - s| + 2C \cdot \left(\sum_{n=1}^N n 2^{n/2} \cdot |t - s| + \sum_{n=N+1}^{\infty} n 2^{-n/2} \right).$$

By (1.4.5) the sums on the right hand side can both be bounded by a constant multiple of $|t - s|^\alpha$ for any $\alpha < 1/2$. This proves that $(B_t)_{t \in [0,1]}$ is almost surely Hölder-continuous of order α . \square

Law of the iterated logarithm

Khintchine's version of the law of the iterated logarithm is a much more precise statement on the local regularity of a typical Brownian path at a fixed time $s \geq 0$. It implies in particular that almost every Brownian path is not Hölder continuous with parameter $\alpha = 1/2$. We state the result without proof:

Theorem 1.20 (Khintchine 1924). *For $s \geq 0$, the following statements hold almost surely:*

$$\limsup_{t \searrow 0} \frac{B_{s+t} - B_s}{\sqrt{2t \log \log(1/t)}} = +1, \quad \text{and} \quad \liminf_{t \searrow 0} \frac{B_{s+t} - B_s}{\sqrt{2t \log \log(1/t)}} = -1.$$

For the proof cf. e.g. Breiman, Probability, Section 12.9.

By a time inversion, the Theorem translates into a statement on the global asymptotics of Brownian paths:

Corollary 1.21. *The following statements hold almost surely:*

$$\limsup_{t \rightarrow \infty} \frac{B_t}{\sqrt{2t \log \log t}} = +1, \quad \text{and} \quad \liminf_{t \rightarrow \infty} \frac{B_t}{\sqrt{2t \log \log t}} = -1.$$

Proof. This follows by applying the Theorem above to the Brownian motion $\widehat{B}_t = t \cdot B_{1/t}$. For example, substituting $h = 1/t$, we have

$$\limsup_{t \rightarrow \infty} \frac{B_t}{\sqrt{2t \log \log(t)}} = \limsup_{h \searrow 0} \frac{h \cdot B_{1/h}}{\sqrt{2h \log \log 1/h}} = +1$$

almost surely. □

The corollary is a continuous time analogue of Kolmogorov's law of the iterated logarithm for Random Walks stating that for $S_n = \sum_{i=1}^n \eta_i$, η_i i.i.d. with $E[\eta_i] = 0$ and $\text{Var}[\eta_i] = 1$, one has

$$\limsup_{n \rightarrow \infty} \frac{S_n}{\sqrt{2n \log \log n}} = +1 \quad \text{and} \quad \liminf_{n \rightarrow \infty} \frac{S_n}{\sqrt{2n \log \log n}} = -1$$

almost surely. In fact, one way to prove Kolmogorov's LIL is to embed the Random Walk into a Brownian motion, cf. e.g. Rogers and Williams, Vol. I, Ch. 7 or Section 3.3

Passage times

We now study the set of passage times to a given level a for a one-dimensional Brownian motion $(B_t)_{t \geq 0}$. This set has interesting properties – in particular it is a random fractal. Fix $a \in \mathbb{R}$, and let

$$\Lambda_a(\omega) = \{t \geq 0 : B_t(\omega) = a\} \subseteq [0, \infty).$$

Assuming that every path is continuous, the random set $\Lambda_a(\omega)$ is *closed* for every ω . Moreover, scale invariance of Brownian motion implies a *statistical self-similarity* property for the sets of passage times: Since the rescaled process $(c^{-1/2}B_{ct})_{t \geq 0}$ has the same distribution as $(B_t)_{t \geq 0}$ for any $c > 0$, we can conclude that the set valued random variable $c \cdot \Lambda_{a/\sqrt{c}}$ has the same distribution as Λ_a . In particular, Λ_0 is a *fractal* in the sense that

$$\Lambda_0 \sim c \cdot \Lambda_0 \quad \text{for any } c > 0.$$




Figure 1.4: Brownian motion with corresponding level set Λ_0 .

Moreover, by Fubini's Theorem one easily verifies that Λ_a has almost surely Lebesgue measure zero. In fact, continuity of $t \mapsto B_t(\omega)$ for any ω implies that $(t, \omega) \mapsto B_t(\omega)$ is product measurable (Exercise). Hence $\{(t, \omega) : B_t(\omega) = a\}$ is contained in the product σ -algebra, and

$$E[\lambda(\Lambda_a)] = E \left[\int_0^\infty I_{\{a\}}(B_t) dt \right] = \int_0^\infty P[B_t = a] dt = 0.$$

Theorem 1.22 (Unbounded oscillations, recurrence).

$$P \left[\sup_{t \geq 0} B_t = +\infty \right] = P \left[\inf_{t \geq 0} B_t = -\infty \right] = 1.$$

In particular, for any $a \in \mathbb{R}$, the random set Λ_a is almost surely unbounded, i.e. Brownian motion is recurrent.

Proof. By scale invariance,

$$\sup_{t \geq 0} B_t \sim c^{-1/2} \sup_{t \geq 0} B_{ct} = c^{-1/2} \sup_{t \geq 0} B_t \quad \text{for any } c > 0.$$

Hence,

$$P \left[\sup_{t \geq 0} B_t \geq a \right] = P \left[\sup_{t \geq 0} B_t \geq a \cdot \sqrt{c} \right]$$

for any $c > 0$, and therefore $\sup B_t \in \{0, \infty\}$ almost surely. The first part of the assertion now follows since $\sup B_t$ is almost surely strictly positive. By reflection symmetry, we also obtain $\inf B_t = -\infty$ with probability one. \square

The last Theorem makes a statement on the global structure of the set Λ_a . By invariance w.r.t. time inversion this again translates into a local regularity result:

Theorem 1.23 (Fine structure of Λ_a). *The set Λ_a is almost surely a **perfect set**, i.e., any $t \in \Lambda_a$ is an accumulation point of Λ_a .*

Proof. We prove the statement for $a = 0$, the general case being left as an exercise. We proceed in three steps:

STEP 1: 0 is almost surely an accumulation point of Λ_0 : This holds by time-reversal.

Setting $\widehat{B}_t = t \cdot B_{1/t}$, we see that 0 is an accumulation point of Λ_0 if and only if for any $n \in \mathbb{N}$ there exists $t > n$ such that $\widehat{B}_t = 0$, i.e., if and only if the zero set of \widehat{B}_t is unbounded. By Theorem 1.22, this holds almost surely.

STEP 2: For any $s \geq 0$, $T_s := \min(\Lambda_a \cap [s, \infty)) = \min\{t \geq s : B_t = a\}$ is almost surely an accumulation point of Λ_a : For the proof we need the strong Markov property of Brownian motion which will be proved in the next section. By Theorem 1.22, the random variable T_s is almost surely finite. Hence, by continuity, $B_{T_s} = a$ almost surely. The strong Markov property says that the process

$$\widetilde{B}_t := B_{T_s+t} - B_{T_s}, \quad t \geq 0,$$

is again a Brownian motion starting at 0 . Therefore, almost surely, 0 is an accumulation point of the zero set of \widetilde{B}_t by Step 1. The claim follows since almost surely

$$\{t \geq 0 : \widetilde{B}_t = 0\} = \{t \geq 0 : B_{T_s+t} = B_{T_s}\} = \{t \geq T_s : B_t = a\} \subseteq \Lambda_a.$$

STEP 3: To complete the proof note that we have shown that the following properties hold with probability one:

- (1). Λ_a is closed.
- (2). $\min(\Lambda_a \cap [s, \infty))$ is an accumulation point of Λ_a for any $s \in \mathbb{Q}_+$.

Since \mathbb{Q}_+ is a dense subset of \mathbb{R}_+ , (1) and (2) imply that any $t \in \Lambda_a$ is an accumulation point of Λ_a . In fact, for any $s \in [0, t] \cap \mathbb{Q}$, there exists an accumulation point of Λ_a in $(s, t]$ by (2), and hence t is itself an accumulation point.

□

Remark. It can be shown that the set Λ_a has Hausdorff dimension $1/2$.

1.5 Strong Markov property and reflection principle

In this section we prove a strong Markov property for Brownian motion. Before, we give another motivation for our interest in an extension of the Markov property to random times.

Maximum of Brownian motion


Suppose that $(B_t)_{t \geq 0}$ is a one-dimensional continuous Brownian motion starting at 0 defined on a probability space (Ω, \mathcal{A}, P) . We would like to compute the distribution of the maximal value

$$M_s = \max_{t \in [0, s]} B_t$$

attained before a given time $s \in \mathbb{R}_+$. The idea is to proceed similarly as for Random Walks, and to reflect the Brownian path after the first passage time

$$T_a = \min\{t \geq 0 : B_t = a\}$$

to a given level $a > 0$:



It seems plausible (e.g. by the heuristic path integral representation of Wiener measure, or by a Random Walk approximation) that the reflected process $(\widehat{B}_t)_{t \geq 0}$ defined by

$$\widehat{B}_t := \begin{cases} B_t & \text{for } t \leq T_a \\ a - (B_t - a) & \text{for } t > T_a \end{cases}$$

is again a Brownian motion. At the end of this section, we will prove this reflection principle rigorously by the strong Markov property. Assuming the reflection principle is true, we can compute the distribution of M_s in the following way:

$$\begin{aligned} P[M_s \geq a] &= P[M_s \geq a, B_s \leq a] + P[M_s \geq a, B_s > a] \\ &= P[\widehat{B}_s \geq a] + P[B_s > a] \\ &= 2 \cdot P[B_s \geq a] \\ &= P[|B_s| \geq a]. \end{aligned}$$

Thus M_s has the same distribution as $|B_s|$.

Furthermore, since $M_s \geq a$ if and only if $\widehat{M}_s = \max\{\widehat{B}_t : t \in [0, s]\} \geq a$, we obtain the stronger statement

$$\begin{aligned} P[M_s \geq a, B_s \leq c] &= P[\widehat{M}_s \geq a, \widehat{B}_s \geq 2a - c] = P[\widehat{B}_s \geq 2a - c] \\ &= \frac{1}{\sqrt{2\pi s}} \int_{2a-c}^{\infty} \exp(-x^2/2s) dx \end{aligned}$$

for any $a \geq 0$ and $c \leq a$. As a consequence, we have:

Theorem 1.24 (Maxima of Brownian paths).

(1). For any $s \geq 0$, the distribution of M_s is absolutely continuous with density

$$f_{M_s}(x) = \frac{2}{\sqrt{2\pi s}} \exp(-x^2/2s) \cdot I_{(0,\infty)}(x).$$

(2). The joint distribution of M_s and B_s is absolutely continuous with density

$$f_{M_s, B_s}(x, y) = 2 \frac{2x - y}{\sqrt{2\pi s^3}} \exp\left(-\frac{(2x - y)^2}{2s}\right) I_{(0,\infty)}(x) I_{(-\infty, x)}(y).$$

Proof. (1) holds since $M_s \sim |B_s|$. For the proof of (2) we assume w.l.o.g. $s = 1$. The general case can be reduced to this case by the scale invariance of Brownian motion (Exercise). For $a \geq 0$ and $c \leq a$ let

$$G(a, c) := P[M_1 \geq a, B_1 \leq c].$$

By the reflection principle,

$$G(a, c) = P[B_1 \geq 2a - c] = 1 - \Phi(2a - c),$$

where Φ denotes the standard normal distribution function. Since $\lim_{a \rightarrow \infty} G(a, c) = 0$ and $\lim_{c \rightarrow -\infty} G(a, c) = 0$, we obtain

$$\begin{aligned} P[M_1 \geq a, B_1 \leq c] = G(a, c) &= - \int_{x=a}^{\infty} \int_{y=-\infty}^c \frac{\partial^2 G}{\partial x \partial y}(x, y) dy dx \\ &= \int_{x=a}^{\infty} \int_{y=-\infty}^c 2 \cdot \frac{2x-y}{\sqrt{2\pi}} \cdot \exp\left(-\frac{(2x-y)^2}{2}\right) dy dx. \end{aligned}$$

This implies the claim for $s = 1$, since $M_1 \geq 0$ and $B_1 \leq M_1$ by definition of M_1 . \square

The Theorem enables us to compute the distributions of the first passage times T_a . In fact, for $a > 0$ and $s \in [0, \infty)$ we obtain

$$\begin{aligned} P[T_a \leq s] = P[M_s \geq a] &= 2 \cdot P[B_s \geq a] = 2 \cdot P[B_1 \geq a/\sqrt{s}] \\ &= \sqrt{\frac{2}{\pi}} \int_{a/\sqrt{s}}^{\infty} e^{-x^2/2} dx. \end{aligned} \tag{1.5.1}$$

Corollary 1.25 (Distribution of T_a). For any $a \in \mathbb{R} \setminus \{0\}$, the distribution of T_a is absolutely continuous with density

$$f_{T_a}(s) = \frac{|a|}{\sqrt{2\pi s^3}} \cdot e^{-a^2/2s}.$$

Proof. For $a > 0$, we obtain

$$f_{T_a}(s) = F'_{T_a}(s) = \frac{a}{\sqrt{2\pi s^3}} e^{-a^2/2s}$$

by (1.5.1). For $a < 0$ the assertion holds since $T_a \sim T_{-a}$ by reflection symmetry of Brownian motion. \square

Next, we prove a strong Markov property for Brownian motion. Below we will then complete the proof of the reflection principle and the statements above by applying the strong Markov property to the passage time T_a .

Strong Markov property for Brownian motion

Suppose again that $(B_t)_{t \geq 0}$ is a d -dimensional continuous Brownian motion starting at 0 on a probability space (Ω, \mathcal{A}, P) , and let

$$\mathcal{F}_t^B = \sigma(B_s : 0 \leq s \leq t), \quad t \geq 0,$$

denote the σ -algebras generated by the process up to time t .

Definition. A random variable $T : \Omega \rightarrow [0, \infty]$ is called an (\mathcal{F}_t^B) -**stopping time** if and only if

$$\{T \leq t\} \in \mathcal{F}_t^B \quad \text{for any } t \geq 0.$$

Example. Clearly, for any $a \in \mathbb{R}$, the first passage time

$$T_a = \min\{t \geq 0 : B_t = a\}$$

to a level a is an (\mathcal{F}_t^B) -stopping time.

The σ -algebra \mathcal{F}_T^B describing the information about the process up to a stopping time T is defined by

$$\mathcal{F}_T^B = \{A \in \mathcal{A} : A \cap \{T \leq t\} \in \mathcal{F}_t^B \text{ for any } t \geq 0\}.$$

Note that for (\mathcal{F}_t^B) stopping times S and T with $S \leq T$ we have $\mathcal{F}_S^B \subseteq \mathcal{F}_T^B$, since for $t \geq 0$

$$A \cap \{S \leq t\} \in \mathcal{F}_t^B \quad \implies \quad A \cap \{T \leq t\} = A \cap \{S \leq t\} \cap \{T \leq t\} \in \mathcal{F}_t^B.$$

For any constant $s \in \mathbb{R}_+$, the process $(B_{s+t} - B_s)_{t \geq 0}$ is a Brownian motion independent of \mathcal{F}_s^B .

A corresponding statement holds for stopping times:

Theorem 1.26 (Strong Markov property). *Suppose that T is an almost surely finite (\mathcal{F}_t^B) stopping time. Then the process $(\tilde{B}_t)_{t \geq 0}$ defined by*

$$\tilde{B}_t = B_{T+t} - B_T \quad \text{if } T < \infty, \quad 0 \quad \text{otherwise,}$$

is a Brownian motion independent of \mathcal{F}_T^B .

Proof. We first assume that T takes values only in $C \cup \{\infty\}$ where C is a countable subset of $[0, \infty)$. Then for $A \in \mathcal{F}_T^B$ and $s \in C$, we have $A \cap \{T = s\} \in \mathcal{F}_s^B$ and $\tilde{B}_t = B_{t+s} - B_s$ on $A \cap \{T = s\}$. By the Markov property, $(B_{t+s} - B_s)_{t \geq 0}$ is a Brownian motion independent of \mathcal{F}_s^B . Hence for any measurable subset Γ of $C([0, \infty], \mathbb{R}^d)$, we have

$$\begin{aligned} P[\{(\tilde{B}_t)_{t \geq 0} \in \Gamma\} \cap A] &= \sum_{s \in C} P[\{(B_{t+s} - B_s)_{t \geq 0} \in \Gamma\} \cap A \cap \{T = s\}] \\ &= \sum_{s \in C} \mu_0[\Gamma] \cdot P[A \cap \{T = s\}] = \mu_0[\Gamma] \cdot P[A] \end{aligned}$$

where μ_0 denotes the distribution of Brownian motion starting at 0. This proves the assertion for discrete stopping times.

For an arbitrary (\mathcal{F}_t^B) stopping time T that is almost surely finite and $n \in \mathbb{N}$, we set $T_n = \frac{1}{n} \lceil nT \rceil$, i.e.,

$$T_n = \frac{k}{n} \quad \text{on} \quad \left\{ \frac{k-1}{n} < T \leq \frac{k}{n} \right\} \quad \text{for any } k \in \mathbb{N}.$$

Since the event $\{T_n = k/n\}$ is $\mathcal{F}_{k/n}^B$ -measurable for any $k \in \mathbb{N}$, T_n is a discrete (\mathcal{F}_t^B) stopping time. Therefore, $(B_{T_n+t} - B_{T_n})_{t \geq 0}$ is a Brownian motion that is independent of $\mathcal{F}_{T_n}^B$, and hence of the smaller σ -algebra \mathcal{F}_T^B . As $n \rightarrow \infty$, $T_n \rightarrow T$, and thus, by continuity,

$$\tilde{B}_t = B_{T+t} - B_T = \lim_{n \rightarrow \infty} (B_{T_n+t} - B_{T_n}).$$

Now it is easy to verify that $(\tilde{B}_t)_{t \geq 0}$ is again a Brownian motion that is independent of \mathcal{F}_T^B . \square

A rigorous reflection principle

We now apply the strong Markov property to prove a reflection principle for Brownian motion. Consider a one-dimensional continuous Brownian motion $(B_t)_{t \geq 0}$ starting at 0. For $a \in \mathbb{R}$ let

$$\begin{aligned} T_a &= \min\{t \geq 0 : B_t = a\} && \text{(first passage time),} \\ B_t^{T_a} &= B_{\min\{t, T_a\}} && \text{(process stopped at } T_a), \quad \text{and} \\ \tilde{B}_t &= B_{T_a+t} - B_{T_a} && \text{(process after } T_a). \end{aligned}$$


Theorem 1.27 (Reflection principle). *The joint distributions of the following random variables with values in $\mathbb{R}_+ \times C([0, \infty)) \times C([0, \infty))$ agree:*

$$(T_a, (B_t^{T_a})_{t \geq 0}, (\tilde{B}_t)_{t \geq 0}) \sim (T_a, (B_t^{T_a})_{t \geq 0}, (-\tilde{B}_t)_{t \geq 0})$$

Proof. By the strong Markov property, the process \tilde{B} is a Brownian motion starting at 0 independent of \mathcal{F}_{T_a} , and hence of T_a and $B^{T_a} = (B_t^{T_a})_{t \geq 0}$. Therefore,

$$P \circ (T_a, B^{T_a}, \tilde{B})^{-1} = P \circ (T_a, B^{T_a})^{-1} \otimes \mu_0 = P \circ (T_a, B^{T_a}, -\tilde{B})^{-1}.$$

\square



As a consequence of the theorem, we can complete the argument given at the beginning of this section: The "shadow path" \widehat{B}_t of a Brownian path B_t with reflection when reaching the level a is given by

$$\widehat{B}_t = \begin{cases} B_t^{T_a} & \text{for } t \leq T_a \\ a - \widetilde{B}_{t-T_a} & \text{for } t > T_a \end{cases},$$

whereas

$$B_t = \begin{cases} B_t^{T_a} & \text{for } t \leq T_a \\ a + \widetilde{B}_{t-T_a} & \text{for } t > T_a \end{cases}.$$

By the Theorem 1.27, $(\widehat{B}_t)_{t \geq 0}$ has the same distribution as $(B_t)_{t \geq 0}$. Therefore, and since $\max_{t \in [0, s]} B_t \geq a$ if and only if $\max_{t \in [0, s]} \widehat{B}_t \geq a$, we obtain for $a \geq c$:

$$\begin{aligned} P \left[\max_{t \in [0, s]} B_t \geq a, B_s \leq c \right] &= P \left[\max_{t \in [0, s]} \widehat{B}_t \geq a, \widehat{B}_s \geq 2a - c \right] \\ &= P \left[\widehat{B}_s \geq 2a - c \right] \\ &= \frac{1}{\sqrt{2\pi s}} \int_{2a-c}^{\infty} e^{-x^2/2s} dx. \end{aligned}$$

Part II

Introduction to Stochastic Analysis

Chapter 2

Martingales in discrete time

Classical analysis starts with studying convergence of sequences of real numbers. Similarly, stochastic analysis relies on basic statements about sequences of real-valued random variables. Any such sequence can be decomposed uniquely into a martingale, i.e., a real-valued stochastic process that is “constant on average”, and a predictable part. Therefore, estimates and convergence theorems for martingales are crucial in stochastic analysis.

2.1 Definitions and examples

We fix a probability space (Ω, \mathcal{A}, P) . Moreover, we assume that we are given an increasing sequence \mathcal{F}_n ($n = 0, 1, 2, \dots$) of sub- σ -algebras of \mathcal{A} . Intuitively, we often think of \mathcal{F}_n as describing the information available to us at time n . Formally, we define:

Definition (Filtration, adapted process). (1). A *filtration* on (Ω, \mathcal{A}) is an increasing sequence

$$\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \mathcal{F}_2 \subseteq \dots$$

of σ -algebras $\mathcal{F}_n \subseteq \mathcal{A}$.

(2). A stochastic process $(X_n)_{n \geq 0}$ is **adapted** to a filtration $(\mathcal{F}_n)_{n \geq 0}$ iff each X_n is \mathcal{F}_n -measurable.

Example. (1). The **canonical filtration** (\mathcal{F}_n^X) generated by a stochastic process (X_n) is given by

$$\mathcal{F}_n^X = \sigma(X_0, X_1, \dots, X_n).$$

If the filtration is not specified explicitly, we will usually consider the canonical filtration.

(2). Alternatively, filtrations containing additional information are of interest, for example the filtration

$$\mathcal{F}_n = \sigma(Z, X_0, X_1, \dots, X_n)$$

generated by the process (X_n) and an additional random variable Z , or the filtration

$$\mathcal{F}_n = \sigma(X_0, Y_0, X_1, Y_1, \dots, X_n, Y_n)$$

generated by the process (X_n) and a further process (Y_n) .

Clearly, the process (X_n) is adapted to any of these filtrations. In general, (X_n) is adapted to a filtration (\mathcal{F}_n) if and only if $\mathcal{F}_n^X \subseteq \mathcal{F}_n$ for any $n \geq 0$.

Martingales and supermartingales

We can now formalize the notion of a real-valued stochastic process that is constant (respectively decreasing or increasing) on average:

Definition (Martingale, supermartingale, submartingale). (1). A sequence of real-valued random variables $M_n : \Omega \rightarrow \mathbb{R}$ ($n = 0, 1, \dots$) on the probability space (Ω, \mathcal{A}, P) is called a **martingale w.r.t. the filtration** (\mathcal{F}_n) if and only if

- (a) (M_n) is adapted w.r.t. (\mathcal{F}_n) ,
- (b) M_n is integrable for any $n \geq 0$, and
- (c) $E[M_n | \mathcal{F}_{n-1}] = M_{n-1}$ for any $n \in \mathbb{N}$.

(2). Similarly, (M_n) is called a **supermartingale** (resp. a **submartingale**) w.r.t. (\mathcal{F}_n) if and only if (a) holds, the positive part M_n^+ (resp. the negative part M_n^-) is integrable for any $n \geq 0$, and (c) holds with “=” replaced by “ \leq ”, “ \geq ” respectively.

Condition (c) in the martingale definition can equivalently be written as

$$(c') \quad E[M_{n+1} - M_n \mid \mathcal{F}_n] = 0 \quad \text{for any } n \in \mathbb{Z}_+,$$

and correspondingly with “=” replaced by “ \leq ” or “ \geq ” for super- or submartingales.

Intuitively, a martingale is a “fair game”, i.e., M_{n-1} is the best prediction (w.r.t. the mean square error) for the next value M_n given the information up to time $n - 1$. A supermartingale is “decreasing on average”, a submartingale is “increasing on average”, and a martingale is both “decreasing” and “increasing”, i.e., “constant on average”. In particular, by induction on n , a martingale satisfies

$$E[M_n] = E[M_0] \quad \text{for any } n \geq 0.$$

Similarly, for a supermartingale, the expectation values $E[M_n]$ are decreasing. More generally, we have:

Lemma 2.1. *If (M_n) is a martingale (respectively a supermartingale) w.r.t. a filtration (\mathcal{F}_n) then*

$$E[M_{n+k} \mid \mathcal{F}_n] \stackrel{(\leq)}{=} M_n \quad P\text{-almost surely for any } n, k \geq 0.$$

Proof. By induction on k : The assertion holds for $k = 0$, since M_n is \mathcal{F}_n -measurable. Moreover, the assertion for $k - 1$ implies

$$\begin{aligned} E[M_{n+k} \mid \mathcal{F}_n] &= E[E[M_{n+k} \mid \mathcal{F}_{n+k-1}] \mid \mathcal{F}_n] \\ &= E[M_{n+k-1} \mid \mathcal{F}_n] = M_n \quad P\text{-a.s.} \end{aligned}$$

by the tower property for conditional expectations. □

Remark (Supermartingale Convergence Theorem). A key fact in analysis is that any lower bounded decreasing sequence of real numbers converges to its infimum. The counterpart of this result in stochastic analysis is the Supermartingale Convergence Theorem: Any lower bounded supermartingale converges almost surely, cf. Theorem 4.5 below.

Some fundamental examples

a) Sums of independent random variables

A Random Walk

$$S_n = \sum_{i=1}^n \eta_i, \quad n = 0, 1, 2, \dots,$$

with independent increments $\eta_i \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$ is a martingale w.r.t. to the filtration

$$\mathcal{F}_n = \sigma(\eta_1, \dots, \eta_n) = \sigma(S_0, S_1, \dots, S_n)$$

if and only if the increments η_i are centered random variables. In fact, for any $n \in \mathbb{N}$,

$$E[S_n - S_{n-1} | \mathcal{F}_{n-1}] = E[\eta_n | \mathcal{F}_{n-1}] = E[\eta_n]$$

by independence of the increments. Correspondingly, (S_n) is an (\mathcal{F}_n) supermartingale if and only if $E[\eta_i] \leq 0$ for any $i \in \mathbb{N}$.

b) Products of independent non-negative random variables

A stochastic process

$$M_n = \prod_{i=1}^n Y_i, \quad n = 0, 1, 2, \dots,$$

with independent non-negative factors $Y_i \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$ is a martingale respectively a supermartingale w.r.t. the filtration

$$\mathcal{F}_n = \sigma(Y_1, \dots, Y_n)$$

if and only if $E[Y_i] = 1$ for any $i \in \mathbb{N}$, or $E[Y_i] \leq 1$ for any $i \in \mathbb{N}$ respectively. In fact, as M_n is \mathcal{F}_n -measurable and Y_{n+1} is independent of \mathcal{F}_n , we have

$$E[M_{n+1} | \mathcal{F}_n] = E[M_n \cdot Y_{n+1} | \mathcal{F}_n] = M_n \cdot E[Y_{n+1}] \quad \text{for any } n \geq 0.$$

Martingales and supermartingales of this type occur naturally in stochastic growth models.

Example (Exponential martingales). Consider a Random Walk $S_n = \sum_{i=1}^n \eta_i$ with i.i.d. increments η_i , and let

$$Z(\lambda) = E[\exp(\lambda\eta_i)], \quad \lambda \in \mathbb{R},$$

denote the moment generating function of the increments. Then for any $\lambda \in \mathbb{R}$ with $Z(\lambda) < \infty$, the process

$$M_n^\lambda := e^{\lambda S_n} / Z(\lambda)^n = \prod_{i=1}^n (e^{\lambda\eta_i} / Z(\lambda))$$

is a martingale. This martingale can be used to prove exponential bounds for Random Walks, cf. e.g. Chernov's theorem ["Einführung in die Wahrscheinlichkeitstheorie", Theorem 8.3].

Example (CRR model of stock market). In the Cox-Ross-Rubinstein binomial model of mathematical finance, the price of an asset is changing during each period either by a factor $1 + a$ or by a factor $1 + b$ with $a, b \in (-1, \infty)$ such that $a < b$. We can model the price evolution in a fixed number N of periods by a stochastic process

$$S_n = S_0 \cdot \prod_{i=1}^n X_i, \quad n = 0, 1, 2, \dots, N,$$

defined on $\Omega = \{1 + a, 1 + b\}^N$, where the initial price S_0 is a given constant, and $X_i(\omega) = \omega_i$. Taking into account a constant interest rate $r > 0$, the discounted stock price after n periods is

$$\tilde{S}_n = S_n / (1 + r)^n = S_0 \cdot \prod_{i=1}^n \frac{X_i}{1 + r}.$$

A probability measure P on Ω is called a **martingale measure** if the discounted stock price is a martingale w.r.t. P and the filtration $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$. Martingale measures are important for option pricing under no arbitrage assumptions, cf. Section 2.3 below. For $1 \leq n \leq N$,

$$E[\tilde{S}_n | \mathcal{F}_{n-1}] = E \left[\tilde{S}_{n-1} \cdot \frac{X_n}{1 + r} \middle| \mathcal{F}_{n-1} \right] = \tilde{S}_{n-1} \cdot \frac{E[X_n | \mathcal{F}_{n-1}]}{1 + r}.$$

Hence (\tilde{S}_n) is an (\mathcal{F}_n) martingale w.r.t. P if and only if

$$E[X_n | \mathcal{F}_{n-1}] = 1 + r \quad \text{for any } 1 \leq n \leq N. \quad (2.1.1)$$

On the other hand, since in the CRR model X_n only takes the values $1 + a$ and $1 + b$, we have

$$\begin{aligned} E[X_n | \mathcal{F}_{n-1}] &= (1 + a) \cdot P[X_n = 1 + a | \mathcal{F}_{n-1}] + (1 + b) \cdot P[X_n = 1 + b | \mathcal{F}_{n-1}] \\ &= 1 + a + (b - a) \cdot P[X_n = 1 + b | \mathcal{F}_{n-1}]. \end{aligned}$$

Therefore, by (2.1.1), (\tilde{S}_n) is a martingale if and only if

$$P[X_n = 1 + b | \mathcal{F}_{n-1}] = \frac{r - a}{b - a} \quad \text{for any } n = 1, \dots, N,$$

i.e., if and only if the growth factors X_1, \dots, X_N are independent with

$$P[X_n = 1 + b] = \frac{r - a}{b - a} \quad \text{and} \quad P[X_n = 1 + a] = \frac{b - r}{b - a}. \quad (2.1.2)$$

Hence for $r \notin [a, b]$, a martingale measure does not exist, and for $r \in [a, b]$, the product measure P on Ω satisfying (2.1.2) is the unique martingale measure. Intuitively this is plausible: If $r < a$ or $r > b$ respectively, then the stock price is always growing more or less than the discount factor $(1 + r)^n$, so the discounted stock price can not be a martingale. If, on the other hand, $a < r < b$, then (\tilde{S}_n) is a martingale provided the growth factors are independent with

$$\frac{P[X_n = 1 + b]}{P[X_n = 1 + a]} = \frac{(1 + r) - (1 + a)}{(1 + b) - (1 + r)}.$$

We remark, however, that uniqueness of the martingale measure only follows from (2.1.1) since we have assumed that each X_n takes only two possible values (binomial model). In a corresponding trinomial model there are infinitely many martingale measures!

c) Successive prediction values

Let F be an integrable random variable, and let (\mathcal{F}_n) be a filtration on a probability space (Ω, \mathcal{A}, P) . Then the process

$$M_n := E[F | \mathcal{F}_n], \quad n = 0, 1, 2, \dots,$$

of successive prediction values for F based on the information up to time n is a martingale. Indeed, by the tower property for conditional expectations, we have

$$E[M_n | \mathcal{F}_{n-1}] = E[E[F | \mathcal{F}_n] | \mathcal{F}_{n-1}] = E[F | \mathcal{F}_{n-1}] = M_{n-1}$$

almost surely for any $n \in \mathbb{N}$.

Remark (Representing martingales as successive prediction values). The class of martingales that have a representation as successive prediction values almost contains general martingales. In fact, for an arbitrary (\mathcal{F}_n) martingale (M_n) and any finite integer $m \geq 0$, the representation

$$M_n = E[M_m | \mathcal{F}_n]$$

holds for any $n = 0, 1, \dots, m$. Moreover, the L^1 Martingale Convergence Theorem implies that under a uniform integrability assumption, the limit $M_\infty = \lim_{m \rightarrow \infty} M_m$ exists in \mathcal{L}^1 , and the representation

$$M_n = E[M_\infty | \mathcal{F}_n]$$

holds for any $n \geq 0$, see Section 4.3 below .

d) Functions of martingales

By Jensen's inequality for conditional expectations, convex functions of martingales are submartingales, and concave functions of martingales are supermartingales:

Theorem 2.2 (Convex functions of martingales). *Suppose that $(M_n)_{n \geq 0}$ is an (\mathcal{F}_n) martingale, and $u : \mathbb{R} \rightarrow \mathbb{R}$ is a convex function that is bounded from below. Then $(u(M_n))$ is an (\mathcal{F}_n) submartingale.*

Proof. Since u is lower bounded, $u(M_n)^-$ is integrable for any n . Jensen's inequality for conditional expectations now implies

$$E[u(M_{n+1}) | \mathcal{F}_n] \geq u(E[M_{n+1} | \mathcal{F}_n]) = u(M_n)$$

almost surely for any $n \geq 0$. □

Example. If (M_n) is a martingale then $(|M_n|^p)$ is a submartingale for any $p \geq 1$.

e) Functions of Markov chains

Let $p(x, dy)$ be a transition kernel on a measurable space (S, \mathcal{B}) .

Definition (Markov chain, superharmonic function). (1). A discrete time stochastic process $(X_n)_{n \geq 0}$ with state space (S, \mathcal{B}) defined on the probability space (Ω, \mathcal{A}, P) is called a **(time-homogeneous) Markov chain with transition kernel p w.r.t. the filtration (\mathcal{F}_n)** , if and only if

(a) (X_n) is (\mathcal{F}_n) adapted, and

(b) $P[X_{n+1} \in B | \mathcal{F}_n] = p(X_n, B)$ P -almost surely for any $B \in \mathcal{B}$ and $n \geq 0$.

(2). A measurable function $h : S \rightarrow \mathbb{R}$ is called **superharmonic** (resp. **subharmonic**) w.r.t. p if and only if the integrals

$$(ph)(x) := \int p(x, dy)h(y), \quad x \in S,$$

exist, and

$$(ph)(x) \leq h(x) \quad (\text{respectively } (ph)(x) \geq h(x))$$

holds for any $x \in S$.

The function h is called **harmonic** iff it is both super- and subharmonic, i.e., iff

$$(ph)(x) = h(x) \quad \text{for any } x \in S.$$

By the tower property for conditional expectations, any (\mathcal{F}_n) Markov chain is also a Markov chain w.r.t. the canonical filtration generated by the process.

Example (Classical Random Walk on \mathbb{Z}^d). The standard Random Walk $(X_n)_{n \geq 0}$ on \mathbb{Z}^d is a Markov chain w.r.t. the filtration $\mathcal{F}_n^X = \sigma(X_0, \dots, X_n)$ with transition probabilities $p(x, x + e) = 1/2d$ for any unit vector $e \in \mathbb{Z}^d$. The coordinate processes $(X_n^i)_{n \geq 0}$, $i = 1, \dots, d$, are Markov chains w.r.t. the same filtration with transition probabilities

$$\bar{p}(x, x + 1) = \bar{p}(x, x - 1) = \frac{1}{2d}, \quad \bar{p}(x, x) = \frac{2d - 2}{2d}.$$

A function $h : \mathbb{Z}^d \rightarrow \mathbb{R}$ is superharmonic w.r.t. p if and only if

$$\Delta_{\mathbb{Z}^d} h(x) = \sum_{i=1}^d (h(x + e_i) - 2h(x) + h(x - e_i)) = 2d((ph)(x) - h(x)) \leq 0$$

for any $x \in \mathbb{Z}^d$.

A function $h : \mathbb{Z} \rightarrow \mathbb{R}$ is harmonic w.r.t. \bar{p} if and only if $h(x) = ax + b$ with $a, b \in \mathbb{R}$, and h is superharmonic if and only if it is concave.

It is easy to verify that (super-)harmonic functions of Markov chains are (super-)martingales:

Theorem 2.3 (Superharmonic functions of Markov chains are supermartingales).

Suppose that (X_n) is an (\mathcal{F}_n) Markov chain. Then the real-valued process

$$M_n := h(X_n), \quad n = 0, 1, 2, \dots,$$

is a martingale (resp. a supermartingale) w.r.t. (\mathcal{F}_n) for every harmonic (resp. superharmonic) function $h : S \rightarrow \mathbb{R}$ such that $h(X_n)$ (resp. $h(X_n)^+$) is integrable for all n .

Proof. Clearly, (M_n) is again (\mathcal{F}_n) adapted. Moreover,

$$E[M_{n+1} | \mathcal{F}_n] = E[h(X_{n+1}) | \mathcal{F}_n] = (ph)(X_n) \quad P\text{-a.s.}$$

The assertion now follows immediately from the definitions. \square

Below, we will show how to construct more general martingales from Markov chains, cf. Theorem 2.5. At first, however, we consider a simple example that demonstrates the usefulness of martingale methods in analyzing Markov chains:

Example (Wright model for evolution). In the Wright model for a population of N individuals (replicas) with a finite number of possible types, each individual in generation $n + 1$ inherits a type from a randomly chosen predecessor in the n th generation.

The number X_n of individuals of a given type in generation n is a Markov chain with state space $S = \{0, 1, \dots, N\}$ and transition kernel

$$p(k, \bullet) = \text{Bin}(N, k/N).$$




Figure 2.1: Transition function of (X_n) .

Moreover, as the average of this binomial distribution is k , the function $h(x) = x$ is harmonic, and the expected number of individuals in generation $n+1$ given X_0, \dots, X_n is

$$E[X_{n+1} | X_0, \dots, X_n] = X_n.$$

Hence, the process (X_n) is a bounded martingale. The Martingale Convergence Theorem now implies that the limit $X_\infty = \lim X_n$ exists almost surely, cf. Section 4.2 below. Since X_n takes discrete values, we can conclude that $X_n = X_\infty$ eventually with probability one. In particular, X_∞ is almost surely an absorbing state. Hence

$$P[X_n = 0 \text{ or } X_n = N \text{ eventually}] = 1. \quad (2.1.3)$$

In order to compute the probabilities of the events “ $X_n = 0$ eventually” and “ $X_n = N$ eventually” we can apply the Optional Stopping Theorem for martingales, cf. Section 2.3 below. Let

$$T := \min\{n \geq 0 : X_n = 0 \text{ or } X_n = N\}, \quad \min \emptyset := \infty,$$

denote the first hitting time of the absorbing states. If the initial number X_0 of individuals of the given type is k , then by the Optional Stopping Theorem,

$$E[X_T] = E[X_0] = k.$$

Hence by (2.1.3) we obtain

$$\begin{aligned} P[X_n = N \text{ eventually}] &= P[X_T = N] = \frac{1}{N}E[X_T] = \frac{k}{N}, & \text{and} \\ P[X_n = 0 \text{ eventually}] &= 1 - \frac{k}{N} = \frac{N - k}{N}. \end{aligned}$$

Hence eventually all individuals have the same type, and a given type occurs eventually with probability determined by its initial relative frequency in the population.

2.2 Doob Decomposition and Martingale Problem

We will show now that any adapted sequence of real-valued random variables can be decomposed into a martingale and a predictable process. In particular, the variance process of a martingale (M_n) is the predictable part in the corresponding Doob decomposition of the process (M_n^2) . The Doob decomposition for functions of Markov chains implies the martingale problem characterization of Markov chains.

Doob Decomposition

Let (Ω, \mathcal{A}, P) be a probability space and $(\mathcal{F}_n)_{n \geq 0}$ a filtration on (Ω, \mathcal{A}) .

Definition (Predictable process). A stochastic process $(A_n)_{n \geq 0}$ is called *predictable w.r.t.* (\mathcal{F}_n) if and only if A_0 is constant and A_n is measurable w.r.t. \mathcal{F}_{n-1} for any $n \in \mathbb{N}$.

Intuitively, the value $A_n(\omega)$ of a predictable process can be predicted by the information available at time $n - 1$.

Theorem 2.4 (Doob decomposition). Every (\mathcal{F}_n) adapted sequence of integrable random variables Y_n ($n \geq 0$) has a unique decomposition (up to modification on null sets)

$$Y_n = M_n + A_n \tag{2.2.1}$$

into an (\mathcal{F}_n) martingale (M_n) and a predictable process (A_n) such that $A_0 = 0$. Explicitly, the decomposition is given by

$$A_n = \sum_{k=1}^n E[Y_k - Y_{k-1} | \mathcal{F}_{k-1}], \quad \text{and} \quad M_n = Y_n - A_n. \quad (2.2.2)$$

Remark. (1). The increments $E[Y_k - Y_{k-1} | \mathcal{F}_{k-1}]$ of the process (A_n) are the predicted increments of (Y_n) given the previous information.

(2). The process (Y_n) is a supermartingale (resp. a submartingale) if and only if the predictable part (A_n) is decreasing (resp. increasing).

Proof of Theorem 2.4. Uniqueness: For any decomposition as in (2.2.1) we have

$$Y_k - Y_{k-1} = M_k - M_{k-1} + A_k - A_{k-1} \quad \text{for any } k \in \mathbb{N}.$$

If (M_n) is a martingale and (A_n) is predictable then

$$E[Y_k - Y_{k-1} | \mathcal{F}_{k-1}] = E[A_k - A_{k-1} | \mathcal{F}_{k-1}] = A_k - A_{k-1} \quad P\text{-a.s.}$$

This implies that (2.2.2) holds almost surely if $A_0 = 0$.

Existence: Conversely, if (A_n) and (M_n) are defined by (2.2.2) then (A_n) is predictable with $A_0 = 0$ and (M_n) is a martingale, since

$$E[M_k - M_{k-1} | \mathcal{F}_{k-1}] = 0 \quad P\text{-a.s. for any } k \in \mathbb{N}.$$

□

Conditional Variance Process

Consider a martingale (M_n) such that M_n is square integrable for any $n \geq 0$. Then, by Jensen's inequality, (M_n^2) is a submartingale and can again be decomposed into a martingale (\widetilde{M}_n) and a predictable process $\langle M \rangle_n$ such that $\langle M \rangle_0 = 0$:

$$M_n^2 = \widetilde{M}_n + \langle M \rangle_n \quad \text{for any } n \geq 0.$$

The increments of the predictable process are given by

$$\begin{aligned} \langle M \rangle_k - \langle M \rangle_{k-1} &= E[M_k^2 - M_{k-1}^2 \mid \mathcal{F}_{k-1}] \\ &= E[(M_k - M_{k-1})^2 \mid \mathcal{F}_{k-1}] + 2 \cdot E[M_{k-1} \cdot (M_k - M_{k-1}) \mid \mathcal{F}_{k-1}] \\ &= \text{Var}[M_k - M_{k-1} \mid \mathcal{F}_{k-1}] \quad \text{for any } k \in \mathbb{N}. \end{aligned}$$

Here we have used in the last step that $E[M_k - M_{k-1} \mid \mathcal{F}_{k-1}]$ vanishes since (M_n) is a martingale.

Definition (Conditional variance process). *The predictable process*

$$\langle M \rangle_n := \sum_{k=1}^n \text{Var}[M_k - M_{k-1} \mid \mathcal{F}_{k-1}], \quad n \geq 0,$$

is called the conditional variance process of the square integrable martingale (M_n) .

Example (Random Walks). If $M_n = \sum_{i=1}^n \eta_i$ is a sum of independent centered random variables η_i and $\mathcal{F}_n = \sigma(\eta_1, \dots, \eta_n)$ then the conditional variance process is given by $\langle M \rangle_n = \sum_{i=1}^n \text{Var}[\eta_i]$.

The conditional variance process is crucial for generalizations of classical limit theorems such as the Law of Large Numbers or the Central Limit Theorem from sums of independent random variables to martingales. A direct consequence of the fact that $M_n^2 - \langle M \rangle_n$ is a martingale is that

$$E[M_n^2] = E[M_0^2] + E[\langle M \rangle_n] \quad \text{for any } n \geq 0.$$

This can often be used to derive L^2 -estimates for martingales.

Example (Discretizations of stochastic differential equations). Consider an ordinary differential equation

$$\frac{dX_t}{dt} = b(X_t), \quad t \geq 0, \quad (2.2.3)$$

where $b : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a given vector field. In order to take into account unpredictable effects on a system, one is frequently interested in studying random perturbations of the dynamics (2.2.3) of type

$$dX_t = b(X_t) dt + \text{“noise”} \quad (2.2.4)$$

with a random noise term. The solution $(X_t)_{t \geq 0}$ of such a stochastic differential equation (SDE) is a stochastic process in continuous time defined on a probability space (Ω, \mathcal{A}, P) where also the random variables describing the noise effects are defined. The vector field b is called the (deterministic) “drift”. We will make sense of general SDE later, but we can already consider time discretizations.

For simplicity let us assume $d = 1$. Let $b, \sigma : \mathbb{R} \rightarrow \mathbb{R}$ be continuous functions, and let $(\eta_i)_{i \in \mathbb{N}}$ be a sequence of i.i.d. random variables $\eta_i \in \mathcal{L}^2(\Omega, \mathcal{A}, P)$ describing the noise effects. We assume

$$E[\eta_i] = 0 \quad \text{and} \quad \text{Var}[\eta_i] = 1 \quad \text{for any } i \in \mathbb{N}.$$

Here, the values 0 and 1 are just a convenient normalization, but it is an important assumption that the random variables are independent with finite variances. Given an initial value $x_0 \in \mathbb{R}$ and a fine discretization step size $h > 0$, we now define a stochastic process $(X_n^{(h)})$ in discrete time by $X_0^{(h)} = x_0$, and

$$X_{k+1}^{(h)} - X_k^{(h)} = b(X_k^{(h)}) \cdot h + \sigma(X_k^{(h)}) \sqrt{h} \eta_{k+1}, \quad \text{for } k = 0, 1, 2, \dots \quad (2.2.5)$$

One should think of $X_k^{(h)}$ as an approximation for the value of the process (X_t) at time $t = k \cdot h$. The equation (2.2.5) can be rewritten as

$$X_n^{(h)} = x_0 + \sum_{k=0}^{n-1} b(X_k^{(h)}) \cdot h + \sum_{k=0}^{n-1} \sigma(X_k^{(h)}) \cdot \sqrt{h} \cdot \eta_{k+1}. \quad (2.2.6)$$

To understand the scaling factors h and \sqrt{h} we note first that if $\sigma \equiv 0$ then (2.2.5) respectively (2.2.6) is the Euler discretization of the ordinary differential equation (2.2.3). Furthermore, if $b \equiv 0$ and $\sigma \equiv 1$, then the *diffusive scaling* by a factor \sqrt{h} in the second term ensures that the continuous time process $X_{\lfloor t/h \rfloor}^{(h)}, t \in [0, \infty)$, converges in distribution as $h \searrow 0$. Indeed, the functional central limit theorem (Donsker’s invariance

principle) states that the limit process in this case is a Brownian motion $(B_t)_{t \in [0, \infty)}$. In general, (2.2.6) is an Euler discretization of a stochastic differential equation of type

$$dX_t = b(X_t) dt + \sigma(X_t) dB_t$$

where $(B_t)_{t \geq 0}$ is a Brownian motion. Let $\mathcal{F}_n = \sigma(\eta_1, \dots, \eta_n)$ denote the filtration generated by the random variables η_i . The following exercise summarizes basic properties of the process $X^{(h)}$ in the case of normally distributed increments.

Exercise. Suppose that the random variables η_i are standard normally distributed.

- (1). Prove that the process $X^{(h)}$ is a time-homogeneous (\mathcal{F}_n) Markov chain with transition kernel

$$p(x, \bullet) = N(x + b(x)h, \sigma(x)^2 h)[\bullet].$$

- (2). Show that the Doob decomposition $X^{(h)} = M^{(h)} + A^{(h)}$ is given by

$$A_n^{(h)} = \sum_{k=0}^{n-1} b(X_k^{(h)}) \cdot h, \quad M_n^{(h)} = x_0 + \sum_{k=0}^{n-1} \sigma(X_k^{(h)}) \sqrt{h} \eta_{k+1}, \quad (2.2.7)$$

and the conditional variance process of the martingale part is

$$\langle M^{(h)} \rangle_n = \sum_{k=0}^{n-1} \sigma(X_k^{(h)})^2 \cdot h. \quad (2.2.8)$$

- (3). Conclude that

$$E[(M_n^{(h)} - x_0)^2] = \sum_{k=0}^{n-1} E[\sigma(X_k^{(h)})^2] \cdot h. \quad (2.2.9)$$

The last equation can be used in combination with the maximal inequality for martingales to derive bounds for the processes $(X^{(h)})$ in an efficient way, cf. Section 2.4 below.

Remark (Quadratic variation). The quadratic variation of a square integrable martingale (M_n) is the process $[M]_n$ defined by

$$[M]_n = \sum_{k=1}^n (M_k - M_{k-1})^2, \quad n \geq 0.$$

It is easy to verify that $M_n^2 - [M]_n$ is again a martingale. However, $[M]_n$ is not predictable. For continuous martingales in continuous time, the quadratic variation and the conditional variance process coincide. In discrete time or for discontinuous martingales they are usually different.

Martingale problem

For a Markov chain (X_n) we obtain a Doob decomposition

$$f(X_n) = M_n^{[f]} + A_n^{[f]} \quad (2.2.10)$$

for any function f on the state space such that $f(X_n)$ is integrable for each n . Computation of the predictable part leads to the following general result:

Theorem 2.5 (Martingale problem for time-homogeneous Markov chains). *Let p be a stochastic kernel on a measurable space (S, \mathcal{B}) . Then for an (\mathcal{F}_n) adapted stochastic process $(X_n)_{n \geq 0}$ with state space (S, \mathcal{B}) the following statements are equivalent:*

- (1). (X_n) is a time homogeneous (\mathcal{F}_n) Markov chain with transition kernel p .
- (2). (X_n) is a **solution of the martingale problem for the operator $\mathcal{L} = p - I$** , i.e., there is a decomposition

$$f(X_n) = M_n^{[f]} + \sum_{k=0}^{n-1} (\mathcal{L}f)(X_k), \quad n \geq 0,$$

with an (\mathcal{F}_n) martingale $(M_n^{[f]})$ for every function $f : S \rightarrow \mathbb{R}$ such that $f(X_n)$ is integrable for each n , or, equivalently, for every bounded function $f : S \rightarrow \mathbb{R}$.

In particular, we see once more that if $f(X_n)$ is integrable and f is harmonic ($\mathcal{L}f = 0$) then $f(X_n)$ is a martingale, and if f is superharmonic ($\mathcal{L}f \leq 0$), then $f(X_n)$ is a supermartingale. The theorem hence extends Theorem 2.3 above.

Proof. The implication “(i) \Rightarrow (ii)” is just the Doob decomposition for $f(X_n)$. In fact, by Theorem 2.4, the predictable part is given by

$$\begin{aligned} A_n^{[f]} &= \sum_{k=0}^{n-1} E[f(X_{k+1}) - f(X_k) \mid \mathcal{F}_k] \\ &= \sum_{k=0}^{n-1} (pf(X_k) - f(X_k)) = \sum_{k=0}^{n-1} (\mathcal{L}f)(X_k), \end{aligned}$$

and $M_n^{[f]} = f(X_n) - A_n^{[f]}$ is a martingale.

To prove the converse implication “(ii) \Rightarrow (i)” suppose that $M_n^{[f]}$ is a martingale for any bounded $f : S \rightarrow \mathbb{R}$. Then

$$\begin{aligned} 0 &= E[M_{n+1}^{[f]} - M_n^{[f]} \mid \mathcal{F}_n] \\ &= E[f(X_{n+1}) - f(X_n) \mid \mathcal{F}_n] - ((pf)(X_n) - f(X_n)) \\ &= E[f(X_{n+1}) \mid \mathcal{F}_n] - (pf)(X_n) \end{aligned}$$

almost surely for any bounded function f . Hence (X_n) is an (\mathcal{F}_n) Markov chain with transition kernel p . \square

Example (One dimensional Markov chains). Suppose that under P_x , the process (X_n) is a time homogeneous Markov chain with state space $S = \mathbb{R}$ or $S = \mathbb{Z}$, initial state $X_0 = x$, and transition kernel p . Assuming $X_n \in \mathcal{L}^2(\Omega, \mathcal{A}, P)$ for each n , we define the “drift” and the “fluctuations” of the process by

$$\begin{aligned} b(x) &:= E_x[X_1 - X_0] \\ a(x) &= \text{Var}_x[X_1 - X_0]. \end{aligned}$$

We now compute the Doob decomposition of (X_n) . Choosing $f(x) = x$ we have

$$(p - I)f(x) = \int y p(x, dy) - x = E_x[X_1 - X_0] = b(x).$$

Hence by Theorem 2.5,

$$X_n = M_n + \sum_{k=0}^{n-1} b(X_k) \tag{2.2.11}$$

with an (\mathcal{F}_n) martingale (M_n) . To obtain detailed information on M_n , we compute the variance process: By (2.2.11) and the Markov property, we obtain

$$\langle M \rangle_n = \sum_{k=0}^{n-1} \text{Var}[M_{k+1} - M_k \mid \mathcal{F}_k] = \sum_{k=0}^{n-1} \text{Var}[X_{k+1} - X_k \mid \mathcal{F}_k] = \sum_{k=0}^{n-1} a(X_k).$$

Therefore

$$M_n^2 = \widetilde{M}_n + \sum_{k=0}^{n-1} a(X_k) \quad (2.2.12)$$

with another (\mathcal{F}_n) martingale (\widetilde{M}_n) . The functions $a(x)$ and $b(x)$ can now be used in connection with fundamental results for martingales as e.g. the maximal inequality (cf. 2.4 below) to derive bounds for Markov chains in an efficient way.

2.3 Gambling strategies and stopping times

Throughout this section, we fix a filtration $(\mathcal{F}_n)_{n \geq 0}$ on a probability space (Ω, \mathcal{A}, P) .

Martingale transforms

Suppose that $(M_n)_{n \geq 0}$ is a martingale w.r.t. (\mathcal{F}_n) , and $(C_n)_{n \in \mathbb{N}}$ is a predictable sequence of real-valued random variables. For example, we may think of C_n as the stake in the n -th round of a fair game, and of the martingale increment $M_n - M_{n-1}$ as the net gain (resp. loss) per unit stake. In this case, the capital I_n of a player with gambling strategy (C_n) after n rounds is given recursively by

$$I_n = I_{n-1} + C_n \cdot (M_n - M_{n-1}) \quad \text{for any } n \in \mathbb{N},$$

i.e.,

$$I_n = I_0 + \sum_{k=1}^n C_k \cdot (M_k - M_{k-1}).$$

Definition (Martingale transform). *The stochastic process $C \bullet M$ defined by*

$$(C \bullet M)_n := \sum_{k=1}^n C_k \cdot (M_k - M_{k-1}) \quad \text{for any } n \geq 0,$$

*is called the **martingale transform** of the martingale $(M_n)_{n \geq 0}$ w.r.t. the predictable sequence $(C_n)_{n \geq 1}$, or the discrete stochastic integral of C w.r.t. M .*

We will see later that the process $C \bullet M$ is a time-discrete version of the stochastic integral $\int C_s dM_s$ of a predictable continuous-time process C w.r.t. a continuous-time martingale M . To be precise, $(C \bullet M)_n$ coincides with the Itô integral $\int_0^n C_{[t]} dM_{[t]}$ of the left continuous jump process $t \mapsto C_{[t]}$ w.r.t. the right continuous martingale $t \mapsto M_{[t]}$.

Example (Martingale strategy). One origin of the word “martingale” is the name of a well-known gambling strategy: In a standard coin-tossing game, the stake is doubled each time a loss occurs, and the player stops the game after the first time he wins. If the net gain in n rounds with unit stake is given by a standard Random Walk


$$M_n = \eta_1 + \dots + \eta_n, \quad \eta_i \text{ i.i.d. with } P[\eta_i = 1] = P[\eta_i = -1] = 1/2,$$

then the stake in the n -th round is

$$C_n = 2^{n-1} \quad \text{if } \eta_1 = \dots = \eta_{n-1} = -1, \quad \text{and} \quad C_n = 0 \quad \text{otherwise.}$$

Clearly, with probability one, the game terminates in finite time, and at that time the player has always won one unit, i.e.,

$$P[(C \bullet M)_n = 1 \text{ eventually}] = 1.$$



At first glance this looks like a safe winning strategy, but of course this would only be the case, if the player had unlimited capital and time available.

Theorem 2.6 (You can't beat the system!). (1). If $(M_n)_{n \geq 0}$ is an (\mathcal{F}_n) martingale, and $(C_n)_{n \geq 1}$ is predictable with $C_n \cdot (M_n - M_{n-1}) \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$ for any $n \geq 1$, then $C \bullet M$ is again an (\mathcal{F}_n) martingale.

(2). If (M_n) is an (\mathcal{F}_n) supermartingale and $(C_n)_{n \geq 1}$ is non-negative and predictable with $C_n \cdot (M_n - M_{n-1}) \in \mathcal{L}^1$ for any n , then $C \bullet M$ is again a supermartingale.

Proof. For $n \geq 1$ we have

$$\begin{aligned} E[(C \bullet M)_n - (C \bullet M)_{n-1} \mid \mathcal{F}_{n-1}] &= E[C_n \cdot (M_n - M_{n-1}) \mid \mathcal{F}_{n-1}] \\ &= C_n \cdot E[M_n - M_{n-1} \mid \mathcal{F}_{n-1}] = 0 \quad P\text{-a.s.} \end{aligned}$$

This proves the first part of the claim. The proof of the second part is similar. \square

The theorem shows that a fair game (a martingale) can not be transformed by choice of a clever gambling strategy into an unfair (or “superfair”) game. In models of financial markets this fact is crucial to exclude the existence of arbitrage possibilities (riskless profit).

Example (Martingale strategy, cont.). For the classical martingale strategy, we obtain

$$E[(C \bullet M)_n] = E[(C \bullet M)_0] = 0 \quad \text{for any } n \geq 0$$

by the martingale property, although

$$\lim_{n \rightarrow \infty} (C \bullet M)_n = 1 \quad P\text{-almost surely.}$$

This is a classical example showing that the assertion of the dominated convergence theorem may not hold if the assumptions are violated.

Remark. The integrability assumption in Theorem 2.6 is always satisfied if the random variables C_n are bounded, or if both C_n and M_n are square-integrable for any n .

Example (Financial market model with one risky asset). Suppose that during each time interval $(n - 1, n)$, an investor is holding Φ_n units of an asset with price S_n per unit at time n . We assume that (S_n) is an adapted and (Φ_n) is a predictable stochastic process w.r.t. a filtration (\mathcal{F}_n) . If the investor always puts his remaining capital onto a bank account with guaranteed interest rate r (“riskless asset”) then the change of his capital V_n during the time interval $(n - 1, n)$ is given by

$$V_n = V_{n-1} + \Phi_n \cdot (S_n - S_{n-1}) + (V_{n-1} - \Phi_n \cdot S_{n-1}) \cdot r. \quad (2.3.1)$$

Considering the discounted quantity $\tilde{V}_n = V_n / (1 + r)^n$, we obtain the equivalent recursion

$$\tilde{V}_n = \tilde{V}_{n-1} + \Phi_n \cdot (\tilde{S}_n - \tilde{S}_{n-1}) \quad \text{for any } n \geq 1. \quad (2.3.2)$$

In fact, (2.3.1) holds if and only if

$$V_n - (1 + r)V_{n-1} = \Phi_n \cdot (S_n - (1 + r)S_{n-1}),$$

which is equivalent to (2.3.2). Therefore, the discounted capital at time n is given by

$$\tilde{V}_n = V_0 + (\Phi \bullet \tilde{S})_n.$$

By Theorem 2.6, we can conclude that, if the discounted price process (\tilde{S}_n) is an (\mathcal{F}_n) martingale w.r.t. a given probability measure, then (\tilde{V}_n) is a martingale as well. In this case, assuming that V_0 is constant, we obtain in particular

$$E[\tilde{V}_n] = V_0,$$

or, equivalently,

$$E[V_n] = (1+r)^n V_0 \quad \text{for any } n \geq 0. \quad (2.3.3)$$

This fact, together with the existence of a martingale measure, can now be used for option pricing under a **no-arbitrage assumption**. To this end we assume that the payoff of an option at time N is given by an (\mathcal{F}_N) -measurable random variable F . For example, the payoff of a European call option with strike price K based on the asset with price process (S_n) is $S_N - K$ if the price S_N at maturity exceeds K , and 0 otherwise, i.e.,

$$F = (S_N - K)^+.$$

Suppose further that the option can be *replicated by a hedging strategy* (Φ_n) , i.e., there exists an \mathcal{F}_0 -measurable random variable V_0 and a predictable sequence of random variables $(\Phi_n)_{1 \leq n \leq N}$ such that

$$F = V_N$$

is the value at time N of a portfolio with initial value V_0 w.r.t. the trading strategy (Φ_n) . Then, assuming the non-existence of arbitrage possibilities, the option price at time 0 has to be V_0 , since otherwise one could construct an arbitrage strategy by selling the option and investing money in the stock market with strategy (Φ_n) , or conversely. Therefore, if a martingale measure exists (i.e., an underlying probability measure such that the discounted stock price (\tilde{S}_n) is a martingale), then the no-arbitrage price of the option at time 0 can be computed by (2.3.3) where the expectation is taken w.r.t. the martingale measure.

The following exercise shows how this works out in the Cox-Ross-Rubinstein binomial model:

Exercise (No-Arbitrage Pricing in the CRR model). Consider the CRR binomial model, i.e., $\Omega = \{1 + a, 1 + b\}^N$ with $-1 < a < r < b < \infty$, $X_i(\omega_1, \dots, \omega_N) = \omega_i$, $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$, and

$$S_n = S_0 \cdot \prod_{i=1}^n X_i, \quad n = 0, 1, \dots, N,$$

where S_0 is a constant.

- (1). *Completeness of the CRR model:* Prove that for any function $F : \Omega \rightarrow \mathbb{R}$ there exists a constant V_0 and a predictable sequence $(\Phi_n)_{1 \leq n \leq N}$ such that $F = V_N$ where $(V_n)_{1 \leq n \leq N}$ is defined by (2.3.1), or, equivalently,

$$\frac{F}{(1+r)^N} = \tilde{V}_N = V_0 + (\Phi \bullet \tilde{S})_N.$$

Hence in the CRR model, any \mathcal{F}_N -measurable function F can be replicated by a predictable trading strategy. Market models with this property are called *complete*.

Hint: Prove inductively that for $n = N, N-1, \dots, 0$, $\tilde{F} = F/(1+r)^N$ can be represented as

$$\tilde{F} = \tilde{V}_n + \sum_{i=n+1}^N \Phi_i \cdot (\tilde{S}_i - \tilde{S}_{i-1})$$

with an \mathcal{F}_n -measurable function \tilde{V}_n and a predictable sequence $(\Phi_i)_{n+1 \leq i \leq N}$.

- (2). *Option pricing:* Derive a general formula for the no-arbitrage price of an option with payoff function $F : \Omega \rightarrow \mathbb{R}$ in the CRR model. Compute the no-arbitrage price for a European call option with maturity N and strike K explicitly.

Stopped Martingales

One possible strategy for controlling a fair game is to terminate the game at a time depending on the previous development. Recall that a random variable $T : \Omega \rightarrow \{0, 1, 2, \dots\} \cup \{\infty\}$ is called a **stopping time** w.r.t. the filtration (\mathcal{F}_n) if and only if the event $\{T = n\}$ is contained in \mathcal{F}_n for any $n \geq 0$, or equivalently, iff $\{T \leq n\} \in \mathcal{F}_n$ for any $n \geq 0$.

Example (Hitting times). (1). The **first hitting time**

$$T_B = \min\{n \geq 0 : X_n \in B\} \quad (\text{where } \min \emptyset := \infty)$$

and the **first passage or return time**

$$S_B = \min\{n \geq 1 : X_n \in B\}$$

to a measurable subset B of the state space by an (\mathcal{F}_n) adapted stochastic process are (\mathcal{F}_n) stopping times. For example, for $n \geq 0$,

$$\{T_B = n\} = \{X_1 \in B^C, \dots, X_{n-1} \in B^C, X_n \in B\} \in \mathcal{F}_n.$$

If one decides to sell an asset as soon as the price S_n exceeds a given level $\lambda > 0$ then the selling time equals $T_{(\lambda, \infty)}$ and is hence a stopping time.

(2). On the other hand, the **last visit time**

$$L_B := \sup\{n \geq 0 : X_n \in B\} \quad (\text{where } \sup \emptyset := 0)$$

is not a stopping time in general. Intuitively, to decide whether $L_B = n$, information on the future development of the process is required.

We consider an (\mathcal{F}_n) -adapted stochastic process $(M_n)_{n \geq 0}$, and an (\mathcal{F}_n) -stopping time T on the probability space (Ω, \mathcal{A}, P) . The process stopped at time T is defined as $(M_{T \wedge n})_{n \geq 0}$ where

$$M_{T \wedge n}(\omega) = M_{T(\omega) \wedge n}(\omega) = \begin{cases} M_n(\omega) & \text{for } n \leq T(\omega), \\ M_{T(\omega)}(\omega) & \text{for } n \geq T(\omega). \end{cases}$$

For example, the process stopped at a hitting time T_B gets stuck at the first time it enters the set B .

Theorem 2.7 (Optional Stopping Theorem, Version 1). *If $(M_n)_{n \geq 0}$ is a martingale (resp. a supermartingale) w.r.t. (\mathcal{F}_n) , and T is an (\mathcal{F}_n) -stopping time, then the stopped process $(M_{T \wedge n})_{n \geq 0}$ is again an (\mathcal{F}_n) -martingale (resp. supermartingale). In particular, we have*

$$E[M_{T \wedge n}] \stackrel{(\leq)}{=} E[M_0] \quad \text{for any } n \geq 0.$$

Proof. Consider the following strategy:

$$C_n = I_{\{T \geq n\}} = 1 - I_{\{T \leq n-1\}},$$

i.e., we put a unit stake in each round before time T and quit playing at time T . Since T is a stopping time, the sequence (C_n) is predictable. Moreover,

$$M_{T \wedge n} - M_0 = (C \bullet M)_n \quad \text{for any } n \geq 0. \quad (2.3.4)$$

In fact, for the increments of the stopped process we have

$$M_{T \wedge n} - M_{T \wedge (n-1)} = \begin{cases} M_n - M_{n-1} & \text{if } T \geq n \\ 0 & \text{if } T \leq n-1 \end{cases} = C_n \cdot (M_n - M_{n-1}),$$

and (2.3.4) follows by summing over n . Since the sequence (C_n) is predictable, bounded and non-negative, the process $C \bullet M$ is a martingale, supermartingale respectively, provided the same holds for M . \square

Remark (IMPORTANT). (1). In general, it is NOT TRUE under the assumptions in Theorem 2.7 that

$$E[M_T] = E[M_0], \quad E[M_T] \leq E[M_0] \quad \text{respectively.} \quad (2.3.5)$$

Suppose for example that (M_n) is the classical Random Walk starting at 0 and $T = T_{\{1\}}$ is the first hitting time of the point 1. Then, by recurrence of the Random Walk, $T < \infty$ and $M_T = 1$ hold almost surely although $M_0 = 0$.

(2). If, on the other hand, T is a *bounded stopping time*, then there exists $n \in \mathbb{N}$ such that $T(\omega) \leq n$ for any ω . In this case, the optional stopping theorem implies

$$E[M_T] = E[M_{T \wedge n}] \stackrel{(\leq)}{=} E[M_0].$$

More general sufficient conditions for (2.3.5) are given in Theorems 2.8, 2.9 and 2.10 below.

Example (Classical ruin problem). Let $a, b, x \in \mathbb{Z}$ with $a < x < b$. We consider the classical Random Walk

$$X_n = x + \sum_{i=1}^n \eta_i, \quad \eta_i \text{ i.i.d. with } P[\eta_i = \pm 1] = \frac{1}{2},$$

with initial value $X_0 = x$. We now show how to apply the Optional Stopping Theorem to compute the distributions of the exit time

$$T(\omega) = \min\{n \geq 0 : X_n(\omega) \notin (a, b)\},$$

and the exit point X_T . These distributions can also be computed by more traditional methods (first step analysis, reflection principle), but martingales yield an elegant and general approach.

(1). *Ruin probability* $r(x) = P[X_T = a]$.

The process (X_n) is a martingale w.r.t. the filtration $\mathcal{F}_n = \sigma(\eta_1, \dots, \eta_n)$, and $T < \infty$ almost surely holds by elementary arguments. As the stopped process $X_{T \wedge n}$ is bounded ($a \leq X_{T \wedge n} \leq b$), we obtain

$$x = E[X_0] = E[X_{T \wedge n}] \xrightarrow{n \rightarrow \infty} E[X_T] = a \cdot r(x) + b \cdot (1 - r(x))$$

by the Optional Stopping Theorem and the Dominated Convergence Theorem. Hence

$$r(x) = \frac{b - x}{a - x}. \quad (2.3.6)$$

(2). *Mean exit time from* (a, b) .

To compute the expectation $E[T]$, we apply the Optional Stopping Theorem to the (\mathcal{F}_n) martingale

$$M_n := X_n^2 - n.$$

By monotone and dominated convergence, we obtain

$$\begin{aligned} x^2 &= E[M_0] = E[M_{T \wedge n}] = E[X_{T \wedge n}^2] - E[T \wedge n] \\ &\xrightarrow{n \rightarrow \infty} E[X_T^2] - E[T]. \end{aligned}$$

Therefore, by (2.3.6),

$$\begin{aligned} E[T] &= E[X_T^2] - x^2 = a^2 \cdot r(x) + b^2 \cdot (1 - r(x)) - x^2 \\ &= (b - x) \cdot (x - a). \end{aligned} \quad (2.3.7)$$

(3). *Mean passage time of b .*

The first passage time $T_b = \min\{n \geq 0 : X_n = b\}$ is greater or equal than the exit time from the interval (a, b) for any $a < x$. Thus by (2.3.7), we have

$$E[T_b] \geq \lim_{a \rightarrow -\infty} (b - x) \cdot (x - a) = \infty,$$

i.e., T_b is **not integrable!** These and some other related passage times are important examples of random variables with a heavy-tailed distribution and infinite first moment.

(4). *Distribution of passage times.*

We now compute the distribution of the first passage time T_b explicitly in the case $x = 0$ and $b = 1$. Hence let $T = T_1$. As shown above, the process

$$M_n^\lambda := e^{\lambda X_n} / (\cosh \lambda)^n, \quad n \geq 0,$$

is a martingale for each $\lambda \in \mathbb{R}$. Now suppose $\lambda > 0$. By the Optional Stopping Theorem,

$$1 = E[M_0^\lambda] = E[M_{T \wedge n}^\lambda] = E[e^{\lambda X_{T \wedge n}} / (\cosh \lambda)^{T \wedge n}] \quad (2.3.8)$$

for any $n \in \mathbb{N}$. As $n \rightarrow \infty$, the integrands on the right hand side converge to $e^{\lambda(\cosh \lambda)^{-T}} \cdot I_{\{T < \infty\}}$. Moreover, they are uniformly bounded by e^λ , since $X_{T \wedge n} \leq 1$ for any n . Hence by the Dominated Convergence Theorem, the expectation on the right hand side of (2.3.8) converges to $E[e^\lambda / (\cosh \lambda)^T ; T < \infty]$, and we obtain the identity

$$E[(\cosh \lambda)^{-T} ; T < \infty] = e^{-\lambda} \quad \text{for any } \lambda > 0. \quad (2.3.9)$$

Taking the limit as $\lambda \searrow 0$, we see that $P[T < \infty] = 1$. Taking this into account, and substituting $s = 1/\cosh \lambda$ in (2.3.9), we can now compute the generating function of T explicitly:

$$E[s^T] = e^{-\lambda} = (1 - \sqrt{1 - s^2})/s \quad \text{for any } s \in (0, 1). \quad (2.3.10)$$

Developing both sides into a power series finally yields

$$\sum_{n=0}^{\infty} s^n \cdot P[T = n] = \sum_{m=1}^{\infty} (-1)^{m+1} \binom{1/2}{m} s^{2m-1}.$$

Therefore, the distribution of the first passage time of 1 is given by

$$P[T = 2m-1] = (-1)^{m+1} \binom{1/2}{m} = (-1)^{m+1} \cdot \frac{1}{2} \cdot \left(-\frac{1}{2}\right) \cdots \left(\frac{1}{2} - m + 1\right) / m!$$

and $P[T = 2m] = 0$ for any $m \in \mathbb{N}$.

Optional Stopping Theorems

Stopping times occurring in applications are typically not bounded. Therefore, we need more general conditions guaranteeing that (2.3.5) holds nevertheless. A first general criterion is obtained by applying the Dominated Convergence Theorem:

Theorem 2.8 (Optional Stopping Theorem, Version 2). *Suppose that (M_n) is a martingale w.r.t. (\mathcal{F}_n) , T is an (\mathcal{F}_n) -stopping time with $P[T < \infty] = 1$, and there exists a random variable $Y \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$ such that*

$$|M_{T \wedge n}| \leq Y \quad P\text{-almost surely for any } n \in \mathbb{N}.$$

Then

$$E[M_T] = E[M_0].$$

Proof. Since $P[T < \infty] = 1$, we have

$$M_T = \lim_{n \rightarrow \infty} M_{T \wedge n} \quad P\text{-almost surely.}$$

By Theorem 2.7, $E[M_0] = E[M_{T \wedge n}]$, and by the Dominated Convergence Theorem, $E[M_{T \wedge n}] \rightarrow E[M_T]$ as $n \rightarrow \infty$. \square

Remark (Weakening the assumptions). Instead of the existence of an integrable random variable Y dominating the random variables $M_{T \wedge n}$, $n \in \mathbb{N}$, it is enough to assume that these random variables are **uniformly integrable**, i.e.,

$$\sup_{n \in \mathbb{N}} E[|M_{T \wedge n}|; |M_{T \wedge n}| \geq c] \rightarrow 0 \quad \text{as } c \rightarrow \infty.$$

A corresponding generalization of the Dominated Convergence Theorem is proven in Section 4.3 below.

For non-negative supermartingales, we can apply Fatou's Lemma instead of the Dominated Convergence Theorem to pass to the limit as $n \rightarrow \infty$ in the Stopping Theorem. The advantage is that no integrability assumption is required. Of course, the price to pay is that we only obtain an inequality:

Theorem 2.9 (Optional Stopping Theorem, Version 3). *If (M_n) is a non-negative supermartingale w.r.t. (\mathcal{F}_n) , then*

$$E[M_0] \geq E[M_T; T < \infty]$$

holds for any (\mathcal{F}_n) stopping time T .

Proof. Since $M_T = \lim_{n \rightarrow \infty} M_{T \wedge n}$ on $\{T < \infty\}$, and $M_T \geq 0$, Theorem 2.7 combined with Fatou's Lemma implies

$$E[M_0] \geq \liminf_{n \rightarrow \infty} E[M_{T \wedge n}] \geq E\left[\liminf_{n \rightarrow \infty} M_{T \wedge n}\right] \geq E[M_T; T < \infty].$$

□

Example (Dirichlet problem for Markov chains). Suppose that w.r.t. the probability measure P_x , the process (X_n) is a time-homogeneous Markov chain with measurable state space (S, \mathcal{B}) , transition kernel p , and start in x . Let $D \in \mathcal{B}$ be a measurable subset of the state space, and $f : D^C \rightarrow \mathbb{R}$ a measurable function (the given “boundary values”), and let

$$T = \min\{n \geq 0 : X_n \in D^C\}$$


denote the first exit time of the Markov chain from D . By conditioning on the first step of the Markov chain, one can show that if f is non-negative or bounded, then the function

$$h(x) = E_x[f(X_T); T < \infty], \quad (x \in S),$$

is a solution of the *Dirichlet problem*

$$\begin{aligned} (ph)(x) &= h(x) && \text{for } x \in D, \\ h(x) &= f(x) && \text{for } x \in D^C, \end{aligned}$$

see [XXXStochastic Processes].



By considering the martingale $h(X_{T \wedge n})$ for a function h that is harmonic on D , we obtain a converse statement:

Exercise (Uniqueness of the Dirichlet problem). Suppose that $P_x[T < \infty] = 1$ for any $x \in S$.

- (1). Prove that $h(X_{T \wedge n})$ is a martingale w.r.t. P_x for any bounded solution h of the Dirichlet problem and any $x \in S$.
- (2). Conclude that if f is bounded, then

$$h(x) = E_x[f(X_T)] \quad (2.3.11)$$

is the unique bounded solution of the Dirichlet problem.

- (3). Similarly, show that for any non-negative f , the function h defined by (2.3.11) is the minimal non-negative solution of the Dirichlet problem.

We finally state a version of the Optional Stopping Theorem that applies in particular to martingales with bounded increments:

Corollary 2.10 (Optional Stopping for martingales with bounded increments). *Suppose that (M_n) is an (\mathcal{F}_n) martingale, and there exists a finite constant $K \in (0, \infty)$ such that*

$$E[|M_{n+1} - M_n| \mid \mathcal{F}_n] \leq K \quad P\text{-almost surely for any } n \geq 0. \quad (2.3.12)$$

Then for any (\mathcal{F}_n) stopping time T with $E[T] < \infty$, we have

$$E[M_T] = E[M_0].$$

Proof. For any $n \geq 0$,

$$|M_{T \wedge n}| \leq |M_0| + \sum_{i=0}^{\infty} |M_{i+1} - M_i| \cdot I_{\{T > i\}}.$$

Let Y denote the expression on the right hand side. We will show that Y is an integrable random variable – this implies the assertion by Theorem 2.8. To verify integrability of Y note that the event $\{T > i\}$ is contained in \mathcal{F}_i for any $i \geq 0$ since T is a stopping time. Therefore and by (2.3.12),

$$E[|M_{i+1} - M_i| ; T > i] = E[E[|M_{i+1} - M_i| \mid \mathcal{F}_i] ; T > i] \leq k \cdot P[T > i].$$

Summing over i , we obtain

$$E[Y] \leq E[|M_0|] + k \cdot \sum_{i=0}^{\infty} P[T > i] = E[|M_0|] + k \cdot E[T] < \infty$$

by the assumptions. □

Exercise (Integrability of stopping times). Prove that the expectation $E[T]$ of a stopping time T is finite if there exist constants $\varepsilon > 0$ and $k \in \mathbb{N}$ such that

$$P[T \leq n + k \mid \mathcal{F}_n] \geq \varepsilon \quad P\text{-a.s. for any } n \in \mathbb{N}.$$

Wald's identity for random sums

We finally apply the Optional Stopping Theorem to sums of independent random variables with a random number T of summands. The point is that we do not assume that T is independent of the summands but only that it is a stopping time w.r.t. the filtration generated by the summands.

Let $S_n = \eta_1 + \dots + \eta_n$ with i.i.d. random variables $\eta_i \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$. Denoting by m the expectations of the increments η_i , the process

$$M_n = S_n - n \cdot m$$

is a martingale w.r.t. $\mathcal{F}_n = \sigma(\eta_1, \dots, \eta_n)$. By applying Corollary 2.10 to this martingale, we obtain:

Theorem 2.11 (Wald's identity). *Suppose that T is an (\mathcal{F}_n) stopping time with $E[T] < \infty$. Then*

$$E[S_T] = m \cdot E[T].$$

Proof. For any $n \geq 0$, we have

$$E[|M_{n+1} - M_n| | \mathcal{F}_n] = E[|\eta_{n+1} - m| | \mathcal{F}_n] = E[|\eta_{n+1} - m|]$$

by the independence of the η_i . As the η_i are identically distributed and integrable, the right hand side is a finite constant. Hence Corollary 2.10 applies, and we obtain

$$0 = E[M_0] = E[M_T] = E[S_T] - m \cdot E[T].$$

□

2.4 Maximal inequalities

For a standard Random Walk $S_n = \eta_1 + \dots + \eta_n$, η_i i.i.d. with $P[\eta_i = \pm 1] = 1/2$, the reflection principle implies the identity

$$\begin{aligned} P[\max(S_0, S_1, \dots, S_n) \geq c] &= P[S_n \geq c] + P[S_n < c; \max(S_0, S_1, \dots, S_n) \geq c] \\ &= P[|S_n| > c] + P[S_n > c] \end{aligned}$$

for any $c \in \mathbb{N}$. In combination with the Markov-Čebyšev inequality this can be used to control the running maximum of the Random Walk in terms of the moments of the last value S_n .

Maximal inequalities are corresponding estimates for $\max(M_0, M_1, \dots, M_n)$ or $\sup_{k \geq 0} M_k$ when (M_n) is a sub- or supermartingale respectively. These estimates are an important tool in stochastic analysis. They are a consequence of the Optional Stopping Theorem.

Doob's inequality

We first prove the basic version of maximal inequalities for sub- and supermartingales:

Theorem 2.12 (Doob).

(1). Suppose that $(M_n)_{n \geq 0}$ is a non-negative supermartingale. Then

$$P \left[\sup_{k \geq 0} M_k \geq c \right] \leq \frac{1}{c} \cdot E[M_0] \quad \text{for any } c > 0.$$

(2). Suppose that $(M_n)_{n \geq 0}$ is a non-negative submartingale. Then

$$P \left[\max_{0 \leq k \leq n} M_k \geq c \right] \leq \frac{1}{c} \cdot E \left[M_n ; \max_{0 \leq k \leq n} M_k \geq c \right] \leq \frac{1}{c} \cdot E[M_n] \quad \text{for any } c > 0.$$

Proof. (1). For $c > 0$ we consider the stopping time

$$T_c = \min\{k \geq 0 : M_k \geq c\}, \quad \min \emptyset = \infty.$$

Note that $T_c < \infty$ whenever $\sup M_k > c$. Hence by the version of the Optional Stopping Theorem for non-negative supermartingales, we obtain

$$P[\sup M_k > c] \leq P[T_c < \infty] \leq \frac{1}{c} E[M_{T_c} ; T_c < \infty] \leq \frac{1}{c} E[M_0].$$

Here we have used in the second and third step that (M_n) is non-negative. Replacing c by $c - \varepsilon$ and letting ε tend to zero we can conclude

$$P[\sup M_k \geq c] = \lim_{\varepsilon \searrow 0} P[\sup M_k > c - \varepsilon] \leq \liminf_{\varepsilon \searrow 0} \frac{1}{c - \varepsilon} E[M_0] = \frac{1}{c} \cdot E[M_0].$$

(2). For a non-negative supermartingale, we obtain

$$\begin{aligned}
 P \left[\max_{0 \leq k \leq n} M_k \geq c \right] &= P[T_c \leq n] \leq \frac{1}{c} E[M_{T_c}; T_c \leq n] \\
 &= \frac{1}{c} \sum_{k=0}^n E[M_k; T_c = k] \leq \frac{1}{c} \sum_{k=0}^n E[M_n; T_c = k] \\
 &= \frac{1}{c} \cdot E[M_n; T_c \leq n].
 \end{aligned}$$

Here we have used in the second last step that $E[M_k; T_c = k] \leq E[M_n; T_c = k]$ since (M_n) is a supermartingale and $\{T_c = k\}$ is in \mathcal{F}_k . □

First consequences of Doob's maximal inequality for submartingales are extensions of the classical Markov-Čebyšev inequalities:

Corollary 2.13. (1). *Suppose that $(M_n)_{n \geq 0}$ is an arbitrary submartingale (not necessarily non-negative!). Then*

$$\begin{aligned}
 P \left[\max_{k \leq n} M_k \geq c \right] &\leq \frac{1}{c} E \left[M_n^+; \max_{k \leq n} M_k \geq c \right] \quad \text{for any } c > 0, \text{ and} \\
 P \left[\max_{k \leq n} M_k \geq c \right] &\leq e^{-\lambda c} E \left[e^{\lambda M_n}; \max_{k \leq n} M_k \geq c \right] \quad \text{for any } \lambda, c > 0.
 \end{aligned}$$

(2). *If (M_n) is a martingale then, moreover, the estimates*

$$P \left[\max_{k \leq n} |M_k| \geq c \right] \leq \frac{1}{c^p} E \left[|M_n|^p; \max_{k \leq n} |M_k| \geq c \right]$$

hold for any $c > 0$ and $p \in [1, \infty)$.

Proof. The corollary follows by applying the maximal inequality to the non-negative submartingales $M_n^+, \exp(\lambda M_n), |M_n|^p$ respectively. These processes are indeed submartingales, as the functions $x \mapsto x^+$ and $x \mapsto \exp(\lambda x)$ are convex and non-decreasing for any $\lambda > 0$, and the functions $x \mapsto |x|^p$ are convex for any $p \geq 1$. □

L^p inequalities

The last estimate in Corollary 2.13 can be used to bound the L^p norm of the running maximum of a martingale in terms of the L^p -norm of the last value. The resulting bound, known as Doob's L^p -inequality, is crucial for stochastic analysis. We first remark:

Lemma 2.14. *If $Y : \Omega \rightarrow \mathbb{R}_+$ is a non-negative random variable, and $G(y) = \int_0^y g(x) dx$ is the integral of a non-negative function $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, then*

$$E[G(Y)] = \int_0^\infty g(c) \cdot P[Y \geq c] dc.$$

Proof. By Fubini's theorem we have

$$\begin{aligned} E[G(Y)] &= E \left[\int_0^Y g(c) dc \right] = E \left[\int_0^\infty I_{[0,Y]}(c) g(c) dc \right] \\ &= \int_0^\infty g(c) \cdot P[Y \geq c] dc. \end{aligned}$$

□

Theorem 2.15 (Doob's L^p inequality). *Suppose that $(M_n)_{n \geq 0}$ is a martingale, and let*

$$M_n^* := \max_{k \leq n} |M_k|, \quad \text{and} \quad M^* := \sup_k |M_k|.$$

Then, for any $p, q \in (1, \infty)$ such that $\frac{1}{p} + \frac{1}{q} = 1$, we have

$$\|M_n^*\|_{L^p} \leq q \cdot \|M_n\|_{L^p}, \quad \text{and} \quad \|M^*\|_{L^p} \leq q \cdot \sup_n \|M_n\|_{L^p}.$$

In particular, if (M_n) is bounded in L^p then M^ is contained in L^p .*

Proof. By Lemma 2.14, Corollary 2.13 applied to the martingales M_n and $(-M_n)$, and Fubini's theorem, we have

$$\begin{aligned}
 E[(M_n^*)^p] &\stackrel{2.14}{=} \int_0^\infty pc^{p-1} \cdot P[M_n^* \geq c] \, dc \\
 &\stackrel{2.13}{\leq} \int_0^\infty pc^{p-2} E[|M_n|; M_n^* \geq c] \, dc \\
 &\stackrel{\text{Fub.}}{=} E \left[|M_n| \cdot \int_0^{M_n^*} pc^{p-2} \, dp \right] \\
 &= \frac{p}{p-1} E[|M_n| \cdot (M_n^*)^{p-1}]
 \end{aligned}$$

for any $n \geq 0$ and $p \in (1, \infty)$. Setting $q = \frac{p}{p-1}$ and applying Hölder's inequality to the right hand side, we obtain

$$E[(M_n^*)^p] \leq q \cdot \|M_n\|_{L^p} \cdot \|(M_n^*)^{p-1}\|_{L^q} = q \cdot \|M_n\|_{L^p} \cdot E[(M_n^*)^p]^{1/q},$$

i.e.,

$$\|M_n^*\|_{L^p} = E[(M_n^*)^p]^{1-1/q} \leq q \cdot \|M_n\|_{L^p}. \quad (2.4.1)$$

This proves the first inequality. The second inequality follows as $n \rightarrow \infty$, since

$$\|M^*\|_{L^p} = \left\| \lim_{n \rightarrow \infty} M_n^* \right\|_{L^p} = \liminf_{n \rightarrow \infty} \|M_n^*\|_{L^p} \leq q \cdot \sup_{n \in \mathbb{N}} \|M_n\|_{L^p}$$

by Fatou's Lemma. □

Hoeffding's inequality

For a standard Random Walk (S_n) starting at 0, the reflection principle combined with Bernstein's inequality implies the upper bound

$$P[\max(S_0, \dots, S_n) \geq c] \leq 2 \cdot P[S_n \geq c] \leq 2 \cdot \exp(-2c^2/n)$$

for any $n \in \mathbb{N}$ and $c \in (0, \infty)$. A similar inequality holds for arbitrary martingales with bounded increments:

Theorem 2.16 (Azuma, Hoeffding). *Suppose that (M_n) is a martingale such that*

$$|M_n - M_{n-1}| \leq a_n \quad P\text{-almost surely}$$

for a sequence (a_n) of non-negative constants. Then

$$P \left[\max_{k \leq n} (M_k - M_0) \geq c \right] \leq \exp \left(-\frac{1}{2} c^2 / \sum_{i=1}^n a_i^2 \right) \quad (2.4.2)$$

for any $n \in \mathbb{N}$ and $c \in (0, \infty)$.

Proof. W.l.o.g. we may assume $M_0 = 0$. Let $Y_n = M_n - M_{n-1}$ denote the martingale increments. We will apply the exponential form of the maximal inequality. For $\lambda > 0$ and $n \in \mathbb{N}$, we have,

$$E[e^{\lambda M_n}] = E \left[\prod_{i=1}^n e^{\lambda Y_i} \right] = E \left[e^{\lambda M_{n-1}} \cdot E \left[e^{\lambda Y_n} \mid \mathcal{F}_{n-1} \right] \right]. \quad (2.4.3)$$

To bound the conditional expectation, note that

$$e^{\lambda Y_n} \leq \frac{1}{2} \frac{a_n - Y_n}{a_n} e^{-\lambda a_n} + \frac{1}{2} \frac{a_n + Y_n}{a_n} e^{\lambda a_n}$$

holds almost surely, since $x \mapsto \exp(\lambda x)$ is a convex function, and $-a_n \leq Y_n \leq a_n$. Indeed, the right hand side is the value at Y_n of the secant connecting the points $(-a_n, \exp(-\lambda a_n))$ and $(a_n, \exp(\lambda a_n))$. Since (M_n) is a martingale, we have

$$E[Y_n \mid \mathcal{F}_{n-1}] = 0,$$

and therefore

$$E[e^{\lambda Y_n} \mid \mathcal{F}_{n-1}] \leq (e^{-\lambda a_n} + e^{\lambda a_n}) / 2 = \cosh(\lambda a_n) \leq e^{(\lambda a_n)^2 / 2}$$

almost surely. Now, by (2.4.3), we obtain

$$E[e^{\lambda Y_n}] \leq E[e^{\lambda M_{n-1}}] \cdot e^{(\lambda a_n)^2 / 2}.$$

Hence, by induction on n ,

$$E[e^{\lambda M_n}] \leq \exp\left(\frac{1}{2}\lambda^2 \sum_{i=1}^n a_i^2\right) \quad \text{for any } n \in \mathbb{N}, \quad (2.4.4)$$

and, by the exponential maximal inequality (cf. Corollary 2.13),

$$P[\max_{k \leq n} M_k \geq c] \leq \exp\left(-\lambda c + \frac{1}{2}\lambda^2 \sum_{i=1}^n a_i^2\right) \quad (2.4.5)$$

holds for any $n \in \mathbb{N}$ and $c, \lambda > 0$.

For a given c and n , the expression on the right hand side of (2.4.5) is minimal for $\lambda = c / \sum_{i=1}^n a_i^2$. Choosing λ correspondingly, we finally obtain the upper bound (2.4.2). \square

Hoeffding's concentration inequality has numerous applications, for example in the analysis of algorithms, cf. [Mitzenmacher, Upful: Probability and Computing]. Here, we just consider one simple example to illustrate the way it typically is applied:

Example (Pattern Matching). Suppose that X_1, X_2, \dots, X_n is a sequence of independent, uniformly distributed random variables ("letters") taking value in a finite set S (the underlying "alphabet"), and let

$$N = \sum_{i=0}^{n-l} I_{\{X_{i+1}=a_1, X_{i+2}=a_2, \dots, X_{i+l}=a_l\}} \quad (2.4.6)$$

denote the number of occurrences of a given "word" $a_1 a_2 \cdots a_l$ with l letters in the random text. In applications, the "word" could for example be a DNA sequence. We easily obtain

$$E[N] = \sum_{i=0}^{n-l} P[X_{i+k} = a_k \text{ for } k = 1, \dots, l] = (n-l+1)/|S|^l. \quad (2.4.7)$$

To estimate the fluctuations of the random variable N around its mean value, we consider the martingale

$$M_i = E[N \mid \sigma(X_1, \dots, X_i)], \quad (i = 0, 1, \dots, n)$$

with initial value $M_0 = E[N]$ and terminal value $M_n = N$. Since at most l of the summands in (2.4.6) are not independent of i , and each summand takes values 0 and 1 only, we have

$$|M_i - M_{i-1}| \leq l \quad \text{for each } i = 0, 1, \dots, n.$$

Therefore, by Hoeffding's inequality, applied in both directions, we obtain

$$P[|N - E[N]| \geq c] = P[|M_n - M_0| \geq c] \leq 2 \exp(-c^2/(2nl^2))$$

for any $c > 0$, or equivalently,

$$P[|N - E[N]| \geq \varepsilon \cdot l\sqrt{n}] \leq 2 \cdot \exp(-\varepsilon^2/2) \quad \text{for any } \varepsilon > 0. \quad (2.4.8)$$

The equation (2.4.7) and the bound (2.4.8) show that N is highly concentrated around its mean if l is small compared to \sqrt{n} .

Chapter 3

Martingales in continuous time

The notion of a martingale, sub- and supermartingale in continuous time can be defined similarly as in the discrete parameter case. Fundamental results such as the optional stopping theorem or the maximal inequality carry over from discrete parameter to continuous time martingales under additional regularity conditions as, for example, continuity of the sample paths. Similarly as for Markov chains in discrete time, martingale methods can be applied to derive explicit expressions and bounds for probabilities and expectations of Brownian motion in a clear and efficient way.

We start with the definition of martingales in continuous time. Let (Ω, \mathcal{A}, P) denote a probability space.

Definition. (1). A continuous-time **filtration** on (Ω, \mathcal{A}) is a family $(\mathcal{F}_t)_{t \in [0, \infty)}$ of σ -algebras $\mathcal{F}_t \subseteq \mathcal{A}$ such that $\mathcal{F}_s \subseteq \mathcal{F}_t$ for any $0 \leq s \leq t$.

(2). A real-valued stochastic process $(M_t)_{t \in [0, \infty)}$ on (Ω, \mathcal{A}, P) is called a **martingale** (or **super-, submartingale**) w.r.t. a filtration (\mathcal{F}_t) if and only if

(a) (M_t) is adapted w.r.t. (\mathcal{F}_t) , i.e., M_t is \mathcal{F}_t measurable for any $t \geq 0$.

(b) For any $t \geq 0$, the random variable M_t (resp. M_t^+, M_t^-) is integrable.

(c) $E[M_t | \mathcal{F}_s] \stackrel{(\leq, \geq)}{=} M_s$ P -almost surely for any $0 \leq s \leq t$.

3.1 Some fundamental martingales of Brownian Motion

In this section, we identify some important martingales that are functions of Brownian motion. Let $(B_t)_{t \geq 0}$ denote a d -dimensional Brownian motion defined on (Ω, \mathcal{A}, P) .

Filtrations generated by Brownian motion

Any stochastic process $(X_t)_{t \geq 0}$ in continuous time generates a filtration

$$\mathcal{F}_t^X = \sigma(X_s : 0 \leq s \leq t), \quad t \geq 0.$$

However, not every hitting time that we are interested in is a stopping time w.r.t. this filtration. For example, for one-dimensional Brownian motion (B_t) , the first hitting time $T = \inf\{t \geq 0 : B_t > c\}$ of the *open* interval (c, ∞) is not an (\mathcal{F}_t^B) stopping time. An intuitive explanation for this fact is that for $t \geq 0$, the event $\{T \leq t\}$ is not contained in \mathcal{F}_t^B , since for a path with $B_s \leq c$ on $[0, t]$ and $B_t = c$, we can not decide at time t , if the path will enter the interval (c, ∞) in the next instant. For this and other reasons, we also consider the right-continuous filtration

$$\mathcal{F}_t := \bigcap_{\varepsilon > 0} \mathcal{F}_{t+\varepsilon}^B, \quad t \geq 0,$$

that takes into account “infinitesimal information on the future development.”

Exercise (Hitting times as stopping times). Prove that the first hitting time $T_A = \inf\{t \geq 0 : B_t \in A\}$ of a set $A \subseteq \mathbb{R}^d$ is an (\mathcal{F}_t^B) stopping time if A is closed, whereas T_A is an (\mathcal{F}_t) stopping time, but not necessarily an (\mathcal{F}_t^B) stopping time if A is open.

It is easy to verify that a d -dimensional Brownian motion (B_t) is also a Brownian motion w.r.t. the right-continuous filtration (\mathcal{F}_t) :

Lemma 3.1. *For any $0 \leq s < t$, the increment $B_t - B_s$ is independent of \mathcal{F}_s with distribution $N(0, (t - s) \cdot I_d)$.*

Proof. Since $t \mapsto B_t$ is almost surely continuous, we have

$$B_t - B_s = \lim_{\substack{\varepsilon \searrow 0 \\ \varepsilon \in \mathbb{Q}}} (B_t - B_{s+\varepsilon}) \quad P\text{-a.s.} \quad (3.1.1)$$

For small $\varepsilon > 0$ the increment $B_t - B_{s+\varepsilon}$ is independent of $\mathcal{F}_{s+\varepsilon}^B$, and hence independent of \mathcal{F}_s . Therefore, by (3.1.1), $B_t - B_s$ is independent of \mathcal{F}_s as well. \square

Another filtration of interest is the completed filtration (\mathcal{F}_t^P) . A σ -algebra \mathcal{F} is called **complete** w.r.t. a probability measure P iff it contains all subsets of P -measure zero sets. The **completion** of a σ -algebra \mathcal{A} w.r.t. a probability measure P on (Ω, \mathcal{A}) is the complete σ -algebra

$$\mathcal{A}^P = \{A \subseteq \Omega : \exists A_1, A_2 \in \mathcal{A} : A_1 \subseteq A \subseteq A_2, P[A_2 \setminus A_1] = 0\}$$

generated by all sets in \mathcal{A} and all subsets of P -measure zero sets in \mathcal{A} .

It can be shown that the completion (\mathcal{F}_t^P) of the right-continuous filtration (\mathcal{F}_t) is again right-continuous. The assertion of Lemma 3.1 obviously carries over to the completed filtration.

Remark (The “usual conditions”). Some textbooks on stochastic analysis consider only complete right-continuous filtrations. A filtration with these properties is said to **satisfy the usual conditions**. A disadvantage of completing the filtration, however, is that (\mathcal{F}_t^P) depends on the underlying probability measure P (or, more precisely, on its null sets). This can cause problems when considering several non-equivalent probability measures at the same time.

Brownian Martingales

We now identify some basic martingales of Brownian motion:

Theorem 3.2 (Elementary martingales of Brownian motion). *For a d -dimensional Brownian motion (B_t) the following processes are martingales w.r.t. each of the filtrations (\mathcal{F}_t^B) , (\mathcal{F}_t) and (\mathcal{F}_t^P) :*

- (1). *The coordinate processes $B_t^{(i)}$, $1 \leq i \leq d$.*
- (2). *$B_t^{(i)} B_t^{(j)} - t \cdot \delta_{ij}$ for any $1 \leq i, j \leq d$.*

(3). $\exp(\alpha \cdot B_t - \frac{1}{2}|\alpha|^2 t)$ for any $\alpha \in \mathbb{R}^d$.

The processes $M_t^\alpha = \exp(\alpha \cdot B_t - \frac{1}{2}|\alpha|^2 t)$ are called **exponential martingales**.

Proof. We only prove the second assertion for $d = 1$ and the right-continuous filtration (\mathcal{F}_t) . The verification of the remaining statements is left as an exercise.

For $d = 1$, since B_t is normally distributed, the \mathcal{F}_t -measurable random variable $B_t^2 - t$ is integrable for any t . Moreover, by Lemma 3.1,

$$\begin{aligned} E[B_t^2 - B_s^2 | \mathcal{F}_s] &= E[(B_t - B_s)^2 | \mathcal{F}_s] + 2B_s \cdot E[B_t - B_s | \mathcal{F}_s] \\ &= E[(B_t - B_s)^2] + 2B_s \cdot E[B_t - B_s] = t - s \end{aligned}$$

almost surely. Hence

$$E[B_t^2 - t | \mathcal{F}_s] = B_s^2 - s \quad P\text{-a.s. for any } 0 \leq s \leq t,$$

i.e., $B_t^2 - t$ is an (\mathcal{F}_t) martingale. \square

Remark (Doob decomposition, variance process of Brownian motion). For a one-dimensional Brownian motion (B_t) , the theorem yields the Doob decomposition

$$B_t^2 = M_t + t$$

of the submartingale (B_t^2) into a martingale (M_t) and the continuous increasing adapted process $\langle B \rangle_t = t$.

A Doob decomposition of the process $f(B_t)$ for general functions $f \in C^2(\mathbb{R})$ will be obtained below as a consequence of Itô's celebrated formula. It states that

$$f(B_t) - f(B_0) = \int_0^t f'(B_s) dB_s + \frac{1}{2} \int_0^t f''(B_s) ds \quad (3.1.2)$$

where the first integral is an Itô stochastic integral, cf. Section 6.3. If, for example, f' is bounded, then the Itô integral is a martingale as a function of t . If f is convex then $f(B_t)$ is a submartingale and the second integral is a continuous increasing adapted process in

t . It is a consequence of (3.1.2) that Brownian motion solves the martingale problem for the operator $\mathcal{L}f = f''/2$ with domain $\text{Dom}(\mathcal{L}) = \{f \in C^2(\mathbb{R}) : f' \text{ bounded}\}$.

Itô's formula (3.1.2) can also be extended to the multi-dimensional case, see Section 6.4 below. The second derivative is then replaced by the Laplacian $\Delta f = \sum_{i=1}^d \frac{\partial^2 f}{\partial x_i^2}$. The multi-dimensional Itô formula implies that a sub- or superharmonic function of d -dimensional Brownian motion is a sub- or supermartingale respectively, if appropriate integrability conditions hold. We now give a direct proof of this fact by the mean value property:

Lemma 3.3 (Mean value property for harmonic function in \mathbb{R}^d). *Suppose that $h \in C^2(\mathbb{R}^d)$ is a (super-)harmonic function, i.e.,*

$$\Delta h(x) \stackrel{(\leq)}{=} 0 \quad \text{for any } x \in \mathbb{R}^d.$$

Then for any $x \in \mathbb{R}^d$ and any rotationally invariant probability measure μ on \mathbb{R}^d ,

$$\int h(x + y) \mu(dy) \stackrel{(\leq)}{=} h(x). \tag{3.1.3}$$

Proof. By the classical mean value property, $h(x)$ is equal to (resp. greater or equal than) the average value $\int_{\partial B_r(x)} h$ of h on any sphere $\partial B_r(x)$ with center at x and radius $r > 0$, cf. e.g. [XXXXKönigsberger: Analysis II]. Moreover, if μ is a rotationally invariant probability measure then the integral in (3.1.3) is an average of average values over spheres:

$$\int h(x + y) \mu(dy) = \int \int_{\partial B_r(x)} h \mu_R(dr) \stackrel{(\leq)}{=} h(x),$$

where μ_R is the distribution of $R(x) = |x|$ under μ . □

Theorem 3.4 (Superharmonic functions of Brownian motion are supermartingales). *If $h \in C^2(\mathbb{R}^d)$ is a (super-) harmonic function then $(h(B_t))$ is a (super-) martingale w.r.t. (\mathcal{F}_t) provided $h(B_t)$ (resp. $h(B_t)^+$) is integrable for any $t \geq 0$.*

Proof. By Lemma 3.1 and the mean value property, we obtain

$$\begin{aligned}
 E[h(B_t) | \mathcal{F}_s](\omega) &= E[h(B_s + B_t - B_s) | \mathcal{F}_s](\omega) \\
 &= E[h(B_s(\omega) + B_t - B_s)] \\
 &= \int h(B_s(\omega) + y) N(0, (t-s)I)(dy) \\
 &\stackrel{(\leq)}{=} h(B_s(\omega))
 \end{aligned}$$

for any $0 \leq s \leq t$ and P -almost every ω . □

3.2 Optional Sampling and Optional Stopping

The Optional Sampling Theorem

The optional stopping theorem can be easily extended to continuous time martingales with continuous sample paths. We directly prove a generalization:

Theorem 3.5 (Optional Sampling Theorem). *Suppose that $(M_t)_{t \in [0, \infty]}$ is a martingale w.r.t. an arbitrary filtration (\mathcal{F}_t) such that $t \mapsto M_t(\omega)$ is continuous for P -almost every ω . Then*

$$E[M_T | \mathcal{F}_S] = M_S \quad P\text{-almost surely} \quad (3.2.1)$$

for any bounded (\mathcal{F}_t) stopping times S and T with $S \leq T$.

We point out that an additional assumption on the filtration (e.g. right-continuity) is not required in the theorem. Stopping times and the σ -algebra \mathcal{F}_S are defined for arbitrary filtrations in complete analogy to the definitions for the filtration (\mathcal{F}_t^B) in Section 1.5.

Remark (Optional Stopping). By taking expectations in the Optional Sampling Theorem, we obtain

$$E[M_T] = E[E[M_T | \mathcal{F}_0]] = E[M_0]$$

for any bounded stopping time T . For unbounded stopping times,

$$E[M_T] = E[M_0]$$

holds by dominated convergence provided $T < \infty$ almost surely, and the random variables $M_{T \wedge n}, n \in \mathbb{N}$, are uniformly integrable.

Proof of Theorem 3.5. We verify the defining properties of the conditional expectation in (3.4) by approximating the stopping times by discrete random variables:

(1). M_S has an \mathcal{F}_S -measurable modification: For $n \in \mathbb{N}$ let $\tilde{S}_n = 2^{-n} \lfloor 2^n S \rfloor$, i.e.,

$$\tilde{S}_n = k \cdot 2^{-n} \quad \text{on} \quad \{k \cdot 2^{-n} \leq S < (k+1)2^{-n}\} \text{ for any } k = 0, 1, 2, \dots$$

We point out that in general, \tilde{S}_n is **not** a stopping time w.r.t. (\mathcal{F}_t) . Clearly, the sequence $(\tilde{S}_n)_{n \in \mathbb{N}}$ is *increasing* with $S = \lim S_n$. By almost sure continuity

$$M_S = \lim_{n \rightarrow \infty} M_{\tilde{S}_n} \quad P\text{-almost surely.} \quad (3.2.2)$$

On the other hand, each of the random variables $M_{\tilde{S}_n}$ is \mathcal{F}_S -measurable. In fact,

$$M_{\tilde{S}_n} \cdot I_{\{S \leq t\}} = \sum_{k: k \cdot 2^{-n} \leq t} M_{k \cdot 2^{-n}} \cdot I_{\{k \cdot 2^{-n} \leq S < (k+1)2^{-n} \text{ and } S \leq t\}}$$

is \mathcal{F}_t -measurable for any $t \geq 0$ since S is an (\mathcal{F}_t) stopping time. Therefore, by (3.2.2), the random variable $\tilde{M}_S := \limsup_{n \rightarrow \infty} M_{\tilde{S}_n}$ is an \mathcal{F}_S -measurable modification of M_S .

(2). $E[M_T; A] = E[M_S; A]$ for any $A \in \mathcal{F}_S$: For $n \in \mathbb{N}$, the discrete random variables $T_n = 2^{-n} \cdot \lceil 2^n T \rceil$ and $S_n = 2^{-n} \cdot \lceil 2^n S \rceil$ are (\mathcal{F}_t) stopping times satisfying $T_n \geq S_n \geq S$, cf. the proof of Theorem 1.26. In particular, $\mathcal{F}_S \subseteq \mathcal{F}_{S_n} \subseteq \mathcal{F}_{T_n}$. Furthermore, (T_n) and (S_n) are *decreasing* sequences with $T = \lim T_n$ and $S = \lim S_n$. As T and S are bounded random variables by assumption, the sequences (T_n) and (S_n) are *uniformly bounded* by a finite constant $c \in (0, \infty)$. Therefore, we obtain




Figure 3.1: Two ways to approximate a continuous stopping time.

$$\begin{aligned}
 E[M_{T_n}; A] &= \sum_{k: k \cdot 2^{-n} \leq c} E[M_{k \cdot 2^{-n}}; A \cap \{T_n = k \cdot 2^{-n}\}] \\
 &= \sum_{k: k \cdot 2^{-n} \leq c} E[M_c; A \cap \{T_n = k \cdot 2^{-n}\}] \quad (3.2.3) \\
 &= E[M_c; A] \quad \text{for any } A \in \mathcal{F}_{T_n},
 \end{aligned}$$

and similarly

$$E[M_{S_n}; A] = E[M_c; A] \quad \text{for any } A \in \mathcal{F}_{S_n}. \quad (3.2.4)$$

In (3.2.3) we have used that (M_t) is an (\mathcal{F}_t) martingale, and $A \cap \{T_n = k \cdot 2^{-n}\} \in \mathcal{F}_{k \cdot 2^{-n}}$. A set $A \in \mathcal{F}_S$ is contained both in \mathcal{F}_{T_n} and \mathcal{F}_{S_n} . Thus by (3.2.3) and (3.2.4),

$$E[M_{T_n}; A] = E[M_{S_n}; A] \quad \text{for any } n \in \mathbb{N} \text{ and any } A \in \mathcal{F}_S. \quad (3.2.5)$$

As $n \rightarrow \infty$, $M_{T_n} \rightarrow M_T$ and $M_{S_n} \rightarrow M_S$ almost surely by continuity. It remains to show that the expectations in (3.2.5) converge as well. To this end note that by (3.2.3) and (3.2.4),

$$M_{T_n} = E[M_c | \mathcal{F}_{T_n}] \quad \text{and} \quad M_{S_n} = E[M_c | \mathcal{F}_{S_n}] \quad P\text{-almost surely.}$$

We will prove in Section 4.3 that any family of conditional expectations of a given random variable w.r.t. different σ -algebras is uniformly integrable, and that for uniformly integrable random variables a generalized Dominated Convergence Theorem holds, cf. Theorem 4.13. Therefore, we finally obtain

$$\begin{aligned} E[M_T; A] &= E[\lim M_{T_n}; A] = \lim E[M_{T_n}; A] \\ &= \lim E[M_{S_n}; A] = E[\lim M_{S_n}; A] = E[M_S; A], \end{aligned}$$

completing the proof of the theorem. □

Remark (Measurability and completion). In general, the random variable M_S is not necessarily \mathcal{F}_S -measurable. However, we have shown in the proof that M_S always has an \mathcal{F}_S -measurable modification \widetilde{M}_S . If the filtration contains all measure zero sets, then this implies that M_S itself is \mathcal{F}_S -measurable and hence a version of $E[M_T | \mathcal{F}_S]$.

Ruin probabilities and passage times revisited

Similarly as for random walks, the Optional Sampling Theorem can be applied to compute distributions of passage times and hitting probabilities for Brownian motion. For a one-dimensional Brownian motion (B_t) starting at 0, and $a, b > 0$, let

$$T = \inf\{t \geq 0 : B_t \notin (-b, a)\} \quad \text{and} \quad T_a = \inf\{t \geq 0 : B_t = a\}$$

denote the first exit time from the interval $(-b, a)$ and the first passage time to the point a , respectively. In Section 1.5 we have computed the distribution of T_a by the reflection principle. This and other results can be recovered by applying optional stopping to the basic martingales of Brownian motion. The advantage of this approach is that it carries over to other diffusion processes.

Exercise (Exit and passage times of Brownian motion). Prove by optional stopping:

- (1). *Law of the exit point:* $P[B_T = a] = b/(a + b)$, $P[B_T = -b] = a/(a + b)$,
- (2). *Mean exit time:* $E[T] = a \cdot b$ and $E[T_a] = \infty$,

(3). *Laplace transform of passage times:* For any $s > 0$,

$$E[\exp(-sT_a)] = \exp(-a\sqrt{2s}).$$

Conclude that the distribution of T_a on $(0, \infty)$ is absolutely continuous with density

$$f_{T_a}(t) = a \cdot (2\pi t^3)^{-1/2} \cdot \exp(-a^2/2t).$$

Exit laws and Dirichlet problem

Applying optional stopping to harmonic functions of a multidimensional Brownian motion yields a generalization of the mean value property and a stochastic representation for solutions of the Dirichlet problem. This will be exploited in full generality in Chapter 7. Here, we only sketch the basic idea.

Suppose that $h \in C^2(\mathbb{R}^d)$ is a harmonic function and that $(B_t)_{t \geq 0}$ is a d -dimensional Brownian motion starting at x w.r.t. the probability measure P_x . Assuming that

$$E_x[h(B_t)] < \infty \quad \text{for any } t \geq 0,$$

the mean value property for harmonic functions implies that $h(B_t)$ is a martingale under P_x , cf. Theorem 3.4. The first hitting time $T = \inf\{t \geq 0 : B_t \in \mathbb{R}^d \setminus D\}$ of the complement of an open set $D \subseteq \mathbb{R}^d$ is a stopping time w.r.t. the filtration (\mathcal{F}_t^B) . Therefore, by Theorem 3.5 and the remark below, we obtain

$$E_x[h(B_{T \wedge n})] = E_x[h(B_0)] = h(x) \quad \text{for any } n \in \mathbb{N}. \quad (3.2.6)$$

Now let us assume in addition that the set D is bounded. Then T is almost surely finite, and the sequence of random variables $h(B_{T \wedge n})$ ($n \in \mathbb{N}$) is uniformly bounded because $B_{T \wedge n}$ takes values in the closure \bar{D} for any $n \in \mathbb{N}$. Applying the Dominated Convergence Theorem to (3.2.6), we obtain the integral representation

$$h(x) = E_x[h(B_T)] = \int_{\partial D} h(y) \mu_x(dy) \quad (3.2.7)$$

where $\mu_x = P_x \circ B_T^{-1}$ denotes the exit law from D for Brownian motion starting at x . In Chapter 7, we show that the representation (3.2.7) still holds true if h is a continuous

function defined on \overline{D} that is C^2 and harmonic on D . The proof requires localization techniques that will be developed below in the context of stochastic calculus. For the moment we note that the representation (3.2.7) has several important aspects and applications:

Generalized mean value property for harmonic functions. For any bounded domain $D \subseteq \mathbb{R}^d$ and any $x \in D$, $h(x)$ is the average of the boundary values of h on ∂D w.r.t. the measure μ_x .

Stochastic representation for solutions of the Dirichlet problem. A solution $h \in C^2(D) \cap C(\overline{D})$ of the Dirichlet problem

$$\begin{aligned} \Delta h(x) &= 0 & \text{for } x \in D, \\ h(x) &= f(x) & \text{for } x \in \partial D, \end{aligned} \quad (3.2.8)$$

has a stochastic representation

$$h(x) = E_x[f(B_T)] \quad \text{for any } x \in \overline{D}. \quad (3.2.9)$$

Monte Carlo solution of the Dirichlet problem. The stochastic representation (3.2.9) can be used as the basis of a Monte Carlo method for computing the harmonic function $h(x)$ approximately by simulating a large number n of sample paths of Brownian motion starting at x , and estimating the expectation by the corresponding empirical average. Although in many cases classical numerical methods are more efficient, the Monte Carlo method is useful in high dimensional cases. Furthermore, it carries over to far more general situations.

Computation of exit law. Conversely, if the Dirichlet problem (3.2.8) has a unique solution h , then computation of h (for example by standard numerical methods) enables us to obtain the expectations in (3.2.8). In particular, the probability $h(x) = P_x[B_T \in A]$ for Brownian motion exiting the domain on a subset $A \subseteq \partial D$ is informally given as the solution of the Dirichlet problem

$$\Delta h = 0 \quad \text{on } D, \quad h = I_A \quad \text{on } \partial D.$$

This can be made rigorous under regularity assumptions. The full exit law is the *harmonic measure*, i.e., the probability measure μ_x such that the representation (3.2.7) holds for any function $h \in C^2(D) \cap C(\overline{D})$ with $\Delta h = 0$ on D . For simple domains such as half-spaces, balls and cylinders, this harmonic measure can be computed explicitly.

Example (Exit laws from balls). For $d \geq 2$, the exit law from the unit ball $D = \{y \in \mathbb{R}^d : |y| < 1\}$ for Brownian motion starting at a point $x \in \mathbb{R}^d$ with $|x| < 1$ is given by

$$\mu_x(dy) = \frac{1 - |x|^2}{|y - x|^d} \nu(dy)$$

where ν denotes the normalized surface measure on the unit sphere $S^{d-1} = \{y \in \mathbb{R}^d : |y| = 1\}$. Indeed, the classical Poisson integral formula states that for any $f \in C(S^{d-1})$, the function

$$h(x) = \int f(y) \mu_x(dy)$$

solves the Dirichlet problem on D with boundary values $\lim_{x \rightarrow z} h(x) = f(z)$ for any $z \in S^{d-1}$, cf. e.g. [XXX Karatzas/Shreve, Ch. 4]. Hence by (3.2.9),

$$E_x[f(B_T)] = \int f(y) \frac{1 - |x|^2}{|y - x|^d} \nu(dy)$$

holds for any $f \in C(S^{d-1})$, and thus by a standard approximation argument, for any indicator function of a measurable subset of S^{d-1} .

3.3 Maximal inequalities and the Law of the Iterated Logarithm

The extension of Doob's maximal inequality to the continuous time case is straightforward. As a first application, we give a proof for the upper bound in the law of the iterated logarithm.

Maximal inequalities in continuous time

Theorem 3.6 (Doob's L^p inequality in continuous time). *Suppose that $(M_t)_{t \in [0, \infty)}$ is a martingale with almost surely right continuous sample paths $t \mapsto M_t(\omega)$. Then the following estimates hold for any $a \in [0, \infty)$, $p \in [1, \infty)$, $q \in (1, \infty]$ with $\frac{1}{p} + \frac{1}{q} = 1$, and $c > 0$:*

$$(1). \quad P \left[\sup_{t \in [0, a]} |M_t| \geq c \right] \leq c^{-p} \cdot E[|M_a|^p],$$

$$(2). \quad \left\| \sup_{t \in [0, a]} |M_t| \right\|_{L^p} \leq q \cdot \|M_a\|_{L^p}.$$

Remark. The same estimates hold for non-negative submartingales.

Proof. Let (π_n) denote an increasing sequence of partitions of the interval $[0, a]$ such that the mesh size of π_n goes to 0 as $n \rightarrow \infty$. By Corollary 2.13 applied to the discrete time martingale $(M_t)_{t \in \pi_n}$, we obtain

$$P \left[\max_{t \in \pi_n} |M_t| \geq c \right] \leq E[|M_a|^p] / c^p \quad \text{for any } n \in \mathbb{N}.$$

Moreover, as $n \rightarrow \infty$,


$$\max_{t \in \pi_n} |M_t| \nearrow \sup_{t \in [0, a]} |M_t| \quad \text{almost surely}$$

by right continuity of the sample paths. Hence

$$\begin{aligned} P \left[\sup_{t \in [0, a]} |M_t| > c \right] &= P \left[\bigcup_n \left\{ \max_{t \in \pi_n} |M_t| > c \right\} \right] \\ &= \lim_{n \rightarrow \infty} P \left[\max_{t \in \pi_n} |M_t| > c \right] \leq E[|M_a|^p] / c^p. \end{aligned}$$

The first assertion now follows by replacing c by $c - \varepsilon$ and letting ε tend to 0. The second assertion follows similarly from Theorem 2.15. \square

As a first application of the maximal inequality to Brownian motion, we derive an upper bound for the probability that the graph of one-dimensional Brownian motion passes a line in \mathbb{R}^2 :



Lemma 3.7 (Passage probabilities for lines). *For a one-dimensional Brownian motion (B_t) starting at 0 we have*

$$P[B_t \geq \beta + \alpha t/2 \text{ for some } t \geq 0] \leq \exp(-\alpha\beta) \quad \text{for any } \alpha, \beta > 0.$$

Proof. Applying the maximal inequality to the exponential martingale

$$M_t^\alpha = \exp(\alpha B_t - \alpha^2 t/2)$$

yields

$$\begin{aligned} P[B_t \geq \beta + \alpha t/2 \text{ for some } t \in [0, a]] &= P \left[\sup_{t \in [0, a]} (B_t - \alpha t/2) \geq \beta \right] \\ &= P \left[\sup_{t \in [0, a]} M_t^\alpha \geq \exp(\alpha\beta) \right] \leq \exp(-\alpha\beta) \cdot E[M_a^\alpha] = \exp(-\alpha\beta) \end{aligned}$$

for any $a > 0$. The assertion follows in the limit as $a \rightarrow \infty$. \square

With slightly more effort, it is possible to compute the passage probability and the distribution of the first passage time of a line explicitly, cf. ?? below.

Application to LIL

A remarkable consequence of Lemma 3.7 is a simplified proof for the upper bound half of the Law of the Iterated Logarithm:

Theorem 3.8 (LIL, upper bound). *For a one-dimensional Brownian motion (B_t) starting at 0,*

$$\limsup_{t \searrow 0} \frac{B_t}{\sqrt{2t \log \log t^{-1}}} \leq +1 \quad P\text{-almost surely.} \quad (3.3.1)$$


Proof. Let $\delta > 0$. We would like to show that almost surely,

$$B_t \leq (1 + \delta)h(t) \quad \text{for sufficiently small } t > 0,$$

where $h(t) := \sqrt{2t \log \log t^{-1}}$. Fix $\theta \in (0, 1)$. The idea is to approximate the function $h(t)$ by affine functions

$$l_n(t) = \beta_n + \alpha_n t/2$$

on each of the intervals $[\theta^n, \theta^{n-1}]$, and to apply the upper bounds for the passage probabilities from the lemma.




We choose α_n and β_n in a such way that $l_n(\theta^n) = h(\theta^n)$ and $l_n(0) = h(\theta^n)/2$, i.e.,

$$\beta_n = h(\theta^n)/2 \quad \text{and} \quad \alpha_n = h(\theta^n)/\theta^n.$$

For this choice we have $l_n(\theta^n) \geq \theta \cdot l_n(\theta^{n-1})$, and hence

$$\begin{aligned} l_n(t) &\leq l_n(\theta^{n-1}) \leq \frac{l_n(\theta^n)}{\theta} \\ &= \frac{h(\theta^n)}{\theta} \leq \frac{h(t)}{\theta} \quad \text{for any } t \in [\theta^n, \theta^{n-1}]. \end{aligned} \tag{3.3.2}$$



We now want to apply the Borel-Cantelli lemma to show that with probability one, $B_t \leq (1 + \delta)l_n(t)$ for large n . By Lemma 3.7,

$$\begin{aligned} P[B_t \geq (1 + \delta)l_n(t) \quad \text{for some } t \geq 0] &\leq \exp(-\alpha_n \beta_n \cdot (1 + \delta)^2) \\ &= \exp\left(-\frac{h(\theta^n)^2}{2\theta^n} \cdot (1 + \delta)^2\right). \end{aligned}$$

Choosing $h(t) = \sqrt{2t \log \log t^{-1}}$, the right hand side is equal to a constant multiple of $n^{-(1+\delta)^2}$, which is a summable sequence. Note that we do not have to know the precise form of $h(t)$ in advance to carry out the proof – we just choose $h(t)$ in such a way that the probabilities become summable!

Now, by Borel-Cantelli, for P -almost every ω there exists $N(\omega) \in \mathbb{N}$ such that

$$B_t(\omega) \leq (1 + \delta)l_n(t) \quad \text{for any } t \in [0, 1] \text{ and } n \geq N(\omega). \quad (3.3.3)$$

By (3.3.2), the right hand side of (3.3.3) is dominated by $(1 + \delta)h(t)/\theta$ for $t \in [\theta^n, \theta^{n-1}]$.

Hence

$$B_t \leq \frac{1 + \delta}{\theta} h(t) \quad \text{for any } t \in \bigcup_{n \geq N} [\theta^n, \theta^{n-1}],$$

i.e., for any $t \in (0, \theta^{N-1})$, and therefore,

$$\limsup_{t \searrow 0} \frac{B_t}{h(t)} \leq \frac{1 + \delta}{\theta} \quad P\text{-almost surely.}$$

The assertion then follows in the limit as $\theta \nearrow 1$ and $\delta \searrow 0$. \square

Since $(-B_t)$ is again a Brownian motion starting at 0, the upper bound (3.3.1) also implies

$$\liminf_{t \searrow 0} \frac{B_t}{\sqrt{2t \log \log t^{-1}}} \geq -1 \quad P\text{-almost surely.} \quad (3.3.4)$$

The converse bounds are actually easier to prove since we can use the independence of the increments and apply the second Borel-Cantelli Lemma. We only mention the key steps and leave the details as an exercise:

Exercise (Complete proof of LIL). Prove the Law of the Iterated Logarithm:

$$\limsup_{t \searrow 0} \frac{B_t}{h(t)} = +1 \quad \text{and} \quad \liminf_{t \searrow 0} \frac{B_t}{h(t)} = -1$$

where $h(t) = \sqrt{2t \log \log t^{-1}}$. Proceed in the following way:

- (1). Let $\theta \in (0, 1)$ and consider the increments $Z_n = B_{\theta^n} - B_{\theta^{n+1}}, n \in \mathbb{N}$. Show that if $\varepsilon > 0$, then

$$P[Z_n > (1 - \varepsilon)h(\theta^n) \text{ infinitely often}] = 1.$$

(Hint: $\int_x^\infty \exp(-z^2/2) dz \leq x^{-1} \exp(-x^2/2)$)

- (2). Conclude that by (3.3.4),

$$\limsup_{t \searrow 0} \frac{B_t}{h(t)} \geq 1 - \varepsilon \quad P\text{-almost surely for any } \varepsilon > 0,$$

and complete the proof of the LIL by deriving the lower bounds

$$\limsup_{t \searrow 0} \frac{B_t}{h(t)} \geq 1 \quad \text{and} \quad \liminf_{t \searrow 0} \frac{B_t}{h(t)} \leq -1 \quad P\text{-almost surely.} \quad (3.3.5)$$

Chapter 4

Martingale Convergence Theorems

The strength of martingale theory is partially due to powerful general convergence theorems that hold for martingales, sub- and supermartingales. In this chapter, we study convergence theorems with different types of convergence including almost sure, L^2 and L^1 convergence, and consider first applications.

At first, we will again focus on discrete-parameter martingales – the results can then be easily extended to continuous martingales.

4.1 Convergence in L^2

Already when proving the Law of Large Numbers, L^2 convergence is much easier to show than, for example, almost sure convergence. The situation is similar for martingales: A necessary and sufficient condition for convergence in the Hilbert space $L^2(\Omega, \mathcal{A}, P)$ can be obtained by elementary methods.

Martingales in L^2

Consider a discrete-parameter martingale $(M_n)_{n \geq 0}$ w.r.t. a filtration (\mathcal{F}_n) on a probability space (Ω, \mathcal{A}, P) . Throughout this section we assume:

Assumption (Square integrability). $E[M_n^2] < \infty$ for any $n \geq 0$.

We start with an important remark:

Lemma 4.1. *The increments $Y_n = M_n - M_{n-1}$ of a square-integrable martingale are centered and orthogonal in $L^2(\Omega, \mathcal{A}, P)$ (i.e. uncorrelated).*

Proof. By definition of a martingale, $E[Y_n | \mathcal{F}_{n-1}] = 0$ for any $n \geq 0$. Hence $E[Y_n] = 0$ and $E[Y_m Y_n] = E[Y_m \cdot E[Y_n | \mathcal{F}_{n-1}]] = 0$ for $0 \leq m < n$. \square

Since the increments are also orthogonal to M_0 by an analogue argument, a square integrable martingale sequence consists of partial sums of a sequence of uncorrelated random variables:

$$M_n = M_0 + \sum_{k=1}^n Y_k \quad \text{for any } n \geq 0.$$

The Convergence Theorem

The central result of this section shows that an L^2 -bounded martingale (M_n) can **always** be extended to $n \in \{0, 1, 2, \dots\} \cup \{\infty\}$:

Theorem 4.2 (L^2 Martingale Convergence Theorem). *The martingale sequence (M_n) converges in $L^2(\Omega, \mathcal{A}, P)$ as $n \rightarrow \infty$ if and only if it is bounded in L^2 in the sense that*

$$\sup_{n \geq 0} E[M_n^2] < \infty. \quad (4.1.1)$$

In this case, the representation

$$M_n = E[M_\infty | \mathcal{F}_n]$$

holds almost surely for any $n \geq 0$, where M_∞ denotes the limit of M_n in $L^2(\Omega, \mathcal{A}, P)$.

We will prove in the next section that (M_n) does also converge almost surely to M_∞ . An analogue result to Theorem 4.2 holds with L^2 replaced by L^p for any $p \in (1, \infty)$ but not for $p = 1$, cf. Section 4.3 below.

Proof. (1). Let us first note that

$$E[(M_n - M_m)^2] = E[M_n^2] - E[M_m^2] \quad \text{for } 0 \leq m \leq n. \quad (4.1.2)$$

Indeed,

$$\begin{aligned} E[M_n^2] - E[M_m^2] &= E[(M_n - M_m)(M_n + M_m)] \\ &= E[(M_n - M_m)^2] + 2E[M_m \cdot (M_n - M_m)], \end{aligned}$$

and the last term vanishes since the increment $M_n - M_m$ is orthogonal to M_m in L^2 .

- (2). To prove that (4.1.1) is sufficient for L^2 convergence, note that the sequence $(E[M_n^2])_{n \geq 0}$ is increasing by (4.1.2). If (4.1.1) holds then this sequence is bounded, and hence a Cauchy sequence. Therefore, by (4.1.2), (M_n) is a Cauchy sequence in L^2 . Convergence now follows by completeness of $L^2(\Omega, \mathcal{A}, P)$.
- (3). Conversely, if (M_n) converges in L^2 to a limit M_∞ , then the L^2 norms are bounded. Moreover, by Jensen's inequality, for each fixed $k \geq 0$,

$$E[M_n | \mathcal{F}_k] \longrightarrow E[M_\infty | \mathcal{F}_k] \quad \text{in } L^2(\Omega, \mathcal{A}, P) \text{ as } n \rightarrow \infty.$$

As (M_n) is a martingale, we have $E[M_n | \mathcal{F}_k] = M_k$ for $n \geq k$, and hence

$$M_k = E[M_\infty | \mathcal{F}_k] \quad P\text{-almost surely.}$$

□

Remark (Functional analytic interpretation of L^2 convergence theorem). The assertion of the L^2 martingale convergence theorem can be rephrased as a purely functional analytic statement:

An infinite sum $\sum_{k=1}^{\infty} Y_k$ of orthogonal vectors Y_k in the Hilbert space $L^2(\Omega, \mathcal{A}, P)$ is convergent if and only if the sequence of partial sums $\sum_{k=1}^n Y_k$ is bounded.

How can boundedness in L^2 be verified for martingales? Writing the martingale (M_n) as the sequence of partial sums of its increments $Y_n = M_n - M_{n-1}$, we have

$$E[M_n^2] = \left(M_0 + \sum_{k=1}^n Y_k, M_0 + \sum_{k=1}^n Y_k \right)_{L^2} = E[M_0^2] + \sum_{k=1}^n E[Y_k^2]$$

by orthogonality of the increments and M_0 . Hence

$$\sup_{n \geq 0} E[M_n^2] = E[M_0^2] + \sum_{k=1}^{\infty} E[Y_k^2].$$

Alternatively, we have $E[M_n^2] = E[M_0^2] + E[\langle M \rangle_n]$. Hence by monotone convergence

$$\sup_{n \geq 0} E[M_n^2] = E[M_0^2] + E[\langle M \rangle_{\infty}]$$

where $\langle M \rangle_{\infty} = \sup \langle M \rangle_n$.

Summability of sequences with random signs

As a first application we study the convergence of series with coefficients with random signs. In an introductory analysis course it is shown as an application of the integral and Leibniz criterion for convergence of series that

$$\begin{aligned} \sum_{n=1}^{\infty} n^{-\alpha} \text{ converges} &\iff \alpha > 1, \text{ whereas} \\ \sum_{n=1}^{\infty} (-1)^n n^{-\alpha} \text{ converges} &\iff \alpha > 0. \end{aligned}$$

Therefore, it seems interesting to see what happens if the signs are chosen randomly. The L^2 martingale convergence theorem yields:

Corollary 4.3. *Let (a_n) be a real sequence. If (ε_n) is a sequence of independent random variables on (Ω, \mathcal{A}, P) with $P[\varepsilon_n = +1] = P[\varepsilon_n = -1] = 1/2$, then*

$$\sum_{n=1}^{\infty} \varepsilon_n a_n \text{ converges in } L^2(\Omega, \mathcal{A}, P) \iff \sum_{n=1}^{\infty} a_n^2 < \infty.$$

Proof. The sequence $M_n = \sum_{k=1}^n \varepsilon_k a_k$ of partial sums is a martingale with

$$\sup_{n \geq 0} E[M_n^2] = \sum_{k=1}^{\infty} E[\varepsilon_k^2 a_k^2] = \sum_{k=1}^{\infty} a_k^2.$$

□

Example. The series $\sum_{n=1}^{\infty} \varepsilon_n \cdot n^{-\alpha}$ converges in L^2 if and only if $\alpha > \frac{1}{2}$.

Remark (Almost sure asymptotics). By the Supermartingale Convergence Theorem (cf. Theorem 4.5 below), the series $\sum \varepsilon_n a_n$ also converges almost surely if $\sum a_n^2 < \infty$. On the other hand, if $\sum a_n^2 = \infty$ then the series of partial sums has almost surely unbounded oscillations:

Exercise. Suppose that $\sum a_n = \infty$, and let $M_n = \sum_{k=1}^n \varepsilon_k a_k$.

- (1). Compute the conditional variance process $\langle M \rangle_n$.
- (2). For $c > 0$ let $T_c = \inf\{n \geq 0 : |M_n| \geq c\}$. Apply the Optional Stopping Theorem to the martingale in the Doob decomposition of (M_n^2) , and conclude that $P[T_c = \infty] = 0$.
- (3). Prove that (M_n) has almost surely unbounded oscillations.

L^2 convergence in continuous time

The L^2 convergence theorem directly extends to the continuous-parameter case.

Theorem 4.4 (L^2 Martingale Convergence Theorem in continuous time). Let $a \in (0, \infty]$. If $(M_t)_{t \in [0, a]}$ is a martingale w.r.t. a filtration $(\mathcal{F}_t)_{t \in [0, a]}$ such that

$$\sup_{t \in [0, u]} E[M_t^2] < \infty$$

then $M_u = \lim_{t \nearrow u} M_t$ exists in $L^2(\Omega, \mathcal{A}, P)$ and $(M_t)_{t \in [0, u]}$ is again a square-integrable martingale.

Proof. Choose any increasing sequence $t_n \in [0, u)$ such that $t_n \rightarrow u$. Then (M_{t_n}) is an L^2 -bounded discrete-parameter martingale. Hence the limit $M_u = \lim M_{t_n}$ exists in L^2 , and

$$M_{t_n} = E[M_u | \mathcal{F}_{t_n}] \quad \text{for any } n \in \mathbb{N}. \quad (4.1.3)$$

For an arbitrary $t \in [0, u)$, there exists $n \in \mathbb{N}$ with $t_n \in (t, u)$. Hence

$$M_t = E[M_{t_n} | \mathcal{F}_t] = E[M_u | \mathcal{F}_t]$$

by (4.1.3) and the tower property. In particular, $(M_t)_{t \in [0, u]}$ is a square-integrable martingale. By orthogonality of the increments,

$$E[(M_u - M_{t_n})^2] = E[(M_u - M_t)^2] + E[(M_t - M_{t_n})^2] \geq E[(M_u - M_t)^2]$$

whenever $t_n \leq t \leq u$. Since $M_{t_n} \rightarrow M_u$ in L^2 , we obtain

$$\lim_{t \nearrow u} E[(M_u - M_t)^2] = 0.$$

□

Remark. (1). Note that in the proof it is enough to consider a fixed sequence $t_n \nearrow u$.

(2). To obtain almost sure convergence, an additional regularity condition on the sample paths (e.g. right-continuity) is required, cf. below. This assumption is not needed for L^2 convergence.

4.2 Almost sure convergence of supermartingales

Let $(Z_n)_{n \geq 0}$ be a discrete-parameter supermartingale w.r.t. a filtration $(\mathcal{F}_n)_{n \geq 0}$ on a probability space (Ω, \mathcal{A}, P) . The following theorem yields a stochastic counterpart to the fact that any lower bounded decreasing sequence of reals converges to a finite limit:

Theorem 4.5 (Supermartingale Convergence Theorem, Doob). *If $\sup_{n \geq 0} E[Z_n^-] < \infty$ then (Z_n) converges almost surely to a random variable $Z_\infty \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$.*

In particular, supermartingales that are uniformly bounded from below converge almost surely to an integrable random variable.

Lemma 4.6 (Doob). *If (Z_n) is a supermartingale then*

$$(b - a) \cdot E[U_n^{(a,b)}] \leq E[(Z_n - a)^-] \quad \text{for any } a < b \text{ and } n \geq 0.$$

Proof. We may assume $E[Z_n^-] < \infty$ since otherwise there is nothing to prove. The key idea is to set up a predictable gambling strategy that increases our capital by $(b - a)$ for each completed upcrossing. Since the net gain with this strategy should again be a supermartingale this yields an upper bound for the average number of upcrossings. Here is the strategy:

- repeat $\left\{ \begin{array}{l} \bullet \text{ Wait until } Z_k \leq a. \\ \bullet \text{ Then play unit stakes until } Z_k \geq b. \\ \bullet \end{array} \right.$

The stake C_k in round k is

$$C_1 = \begin{cases} 1 & \text{if } Z_0 \leq a, \\ 0 & \text{otherwise,} \end{cases}$$

and for $k \geq 2$,

$$C_k = \begin{cases} 1 & \text{if } (C_{k-1} = 1 \text{ and } Z_{k-1} \leq b) \text{ or } (C_{k-1} = 0 \text{ and } Z_{k-1} \leq a), \\ 0 & \text{otherwise} \end{cases}.$$

Clearly, (C_k) is a predictable, bounded and non-negative sequence of random variables. Moreover, $C_k \cdot (Z_k - Z_{k-1})$ is integrable for any $k \leq n$, because C_k is bounded and

$$E[|Z_k|] = 2E[Z_k^+] - E[Z_k] \leq 2E[Z_k^+] - E[Z_n] \leq 2E[Z_k^+] - E[Z_n^-]$$

for $k \leq n$. Therefore, by Theorem 2.6 and the remark below, the process

$$(C \bullet Z)_k = \sum_{i=1}^k C_i \cdot (Z_i - Z_{i-1}), \quad 0 \leq k \leq n,$$


is again a supermartingale.

Clearly, the value of the process $C \bullet Z$ increases by at least $(b - a)$ units during each

completed upcrossing. Between upcrossing periods, the value of $(C \bullet Z)_k$ is constant. Finally, if the final time n is contained in an upcrossing period, then the process can decrease by at most $(Z_n - a)^-$ units during that last period (since Z_k might decrease before the next upcrossing is completed). Therefore, we have

$$(C \bullet Z)_n \geq (b - a) \cdot U_n^{(a,b)} - (Z_n - a)^-, \quad \text{i.e.,}$$

$$(b - a) \cdot U_n^{(a,b)} \leq (C \bullet Z)_n + (Z_n - a)^-.$$



Since $C \bullet Z$ is a supermartingale with initial value 0, we obtain the upper bound

$$(b - a)E[U_n^{(a,b)}] \leq E[(C \bullet Z)_n] + E[(Z_n - a)^-] \leq E[(Z_n - a)^-].$$

□

Proof of Doob's Convergence Theorem

We can now complete the proof of Theorem 4.5.

Proof. Let

$$U^{(a,b)} = \sup_{n \in \mathbb{N}} U_n^{(a,b)}$$

denote the total number of upcrossings of the supermartingale (Z_n) over an interval (a, b) with $-\infty < a < b < \infty$. By the upcrossing inequality and monotone convergence,

$$E[U^{(a,b)}] = \lim_{n \rightarrow \infty} E[U_n^{(a,b)}] \leq \frac{1}{b - a} \cdot \sup_{n \in \mathbb{N}} E[(Z_n - a)^-]. \quad (4.2.1)$$

Assuming $\sup E[Z_n^-] < \infty$, the right hand side of (4.2.1) is finite since $(Z_n - a)^- \leq |a| + Z_n^-$. Therefore,

$$U^{(a,b)} < \infty \quad P\text{-almost surely,}$$

and hence the event

$$\{\liminf Z_n \neq \limsup Z_n\} = \bigcup_{\substack{a,b \in \mathbb{Q} \\ a < b}} \{U^{(a,b)} = \infty\}$$

has probability zero. This proves almost sure convergence.

It remains to show that the almost sure limit $Z_\infty = \lim Z_n$ is an integrable random variable (in particular, it is finite almost surely). This holds true as, by the remark below Theorem 4.5, $\sup E[Z_n^-] < \infty$ implies that (Z_n) is bounded in L^1 , and therefore

$$E[|Z_\infty|] = E[\liminf |Z_n|] \leq \liminf E[|Z_n|] < \infty$$

by Fatou's lemma. □

Examples and first applications

We now consider a few prototypic applications of the almost sure convergence theorem:

Example (1. Sums of i.i.d. random variables). Consider a Random Walk

$$S_n = \sum_{i=1}^n \eta_i$$

on \mathbb{R} with centered and bounded increments

$$\eta_i \text{ i.i.d. with } |\eta_i| \leq c \text{ and } E[\eta_i] = 0, \quad c \in \mathbb{R}.$$

Suppose that $P[\eta_i \neq 0] > 0$. Then there exists $\varepsilon > 0$ such that $P[|\eta_i| \geq \varepsilon] > 0$. As the increments are i.i.d., the event $\{|\eta_i| \geq \varepsilon\}$ occurs infinitely often with probability one. Therefore, almost surely, the martingale (S_n) does not converge as $n \rightarrow \infty$.

Now let $a \in \mathbb{R}$. We consider the first hitting time

$$T_a = \inf\{t \geq 0 : S_n \geq a\}$$

of the interval $[a, \infty)$. By the Optional Stopping Theorem, the stopped Random Walk $(S_{T_a \wedge n})_{n \geq 0}$ is again a martingale. Moreover, as $S_k < a$ for any $k < T_a$ and the increments are bounded by c , we obtain the upper bound

$$S_{T_a \wedge n} < a + c \quad \text{for any } n \in \mathbb{N}.$$

Therefore, the stopped Random Walk converges almost surely by the Supermartingale Convergence Theorem. As (S_n) does not converge, we can conclude that

$$P[T_a < \infty] = 1 \quad \text{for any } a > 0, \text{ i.e., } \limsup S_n = \infty \quad \text{almost surely.}$$

Since (S_n) is also a submartingale, we obtain

$$\liminf S_n = -\infty \quad \text{almost surely}$$

by an analogue argument. A generalization of this result is given in Theorem 4.7 below.

Remark (Almost sure vs. L^p convergence). In the last example, the stopped process does not converge in L^p for any $p \in [1, \infty)$. In fact,

$$\lim_{n \rightarrow \infty} E[S_{T_a \wedge n}] = E[S_{T_a}] \geq a \quad \text{whereas} \quad E[S_{T_a \wedge n}] = E[S_0] = 0 \text{ for all } n.$$

Example (2. Products of non-negative i.i.d. random variables). Consider a growth process

$$Z_n = \prod_{i=1}^n Y_i$$

with i.i.d. factors $Y_i \geq 0$ with finite expectation $\alpha \in (0, \infty)$. Then

$$M_n = Z_n / \alpha^n$$

is a martingale. By the almost sure convergence theorem, a finite limit M_∞ exists almost surely, because $M_n \geq 0$ for all n . For the almost sure asymptotics of (Z_n) , we distinguish three different cases:

(1). $\alpha < 1$: In this case,

$$Z_n = M_n \cdot \alpha^n$$

converges to 0 exponentially fast with probability one.

(2). $\alpha = 1$: Here (Z_n) is a martingale and converges almost surely to a finite limit. If $P[Y_i \neq 1] > 0$ then there exists $\varepsilon > 0$ such that $Y_i \geq 1 + \varepsilon$ infinitely often with probability one. This is consistent with convergence of (Z_n) only if the limit is zero. Hence, if (Z_n) is not almost surely constant, then also in the critical case, $Z_n \rightarrow 0$ almost surely.

(3). $\alpha > 1$ (*supercritical*): In this case, on the set $\{M_\infty > 0\}$,

$$Z_n = M_n \cdot \alpha^n \sim M_\infty \cdot \alpha^n,$$

i.e., (Z_n) grows exponentially fast. The asymptotics on the set $\{M_\infty = 0\}$ is not evident and requires separate considerations depending on the model.

Although most of the conclusions in the last example could have been obtained without martingale methods (e.g. by taking logarithms), the martingale approach has the advantage of carrying over to far more general model classes. These include for example branching processes or exponentials of continuous time processes.

Example (3. Boundary behaviors of harmonic functions). Let $D \subseteq \mathbb{R}^d$ be a bounded open domain, and let $h : D \rightarrow \mathbb{R}$ be a harmonic function on D that is bounded from below:


$$\Delta h(x) = 0 \quad \text{for any } x \in D, \quad \inf_{x \in D} h(x) > -\infty. \quad (4.2.2)$$

To study the asymptotic behavior of $h(x)$ as x approaches the boundary ∂D , we construct a Markov chain (X_n) such that $h(X_n)$ is a martingale: Let $r : D \rightarrow (0, \infty)$ be a continuous function such that

$$0 < r(x) < \text{dist}(x, \partial D) \quad \text{for any } x \in D, \quad (4.2.3)$$

and let (X_n) w.r.t P_x denote the canonical time-homogeneous Markov chain with state space D , initial value x , and transition probabilities

$$p(x, dy) = \text{Uniform distribution on } \{y \in \mathbb{R}^d : |y - x| = r(x)\}.$$



By (4.2.3), the function h is integrable w.r.t. $p(x, dy)$, and, by the mean value property,

$$(ph)(x) = h(x) \quad \text{for any } x \in D.$$

Therefore, the process $h(X_n)$ is a martingale w.r.t. P_x for each $x \in D$. As $h(X_n)$ is lower bounded by (4.2.2), the limit as $n \rightarrow \infty$ exists P_x -almost surely by the Supermartingale Convergence Theorem. In particular, since the coordinate functions $x \mapsto x_i$ are also harmonic and lower bounded on \overline{D} , the limit $X_\infty = \lim_{n \rightarrow \infty} X_n$ exists P_x -almost surely. Moreover, X_∞ is in ∂D , because r is bounded from below by a strictly positive constant on any compact subset of D .

Summarizing we have shown:

- (1). *Boundary regularity*: If h is harmonic and bounded from below on D then the limit $\lim_{n \rightarrow \infty} h(X_n)$ exists along almost every trajectory X_n to the boundary ∂D .
- (2). *Representation of h in terms of boundary values*: If h is continuous on \overline{D} , then $h(X_n) \rightarrow h(X_\infty)$ P_x -almost surely and hence

$$h(x) = \lim_{n \rightarrow \infty} E_x[h(X_n)] = E[h(X_\infty)],$$

i.e., the law of X_∞ w.r.t. P_x is the harmonic measure on ∂D .

Note that, in contrast to classical results from analysis, the first statement holds without any smoothness condition on the boundary ∂D . Thus, although boundary values of h may not exist in the classical sense, they do exist along almost every trajectory of the Markov chain!

Generalized Borel-Cantelli Lemma

Another application of the almost sure convergence theorem is a generalization of the Borel-Cantelli lemmas. We first prove a dichotomy for the asymptotic behavior of martingales with L^1 -bounded increments:

Theorem 4.7 (Asymptotics of martingales with L^1 bounded increments). *Suppose that (M_n) is a martingale, and there exists an integrable random variable Y such that*

$$|M_n - M_{n-1}| \leq Y \quad \text{for any } n \in \mathbb{N}.$$

Then for P -almost every ω , the following dichotomy holds:

Either: *The limit $\lim_{n \rightarrow \infty} M_n(\omega)$ exists in \mathbb{R} ,*

or: $\limsup_{n \rightarrow \infty} M_n(\omega) = +\infty$ *and* $\liminf_{n \rightarrow \infty} M_n(\omega) = -\infty$.

The theorem and its proof are a generalization of Example 1 above.

Proof. For $a \in (-\infty, 0)$ let $T_a = \min\{n \geq 0 : M_n \geq a\}$. By the Optional Stopping Theorem, $(M_{T_a \wedge n})$ is a martingale. Moreover,

$$M_{T_a \wedge n} \geq \min(M_0, a - Y) \quad \text{for any } n \geq 0,$$

and the right hand side is an integrable random variable. Therefore, (M_n) converges almost surely on $\{T_a = \infty\}$. Since this holds for every $a < 0$, we obtain almost sure convergence on the set

$$\{\liminf M_n > -\infty\} = \bigcup_{\substack{a < 0 \\ a \in \mathbb{Q}}} \{T_a = \infty\}.$$

Similarly, almost sure convergence follows on the set $\{\limsup M_n < \infty\}$. □

Now let $(\mathcal{F}_n)_{n \geq 0}$ be an arbitrary filtration. As a consequence of Theorem 4.7 we obtain:

Corollary 4.8 (Generalized Borel-Cantelli Lemma). *If (A_n) is a sequence of events with $A_n \in \mathcal{F}_n$ for any n , then the equivalence*

$$\omega \in A_n \text{ infinitely often} \iff \sum_{n=1}^{\infty} P[A_n | \mathcal{F}_{n-1}](\omega) = \infty$$

holds for almost every $\omega \in \Omega$.

Proof. Let $S_n = \sum_{k=1}^n I_{A_k}$ and $T_n = \sum_{k=1}^n E[I_{A_k} | \mathcal{F}_{k-1}]$. Then S_n and T_n are almost surely increasing sequences. Let $S_\infty = \sup S_n$ and $T_\infty = \sup T_n$ denote the limits on $[0, \infty]$. The claim is that almost surely,

$$S_\infty = \infty \iff T_\infty = \infty. \quad (4.2.4)$$

To prove (4.2.4) we note that $S_n - T_n$ is a martingale with bounded increments. Therefore, almost surely, $S_n - T_n$ converges to a finite limit, or $(\limsup(S_n - T_n) = \infty$ and $\liminf(S_n - T_n) = -\infty)$. In the first case, (4.2.4) holds. In the second case, $S_\infty = \infty$ and $T_\infty = \infty$, so (4.2.4) holds, too. \square

The assertion of Corollary 4.8 generalizes both classical Borel-Cantelli Lemmas: If (A_n) is an arbitrary sequence of events in a probability space (Ω, \mathcal{A}, P) then we can consider the filtration $\mathcal{F}_n = \sigma(A_1, \dots, A_n)$. By Corollary 4.8 we obtain:

1st Borel-Cantelli Lemma: If $\sum P[A_n] < \infty$ then $\sum P[A_n | \mathcal{F}_{n-1}] < \infty$ almost surely, and therefore

$$P[A_n \text{ infinitely often}] = 0.$$

2nd Borel-Cantelli Lemma: If $\sum P[A_n] = \infty$ and the A_n are independent then

$$\sum P[A_n | \mathcal{F}_{n-1}] = \sum P[A_n] = \infty \text{ almost surely, and therefore}$$

$$P[A_n \text{ infinitely often}] = 1.$$

Upcrossing inequality and convergence theorem in continuous time

The upcrossing inequality and the supermartingale convergence theorem carry over immediately to the continuous time case if we assume right continuity (or left continuity) of the sample paths. Let $u \in (0, \infty]$, and let $(Z_s)_{s \in [0, u]}$ be a supermartingale in continuous time w.r.t. a filtration (\mathcal{F}_s) . We define the number of upcrossings of (Z_s) over an interval (a, b) before time t as the supremum of the number of upcrossings over all time discretizations $(Z_s)_{s \in \pi}$ where π is a partition of the interval $[0, t]$:

$$U_t^{(a,b)}[Z] := \sup_{\substack{\pi \subset [0, t] \\ \text{finite}}} U^{(a,b)}[(Z_s)_{s \in \pi}].$$

Note that if (Z_s) has right-continuous sample paths and (π_n) is a sequence of partitions of $[0, t]$ such that $0, t \in \pi_0$, $\pi_n \subset \pi_{n+1}$ and $\text{mesh}(\pi_n) \rightarrow 0$ then

$$U_t^{(a,b)}[Z] = \lim_{n \rightarrow \infty} U^{(a,b)}[(Z_s)_{s \in \pi_n}].$$

Theorem 4.9 (Supermartingale Convergence Theorem in continuous time). *Suppose that $(Z_s)_{s \in [0, u]}$ is a right continuous supermartingale.*

(1). *Upcrossing inequality: For any $t \in [0, u)$ and $a < b$,*

$$E[U_t^{(a,b)}] \leq \frac{1}{b-a} E[(Z_t - a)^-].$$

(2). *Convergence Theorem: If $\sup_{s \in [0, u)} E[Z_s^-] < \infty$, then the limit $Z_{u-} = \lim_{s \nearrow u} Z_s$ exists almost surely, and Z_{u-} is an integrable random variable.*

Proof. (1). By the upcrossing inequality in discrete time,

$$E[U^{(a,b)}[(Z_s)_{s \in \pi_n}]] \leq E[(Z_t - a)^-] \quad \text{for any } n \in \mathbb{N},$$

where (π_n) is a sequence of partitions as above. The assertion now follows by the Monotone Convergence Theorem.

- (2). The almost sure convergence can now be proven in the same way as in the discrete time case.

□

More generally than stated above, the upcrossing inequality also implies that for a right-continuous supermartingale $(Z_s)_{s \in [0, u]}$ all the left limits $\lim_{s \nearrow t} Z_s, t \in [0, u)$, exist *simultaneously* with probability one. Thus almost every sample path is *càdlàg* (continue à droite, limites à gauche, i.e., right continuous with left limits). By similar arguments, the existence of a modification with right continuous (and hence *càdlàg*) sample paths can be proven for *any* supermartingale (Z_s) provided the filtration is right continuous and complete, and $s \mapsto E[Z_s]$ is right continuous, cf. e.g. [XXXRevuz/Yor, Ch.II,§2].

4.3 Uniform integrability and L^1 convergence

The Supermartingale Convergence Theorem shows that every supermartingale (Z_n) that is bounded in L^1 converges almost surely to an integrable limit Z_∞ . However, L^1 convergence does not necessarily hold:

Example. (1). Suppose that $Z_n = \prod_{i=1}^n Y_i$ where the Y_i are i.i.d. with $E[Y_i] = 1$, $P[Y_i \neq 1] > 0$. Then, $Z_n \rightarrow 0$ almost surely, cf. Example 2 in Section 4.2. On the other hand, L^1 convergence does not hold as $E[Z_n] = 1$ for any n .

- (2). Similarly, the exponential martingale $M_t = \exp(B_t - t/2)$ of a Brownian motion converges to 0 almost surely, but $E[M_t] = 1$ for any t .

L^1 convergence of martingales is of interest because it implies that a martingale sequence (M_n) can be extended to $n = \infty$, and the random variables M_n are given as conditional expectations of the limit M_∞ . Therefore, we now prove a generalization of the Dominated Convergence Theorem that leads to a necessary and sufficient condition for L^1 convergence.

Uniform integrability

Let (Ω, \mathcal{A}, P) be a probability space. The key condition required to deduce L^1 convergence from convergence in probability is uniform integrability. To motivate the definition we first recall two characterizations of integrable random variables:

Lemma 4.10. *If $X : \Omega \rightarrow \mathbb{R}$ is an integrable random variable on (Ω, \mathcal{A}, P) , then*

$$(1). \lim_{c \rightarrow \infty} E[|X|; |X| \geq c] = 0, \quad \text{and}$$

(2). *for any $\varepsilon > 0$ there exists $\delta > 0$ such that*

$$E[|X|; A] < \varepsilon \quad \text{for any } A \in \mathcal{A} \text{ with } P[A] < \delta.$$

The second statement says that the positive measure

$$Q(A) = E[|X|; A], \quad A \in \mathcal{A},$$

with relative density $|X|$ w.r.t. P is **absolutely continuous** w.r.t. P in the following sense: *For any $\varepsilon > 0$ there exists $\delta > 0$ such that*

$$P[A] < \delta \quad \Rightarrow \quad Q(A) < \varepsilon.$$

Proof. (1). For an integrable random variable X the first assertion holds by the Monotone Convergence Theorem, since $|X| \cdot I_{\{|X| \geq c\}} \searrow 0$ as $c \nearrow \infty$.

(2). Let $\varepsilon > 0$. By (1),

$$\begin{aligned} E[|X|; A] &= E[|X|; A \cap \{|X| \geq c\}] + E[|X|; A \cap \{|X| < c\}] \\ &\leq E[|X|; |X| \geq c] + c \cdot P[A] \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \end{aligned}$$

provided $c \in (0, \infty)$ is chosen appropriately and $P[A] < \varepsilon/2c$.

□

Uniform integrability means that properties (1) and (2) hold uniformly for a family of random variables:

Definition (Uniform integrability). A family $\{X_i : i \in I\}$ of random variables on (Ω, \mathcal{A}, P) is called **uniformly integrable** if and only if

$$\sup_{i \in I} E[|X_i|; |X_i| \geq c] \longrightarrow 0 \quad \text{as } c \rightarrow \infty.$$

Exercise (Equivalent characterization of uniform integrability). Prove that $\{X_i : i \in I\}$ is uniformly integrable if and only if $\sup E[|X_i|; A] < \infty$, and the measures $Q_i(A) = E[|X_i|; A]$ are **uniformly absolutely continuous**, i.e., for any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$P[A] < \delta \quad \Rightarrow \quad \sup_{i \in I} E[|X_i|; A] < \varepsilon.$$

We will prove below that convergence in probability plus uniform integrability is equivalent to L^1 convergence. Before, we state two lemmas giving sufficient conditions for uniform integrability (and hence for L^1 convergence) that can often be verified in applications:

Lemma 4.11 (Sufficient conditions for uniform integrability). A family $\{X_i : i \in I\}$ of random variables is uniformly integrable if one of the following conditions holds:

(1). There exists an integrable random variable Y such that

$$|X_i| \leq Y \quad \text{for any } i \in I.$$

(2). There exists a measurable function $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that

$$\lim_{x \rightarrow \infty} \frac{g(x)}{x} = \infty \quad \text{and} \quad \sup_{i \in I} E[g(|X_i|)] < \infty.$$

Proof. (1). If $|X_i| \leq Y$ then

$$\sup_{i \in I} E[|X_i|; |X_i| \geq c] \leq E[Y; Y \geq c].$$

The right hand side converges to 0 as $c \rightarrow \infty$ if Y is integrable.

(2). The second condition implies uniform integrability, because

$$\sup_{i \in I} E[|X_i|; |X_i| \geq c] \leq \sup_{y \geq c} \frac{y}{g(y)} \cdot \sup_{i \in I} E[g(|X_i|)].$$

□

The first condition in Lemma 4.11 is the classical assumption in the Dominated Convergence Theorem. The second condition holds in particular if

$$\sup_{i \in I} E[|X_i|^p] < \infty \quad \text{for some } p > 1 \quad (\mathbf{L^p \text{ boundedness}}),$$

or, if

$$\sup_{i \in I} E[|X_i|(\log |X_i|)^+] < \infty \quad (\mathbf{Entropy \text{ condition}})$$

is satisfied. Boundedness in L^1 , however, does not imply uniform integrability, cf. the examples at the beginning of this section.

The next observation is crucial for the application of uniform integrability to martingales:

Lemma 4.12 (Conditional expectations are uniformly integrable). *If X is an integrable random variable on (Ω, \mathcal{A}, P) then the family*

$$\{E[X | \mathcal{F}] : \mathcal{F} \subseteq \mathcal{A} \text{ } \sigma\text{-algebra}\}$$

of all conditional expectations of X given sub- σ -algebras of \mathcal{A} is uniformly integrable.

Proof. By Lemma 4.10, for any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\begin{aligned} E[|E[X | \mathcal{F}]|; |E[X | \mathcal{F}]| \geq c] &\leq E[E[|X| | \mathcal{F}]; |E[X | \mathcal{F}]| \geq c] \quad (4.3.1) \\ &= E[|X|; |E[X | \mathcal{F}]| \geq c] < \varepsilon \end{aligned}$$

holds for $c > 0$ with $P[|E[X | \mathcal{F}]| \geq c] < \delta$. Since

$$P[|E[X | \mathcal{F}]| \geq c] \leq \frac{1}{c} E[|E[X | \mathcal{F}]|] \leq \frac{1}{c} E[|X|],$$

(4.3.1) holds simultaneously for all σ -algebras $\mathcal{F} \subseteq \mathcal{A}$ if c is sufficiently large. □

Definitive version of Lebesgue's Dominated Convergence Theorem

Theorem 4.13. *Suppose that $(X_n)_{n \in \mathbb{N}}$ is a sequence of integrable random variables. Then (X_n) converges to a random variable X w.r.t. the L^1 norm if and only if X_n converges to X in probability and the family $\{X_n : n \in \mathbb{N}\}$ is uniformly integrable.*


Proof. (1). We first prove the “if” part of the assertion under the additional assumption that the random variables $|X_n|$ are uniformly bounded by a finite constant c : For $\varepsilon > 0$,

$$\begin{aligned} E[|X_n - X|] &= E[|X_n - X|; |X_n - X| > \varepsilon] + E[|X_n - X|; |X_n - X| \leq \varepsilon] \\ &\leq 2c \cdot P[|X_n - X| > \varepsilon] + \varepsilon. \end{aligned} \quad (4.3.2)$$

Here we have used that $|X_n| \leq c$ and hence $|X| \leq c$ with probability one, because a subsequence of (X_n) converges almost surely to X . For sufficiently large n , the right hand side of (4.3.2) is smaller than 2ε . Therefore, $E[|X_n - X|] \rightarrow 0$ as $n \rightarrow \infty$.

(2). To prove the “if” part under the uniform integrability condition, we consider the cut-off-functions

$$\phi_c(x) = (x \wedge c) \vee (-c)$$



For $c \in (0, \infty)$, the function $\phi_c : \mathbb{R} \rightarrow \mathbb{R}$ is a contraction. Therefore,

$$|\phi_c(X_n) - \phi_c(X)| \leq |X_n - X| \quad \text{for any } n \in \mathbb{N}.$$

If $X_n \rightarrow X$ in probability then $\phi_c(X_n) \rightarrow \phi_c(X)$ in probability. Hence by (1),

$$E[|\phi_c(X_n) - \phi_c(X)|] \rightarrow 0 \quad \text{for any } c > 0. \quad (4.3.3)$$

We would like to conclude that $E[|X_n - X|] \rightarrow 0$ as well. Since (X_n) is uniformly integrable, and a subsequence converges to X almost surely, we have $E[|X|] \leq \liminf E[|X_n|] < \infty$ by Fatou's Lemma. We now estimate

$$\begin{aligned} E[|X_n - X|] &\leq E[|X_n - \phi_c(X_n)|] + E[|\phi_c(X_n) - \phi_c(X)|] + E[|\phi_c(X) - X|] \\ &\leq E[|X_n|; |X_n| \geq c] + E[|\phi_c(X_n) - \phi_c(X)|] + E[|X|; |X| \geq c]. \end{aligned}$$

Let $\varepsilon > 0$ be given. Choosing c large enough, the first and the last summand on the right hand side are smaller than $\varepsilon/3$ for all n by uniform integrability of $\{X_n : n \in \mathbb{N}\}$ and integrability of X . Moreover, by (4.3.3), there exists $n_0(c)$ such that the middle term is smaller than $\varepsilon/3$ for $n \geq n_0(c)$. Hence $E[|X_n - X|] < \varepsilon$ for $n \geq n_0$, and thus $X_n \rightarrow X$ in L^1 .

(3). Now suppose conversely that $X_n \rightarrow X$ in L^1 . Then $X_n \rightarrow X$ in probability by Markov's inequality. To prove uniform integrability, we observe that

$$E[|X_n|; A] \leq E[|X|; A] + E[|X - X_n|] \quad \text{for any } n \in \mathbb{N} \text{ and } A \in \mathcal{A}.$$

For $\varepsilon > 0$, there exist $n_0 \in \mathbb{N}$ and $\delta > 0$ such that

$$\begin{aligned} E[|X - X_n|] &< \varepsilon/2 \quad \text{for any } n > n_0, \text{ and} \\ E[|X|; A] &< \varepsilon/2 \quad \text{whenever } P[A] < \delta, \end{aligned}$$

cf. Lemma 4.10. Hence, if $P[A] < \delta$ then $\sup_{n \geq n_0} E[|X_n|; A] < \varepsilon$.

Moreover, again by Lemma 4.10, there exist $\delta_1, \dots, \delta_{n_0} > 0$ such that for $n \leq n_0$,

$$E[|X_n|; A] < \varepsilon \quad \text{if } P[A] < \delta_n.$$

Choosing $\tilde{\delta} = \min(\delta, \delta_1, \delta_2, \dots, \delta_{n_0})$, we obtain

$$\sup_{n \in \mathbb{N}} E[|X_n|; A] < \varepsilon \quad \text{whenever} \quad P[A] < \tilde{\delta}.$$

Therefore, $\{X_n : n \in \mathbb{N}\}$ is uniformly integrable by the exercise below the definition of uniform integrability on page 143. □

L^1 convergence of martingales

If X is an integrable random variable and (\mathcal{F}_n) is a filtration then $M_n = E[X | \mathcal{F}_n]$ is a martingale w.r.t. (\mathcal{F}_n) . The next result shows that an arbitrary martingale can be represented in this way if and only if it is uniformly integrable:

Theorem 4.14 (L^1 Martingale Convergence Theorem). *Suppose that $(M_n)_{n \geq 0}$ is a martingale w.r.t. a filtration (\mathcal{F}_n) . Then the following statements are equivalent:*

- (1). $\{M_n : n \geq 0\}$ is uniformly integrable.
- (2). The sequence (M_n) converges w.r.t. the L^1 norm.
- (3). There exists an integrable random variable X such that

$$M_n = E[X | \mathcal{F}_n] \quad \text{for any } n \geq 0.$$

Proof.

(3) \Rightarrow (1) holds by Lemma 4.12.

(1) \Rightarrow (2): If the sequence (M_n) is uniformly integrable then it is bounded in L^1 because

$$\sup_n E[|M_n|] \leq \sup_n E[|M_n|; |M_n| \geq c] + c \leq 1 + c$$

for $c \in (0, \infty)$ sufficiently large. Therefore, the limit $M_\infty = \lim M_n$ exists almost surely and in probability by the almost sure convergence theorem. Uniform integrability then implies

$$M_n \rightarrow M_\infty \quad \text{in } L^1$$

by Theorem 4.13.

(2) \Rightarrow (3): If M_n converges to a limit M_∞ in L^1 then

$$M_n = E[M_\infty | \mathcal{F}_n] \quad \text{for any } n \geq 0.$$

Indeed, M_n is a version of the conditional expectation since it is \mathcal{F}_n -measurable and

$$E[M_\infty; A] = \lim_{k \rightarrow \infty} E[M_k; A] = E[M_n; A] \quad \text{for any } A \in \mathcal{F}_n \quad (4.3.4)$$

by the martingale property. □

A first consequence of the L^1 convergence theorem is a limit theorem for conditional expectations:

Corollary 4.15. *If X is an integrable random variable and (\mathcal{F}_n) is a filtration then*

$$E[X | \mathcal{F}_n] \rightarrow E[X | \mathcal{F}_\infty] \quad \text{almost surely and in } L^1,$$

where $\mathcal{F}_\infty := \sigma(\bigcup \mathcal{F}_n)$.

Proof. Let $M_n := E[X | \mathcal{F}_n]$. By the almost sure and the L^1 martingale convergence theorem, the limit $M_\infty = \lim M_n$ exists almost surely and in L^1 . To obtain a measurable function that is defined everywhere, we set $M_\infty := \limsup M_n$. It remains to verify, that M_∞ is a version of the conditional expectation $E[X | \mathcal{F}_\infty]$. Clearly, M_∞ is measurable w.r.t. \mathcal{F}_∞ . Moreover, for $n \geq 0$ and $A \in \mathcal{F}_n$,

$$E[M_\infty; A] = E[M_n; A] = E[X; A]$$

by (4.3.4). Since $\bigcup \mathcal{F}_n$ is stable under finite intersections,

$$E[M_\infty ; A] = E[X ; A]$$

holds for all $A \in \sigma(\bigcup \mathcal{F}_n)$ as well. \square

Example (Existence of conditional expectations). The common existence proof for conditional expectations relies either on the Radon-Nikodym Theorem or on the existence of orthogonal projections onto closed subspaces of the Hilbert space L^2 . Martingale convergence can be used to give an alternative existence proof. Suppose that X is an integrable random variable on a probability space (Ω, \mathcal{A}, P) and \mathcal{F} is a **separable** sub- σ -algebra of \mathcal{A} , i.e., there exists a countable collection $(A_i)_{i \in \mathbb{N}}$ of events $A_i \in \mathcal{A}$ such that $\mathcal{F} = \sigma(A_i : i \in \mathbb{N})$. Let

$$\mathcal{F}_n = \sigma(A_1, \dots, A_n), \quad n \geq 0.$$

Note that for each $n \geq 0$, there exist finitely many atoms $B_1, \dots, B_k \in \mathcal{A}$ (i.e. disjoint events with $\bigcup B_i = \Omega$) such that $\mathcal{F}_n = \sigma(B_1, \dots, B_k)$. Therefore, the conditional expectation given \mathcal{F}_n can be defined in an elementary way:

$$E[X | \mathcal{F}_n] := \sum_{i: P[B_i] \neq 0} E[X | B_i] \cdot I_{B_i}.$$

Moreover, by Corollary 4.15, the limit $M_\infty = \lim E[X | \mathcal{F}_n]$ exists almost surely and in L^1 , and M_∞ is a version of the conditional expectation $E[X | \mathcal{F}]$.

You might (and should) object that the proofs of the martingale convergence theorems require the existence of conditional expectations. Although this is true, it is possible to state the necessary results by using only elementary conditional expectations, and thus to obtain a more constructive proof for existence of conditional expectations given separable σ -algebras.

Another immediate consequence of Corollary 4.15 is an extension of Kolmogorov's 0-1 law:

Corollary 4.16 (0-1 Law of P.Lévy). *If (\mathcal{F}_n) is a filtration on (Ω, \mathcal{A}, P) then for any event $A \in \sigma(\bigcup \mathcal{F}_n)$,*

$$P[A | \mathcal{F}_n] \longrightarrow I_A \quad P\text{-almost surely.} \quad (4.3.5)$$

Example (Kolmogorov's 0-1 Law). Suppose that $\mathcal{F}_n = \sigma(\mathcal{A}_1, \dots, \mathcal{A}_n)$ with independent σ -algebras $\mathcal{A}_i \subseteq \mathcal{A}$. If A is a **tail event**, i.e., A is in $\sigma(\mathcal{A}_{n+1}, \mathcal{A}_{n+2}, \dots)$ for every $n \in \mathbb{N}$, then A is independent of \mathcal{F}_n for any n . Therefore, the corollary implies that $P[A] = I_A$ P -almost surely, i.e.,

$$P[A] \in \{0, 1\} \quad \text{for any tail event } A.$$

The L^1 Martingale Convergence Theorem also implies that any martingale that is L^p bounded for some $p \in (1, \infty)$ converges in L^p :

Exercise (L^p Martingale Convergence Theorem). Let (M_n) be an (\mathcal{F}_n) martingale with $\sup E[|M_n|^p] < \infty$ for some $p \in (1, \infty)$.

- (1). Prove that (M_n) converges almost surely and in L^1 , and $M_n = E[M_\infty | \mathcal{F}_n]$ for any $n \geq 0$.
- (2). Conclude that $|M_n - M_\infty|^p$ is uniformly integrable, and $M_n \rightarrow M_\infty$ in L^p .

Note that uniform integrability of $|M_n|^p$ holds automatically and has not to be assumed !

Backward Martingale Convergence

We finally remark that Doob's upcrossing inequality can also be used to prove that the conditional expectations $E[X | \mathcal{F}_n]$ of an integrable random variable given a *decreasing* sequence (\mathcal{F}_n) of σ -algebras converge almost surely to $E[X | \bigcap \mathcal{F}_n]$. For the proof one considers the martingale $M_{-n} = E[X | \mathcal{F}_n]$ indexed by the negative integers:

Exercise (Backward Martingale Convergence Theorem and LLN). Let $(\mathcal{F}_n)_{n \geq 0}$ be a *decreasing* sequence of sub- σ -algebras on a probability space (Ω, \mathcal{A}, P) .

- (1). Prove that for any random variable $X \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$, the limit $M_{-\infty}$ of the sequence $M_{-n} := E[X | \mathcal{F}_n]$ as $n \rightarrow -\infty$ exists almost surely and in L^1 , and

$$M_{-\infty} = E[X | \bigcap \mathcal{F}_n] \quad \text{almost surely.}$$

- (2). Now let (X_n) be a sequence of i.i.d. random variables in $\mathcal{L}^1(\Omega, \mathcal{A}, P)$, and let $\mathcal{F}_n = \sigma(S_n, S_{n+1}, \dots)$ where $S_n = X_1 + \dots + X_n$. Prove that

$$E[X_1 | \mathcal{F}_n] = \frac{S_n}{n},$$

and conclude that the strong Law of Large Numbers holds:

$$\frac{S_n}{n} \longrightarrow E[X_1] \quad \text{almost surely.}$$

Chapter 5

Stochastic Integration w.r.t. Continuous Martingales

Suppose that we are interested in a continuous-time scaling limit of a stochastic dynamics of type $X_0^{(h)} = x_0$,

$$X_{k+1}^{(h)} - X_k^{(h)} = \sigma(X_k^{(h)}) \cdot \sqrt{h} \cdot \eta_{k+1}, \quad k = 0, 1, 2, \dots, \quad (5.0.1)$$

with i.i.d. random variables $\eta_i \in \mathcal{L}^2$ such that $E[\eta_i] = 0$ and $\text{Var}[\eta_i] = 1$, a continuous function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$, and a scale factor $h > 0$. Equivalently,

$$X_n^{(h)} = X_0^{(h)} + \sqrt{h} \cdot \sum_{k=0}^{n-1} \sigma(X_k^{(h)}) \cdot \eta_{k+1}, \quad n = 0, 1, 2, \dots \quad (5.0.2)$$

If σ is constant then as $h \searrow 0$, the rescaled process $(X_{\lfloor t/h \rfloor}^{(h)})_{t \geq 0}$ converges in distribution to $(\sigma \cdot B_t)$ where (B_t) is a Brownian motion. We are interested in the scaling limit for general σ . One can prove that the rescaled process again converges in distribution, and the limit process is a solution of a stochastic integral equation

$$X_t = X_0 + \int_0^t \sigma(X_s) dB_s, \quad t \geq 0. \quad (5.0.3)$$

Here the integral is an Itô stochastic integral w.r.t. a Brownian motion (B_t) . Usually the equation (5.0.3) is written briefly as

$$dX_t = \sigma(X_t) dB_t, \quad (5.0.4)$$

and interpreted as a stochastic differential equation. Stochastic differential equations occur more generally when considering scaling limits of appropriately rescaled Markov chains on \mathbb{R}^d with finite second moments. The goal of this section is to give a meaning to the stochastic integral, and hence to the equations (5.0.3), (5.0.4) respectively.

Example (Stock prices, geometric Brownian motion). A simple discrete time model for stock prices is given by

$$X_{k+1} - X_k = X_k \cdot \eta_{k+1}, \quad \eta_i \text{ i.i.d.}$$

To set up a corresponding continuous time model we consider the rescaled equation (5.0.1) as $h \searrow 0$. The limit in distribution is a solution of a stochastic differential equation

$$dX_t = X_t dB_t \quad (5.0.5)$$

w.r.t. a Brownian motion (B_t) . Although with probability one, the sample paths of Brownian motion are nowhere differentiable, we can give a meaning to this equation by rewriting it in the form (5.0.3) with an Itô stochastic integral.

A naive guess would be that the solution of (5.0.5) with initial condition $X_0 = 1$ is $X_t = \exp B_t$. However, more careful considerations show that this can not be true! In fact, the discrete time approximations satisfy

$$X_{k+1}^{(h)} = (1 + \sqrt{h}\eta_{k+1}) \cdot X_k^{(h)} \quad \text{for } k \geq 0.$$

Hence $(X_k^{(h)})$ is a product martingale:

$$X_n^{(h)} = \prod_{k=1}^n (1 + \sqrt{h}\eta_k) \quad \text{for any } n \geq 0.$$

In particular, $E[X_n^{(h)}] = 1$. We would expect similar properties for the scaling limit (X_t) , but $\exp B_t$ is not a martingale and $E[\exp(B_t)] = \exp(t/2)$.

It turns out that in fact, the unique solution of (5.0.5) with $X_0 = 1$ is not $\exp(B_t)$ but the exponential martingale

$$X_t = \exp(B_t - t/2),$$

which is also called a geometric Brownian motion. The reason is that the irregularity of Brownian paths enforces a correction term in the chain rule for stochastic differentials leading to Itô's famous formula, which is the fundament of stochastic calculus.

5.1 Defining stochastic integrals: A first attempt

Let us first fix some notation that will be used constantly below: By a **partition** π of \mathbb{R}_+ we mean an increasing sequence $0 = t_0 < t_1 < t_2 < \dots$ such that $\sup t_n = \infty$. The *mesh size* of the partition is

$$\text{mesh}(\pi) = \sup\{|t_i - t_{i-1}| : i \in \mathbb{N}\}.$$

We are interested in defining integrals of type

$$I_t = \int_0^t H_s dX_s, \quad t \geq 0, \quad (5.1.1)$$

for continuous functions and, respectively, continuous adapted processes (H_s) and (X_s) .

For a given $t \geq 0$ and a given partition π of \mathbb{R}_+ , we define the increments of (X_s) up to time t by

$$\delta X_s := X_{s' \wedge t} - X_{s \wedge t} \quad \text{for any } s \in \pi,$$

where $s' := \min\{u \in \pi : u > s\}$ denotes the next partition point after s . Note that the increments δX_s vanish for $s \geq t$. In particular, only finitely many of the increments are not equal to zero. A nearby approach for defining the integral I_t in (5.1.1) would be Riemann sum approximations:

Riemann sum approximations

There are various possibilities to define approximating Riemann sums w.r.t. a given sequence (π_n) of partitions with $\text{mesh}(\pi_n) \rightarrow 0$, for example:

$$\text{Variant 1 (non-anticipative): } I_t^n = \sum_{s \in \pi_n} H_s \delta X_s,$$

$$\text{Variant 2 (anticipative): } \hat{I}_t^n = \sum_{s \in \pi_n} H_{s'} \delta X_s,$$

$$\text{Variant 3 (anticipative): } \overset{\circ}{I}_t^n = \sum_{s \in \pi_n} \frac{1}{2}(H_s + H_{s'}) \delta X_s.$$

Note that for finite t , in each of the sums, only finitely many summands do not vanish. For example,

$$I_t^n = \sum_{\substack{s \in \pi_n \\ s < t}} H_s \delta X_s = \sum_{\substack{s \in \pi_n \\ s < t}} H_s \cdot (X_{s' \wedge t} - X_s).$$

Now let us consider at first the case where $H_s = X_s$ and $t = 1$, i.e., we would like to define the integral $I = \int_0^1 X_s dX_s$. Suppose first that $X : [0, 1] \rightarrow \mathbb{R}$ is a continuous function of finite variation, i.e.,

$$V^{(1)}(X) = \sup \left\{ \sum_{s \in \pi} |\delta X_s| : \pi \text{ partition of } \mathbb{R}_+ \right\} < \infty.$$

Then for $H = X$ and $t = 1$ all the approximations above converge to the same limit as $n \rightarrow \infty$. For example,

$$\|\hat{I}_1^n - I_1^n\| = \sum_{s \in \pi_n} (\delta X_s)^2 \leq V^{(1)}(X) \cdot \sup_{s \in \pi_n} |\delta X_s|,$$

and the right-hand side converges to 0 by uniform continuity of X on $[0, 1]$. In this case the limit of the Riemann sums is a Riemann-Stieltjes integral

$$\lim_{n \rightarrow \infty} I_1^n = \lim_{n \rightarrow \infty} \hat{I}_1^n = \int_0^1 X_s dX_s,$$

which is well-defined whenever the integrand is continuous and the integrator is of finite variation or conversely. The sample paths of Brownian motion, however, are almost surely not of finite variation. Therefore, the reasoning above does not apply, and in fact if $X_t = B_t$ is a one-dimensional Brownian motion and $H_t = X_t$ then

$$E[|\hat{I}_1^n - I_1^n|] = \sum_{s \in \pi_n} E[(\delta B_s)^2] = \sum_{s \in \pi_n} \delta s = 1,$$

i.e., the L^1 -limits of the random sequence (I_1^n) and (\hat{I}_1^n) are different if they exist. Below we will see that indeed the limits of the sequences (I_1^n) , (\hat{I}_1^n) and $(I_1^{\circ n})$ do exist in L^2 , and all the limits are different. The limit of the non-anticipative Riemann sums I_1^n is the *Itô stochastic integral* $\int_0^1 B_s dB_s$, the limit of (\hat{I}_1^n) is the *backward Itô integral* $\int_0^1 B_s \hat{d}B_s$, and the limit of $I_1^{\circ n}$ is the *Stratonovich integral* $\int_0^1 B_s \circ dB_s$. All three notions

of stochastic integrals are relevant. The most important one is the Itô integral because the non-anticipating Riemann sum approximations imply that the Itô integral $\int_0^t H_s dB_s$ is a continuous time martingale transform of Brownian motion if the process (H_s) is adapted.

Itô integrals for continuous bounded integrands

We now give a first existence proof for Itô integrals w.r.t. Brownian motion. We start with a provisional definition that will be made more precise later:

Preliminary Definition. *For continuous functions or continuous stochastic processes (H_s) and (X_s) and a given sequence (π_n) of partitions with $\text{mesh}(\pi_n) \rightarrow 0$, the **Itô integral of H w.r.t. X** is defined by*

$$\int_0^t H_s dX_s = \lim_{n \rightarrow \infty} \sum_{s \in \pi_n} H_s \delta X_s$$

whenever the limit exists in a sense to be specified.

Note that the definition is vague since the mode of convergence is not specified. Moreover, the Itô integral might depend on the sequence (π_n) . In the following sections we will see which kind of convergence holds in different circumstances, and in which sense the limit is independent of (π_n) .

To get started let us consider the convergence of Riemann sum approximations for the Itô integrals $\int_0^t H_s dB_s$ of a bounded continuous (\mathcal{F}_s) adapted process $(H_s)_{s \geq 0}$ w.r.t. an (\mathcal{F}_s) Brownian motion (B_s) . Let (π_n) be a fixed sequence of partitions with $\pi_n \subseteq \pi_{n+1}$ and $\text{mesh}(\pi_n) \rightarrow 0$. Then for the Riemann-Itô sums

$$I_t^n = \sum_{s \in \pi_n} H_s \delta B_s = \sum_{\substack{s \in \pi_n \\ s < t}} H_s (B_{s' \wedge t} - B_s)$$

we have

$$I_t^n - I_t^m = \sum_{\substack{s \in \pi_n \\ s < t}} (H_s - H_{[s]_m}) \delta B_s \quad \text{for any } m \leq n,$$

where $\lfloor s \rfloor_m = \max\{r \in \pi_m : r \leq s\}$ denotes the next partition point in π_m below s . Since Brownian motion is a martingale, we have $E[\delta B_s | \mathcal{F}_s] = 0$ for any $s \in \pi_n$. Moreover, $E[(\delta B_s)^2 | \mathcal{F}_s] = \delta s$. Therefore, we obtain by conditioning on $\mathcal{F}_s, \mathcal{F}_r$ respectively:

$$\begin{aligned} E[(I_t^n - I_t^m)^2] &= \sum_{\substack{s \in \pi_n \\ s < t}} \sum_{\substack{r \in \pi_m \\ r < t}} E[(H_s - H_{\lfloor s \rfloor_m})(H_r - H_{\lfloor r \rfloor_m})\delta B_s \delta B_r] \\ &= \sum_{\substack{s \in \pi_n \\ s < t}} E[(H_s - H_{\lfloor s \rfloor_m})^2 \delta s] \leq E[V_m] \cdot \sum_{\substack{s \in \pi_n \\ s < t}} \delta s = E[V_m] \cdot t, \end{aligned}$$

where

$$V_m := \sup_{|s-r| \leq \text{mesh}(\pi_m)} (H_s - H_r)^2 \longrightarrow 0 \quad \text{as } m \rightarrow \infty$$

by uniform continuity of (H_s) on $[0, t]$. Since H is bounded, $E[V_m] \rightarrow 0$ as $m \rightarrow \infty$, and hence (I_t^n) is a Cauchy sequence in $L^2(\Omega, \mathcal{A}, P)$ for any given $t \geq 0$. Thus we obtain:

Theorem 5.1 (Itô integrals for bounded continuous integrands, Variant 1). *Suppose that $(H_s)_{s \geq 0}$ is a bounded continuous (\mathcal{F}_s) adapted process, and $(B_s)_{s \geq 0}$ is an (\mathcal{F}_s) Brownian motion. Then for any fixed $t \geq 0$, the Itô integral*

$$\int_0^t H_s dB_s = \lim_{n \rightarrow \infty} I_t^n \quad (5.1.2)$$

exists as a limit in $L^2(\Omega, \mathcal{A}, P)$. Moreover, the limit does not depend on the choice of a sequence of partitions (π_n) with $\text{mesh}(\pi_n) \rightarrow 0$.

Proof. An analogue argument as above shows that for any partitions π and $\tilde{\pi}$ such that $\pi \supseteq \tilde{\pi}$, the L^2 distance of the corresponding Riemann sum approximations I_t^π and $I_t^{\tilde{\pi}}$ is bounded by a constant $C(\text{mesh}(\tilde{\pi}))$ that only depends on the maximal mesh size of the two partitions. Moreover, the constant goes to 0 as the mesh sizes go to 0. By choosing a joint refinement and applying the triangle inequality, we see that

$$\|I_t^\pi - I_t^{\tilde{\pi}}\|_{L^2(P)} \leq 2C(\Delta)$$

holds for arbitrary partitions $\pi, \tilde{\pi}$ such that $\max(\text{mesh}(\pi), \text{mesh}(\tilde{\pi})) \leq \Delta$. The assertion now follows by completeness of $L^2(P)$. \square

The definition of the Itô integral suggested by Theorem 5.1 has two obvious drawbacks:

Drawback 1: The integral $\int_0^t H_s dB_s$ is only defined as an equivalence class in $L^2(\Omega, \mathcal{A}, P)$, i.e., uniquely up to modification on P -measure zero sets. In particular, we do not have a *pathwise definition* of $\int_0^t H_s(\omega) dB_s(\omega)$ for a given Brownian sample path $s \mapsto B_s(\omega)$.

Drawback 2: Even worse, the construction above works only for a fixed integration interval $[0, t]$. The exceptional sets may depend on t and therefore, the process $t \mapsto \int_0^t H_s dB_s$ does not have a meaning yet. In particular, we do not know yet if there exists a version of this process that is almost surely continuous.

The first drawback is essential: In certain cases it is indeed possible to define stochastic integrals pathwise, cf. Chapter 6 below. In general, however, pathwise stochastic integrals cannot be defined. The extra impact needed is the Lévy area process, cf. the rough paths theory developed by T. Lyons and others [XXXLyons, Friz and Victoir, Friz and Hairer].

Fortunately, the second drawback can be overcome easily. By extending the Itô isometry to an isometry into the space M_c^2 of continuous L^2 bounded martingales, we can construct the complete process $t \mapsto \int_0^t H_s dB_s$ simultaneously as a continuous martingale. The key observation is that by the maximal inequality, continuous L^2 bounded martingales can be controlled uniformly in t by the L^2 norm of their final value.

The Hilbert space M_c^2

Fix $u \in (0, \infty]$ and suppose that for $t \in [0, u]$, (I_t^n) is a sequence of Riemann sum approximations for $\int_0^t H_s dB_s$ as considered above. It is not difficult to check that for each fixed $n \in \mathbb{N}$, the stochastic process $t \mapsto I_t^n$ is a continuous martingale. Our aim is to prove convergence of these continuous martingales to a further continuous martingale $I_t = \int_0^t H_s dB_s$. Since the convergence holds only almost surely, the limit process

will not necessarily be (\mathcal{F}_t) adapted. To ensure adaptedness, we have to consider the **completed filtration**

$$\mathcal{F}_t^P = \{A \in \mathcal{A} : P[A \Delta B] = 0 \text{ for some } B \in \mathcal{F}_t\}, \quad t \geq 0,$$

where $A \Delta B = (A \setminus B) \cup (B \setminus A)$ is the symmetric difference of the sets A and B . Note that the conditional expectations given \mathcal{F}_t and \mathcal{F}_t^P agree P -almost surely. Hence, if (B_t) is a Brownian motion resp. a martingale w.r.t. the filtration (\mathcal{F}_t) then it is also a Brownian motion or a martingale w.r.t. (\mathcal{F}_t^P) .

Let $\mathcal{M}^2([0, u])$ denote the space of all L^2 -bounded (\mathcal{F}_t^P) martingales $(M_t)_{0 \leq t \leq u}$ on (Ω, \mathcal{A}, P) . By $\mathcal{M}_c^2([0, u])$ and $\mathcal{M}_d^2([0, u])$ we denote the subspaces consisting of all continuous (respectively right continuous) martingales $M \in \mathcal{M}^2([0, u])$. Recall that by the L^2 martingale convergence theorem, any (right) continuous L^2 -bounded martingale (M_t) defined for $t \in [0, u)$ can be extended to a (right) continuous martingale in $\mathcal{M}^2([0, u])$.

Two martingales $M, \widetilde{M} \in \mathcal{M}^2([0, u])$ are called **modifications** of each other if

$$P[M_t = \widetilde{M}_t] = 1 \quad \text{for any } t \in [0, u].$$

If the martingales are right-continuous then two modifications agree almost surely, i.e.,

$$P[M_t = \widetilde{M}_t \forall t \in [0, u]] = 1.$$

In order to obtain norms and not just semi-norms, we consider the spaces

$$M^2([0, u]) := \mathcal{M}^2([0, u]) / \sim \quad \text{and} \quad M_c^2([0, u]) := \mathcal{M}_c^2([0, u]) / \sim$$

of equivalence classes of martingales that are modifications of each other. We will frequently identify equivalence classes and their representatives.

We endow the space $M^2([0, u])$ with the inner product

$$(M, N)_{M^2([0, u])} = (M_u, N_u)_{L^2} = E[M_u N_u].$$

As the process (M_t^2) is a submartingale for any $M \in M^2([0, u])$, the norm corresponding to the inner product is given by

$$\|M\|_{M^2([0, u])}^2 = E[M_u^2] = \sup_{0 \leq t \leq u} E[M_t^2].$$

Moreover, if (M_t) is right continuous then by **Doob's L^2 -maximal inequality**,

$$\left\| \sup_{0 \leq t \leq u} |M_t| \right\|_{L^2(\Omega, \mathcal{A}, P)} \leq 2 \cdot \sup_{0 \leq t \leq u} \|M_t\|_{L^2(\Omega, \mathcal{A}, P)} = 2\|M\|_{M^2([0, u])}. \quad (5.1.3)$$

This crucial estimate shows that on the subspaces M_c^2 and M_d^2 , the M^2 norm is equivalent to the L^2 norm of the supremum of the martingale. Therefore, *the M^2 norm can be used to control (right) continuous martingales uniformly in t !*

Lemma 5.2 (Completeness). (1). *The space $M^2([0, u])$ is a Hilbert space, and the linear map $M \mapsto M_u$ from $M^2([0, u])$ to $L^2(\Omega, \mathcal{F}_u, P)$ is onto and isometric.*

(2). *The spaces $M_c^2([0, u])$ and $M_d^2([0, u])$ are closed subspaces of $M^2([0, u])$, i.e., if (M^n) is a Cauchy sequence in $M_c^2([0, u])$, or in $M_d^2([0, u])$ respectively, then there exists a (right) continuous martingale $M \in M^2([0, u])$ such that*

$$\sup_{t \in [0, u]} |M_t^n - M_t| \longrightarrow 0 \quad \text{in } L^2(\Omega, \mathcal{A}, P).$$

Proof. (1). By definition of the inner product on $M^2([0, u])$, the map $M \mapsto M_u$ is an isometry. Moreover, for $X \in L^2(\Omega, \mathcal{F}_u, P)$, the process $M_t = E[X | \mathcal{F}_t]$ is in $M^2([0, u])$ with $M_u = X$. Hence, the range of the isometry is the whole space $L^2(\Omega, \mathcal{F}_u, P)$. Since $L^2(\Omega, \mathcal{F}_u, P)$ is complete w.r.t. the L^2 norm, the space $M^2([0, u])$ is complete w.r.t. the M^2 norm.

(2). If (M^n) is a Cauchy sequence in $M_c^2([0, u])$ or in $M_d^2([0, u])$ respectively, then by (5.1.3),

$$\|M^n - M^m\|_{\text{sup}} := \sup_{0 \leq t \leq u} |M_t^n - M_t^m| \longrightarrow 0 \quad \text{in } L^2(\Omega, \mathcal{A}, P).$$

In particular, we can choose a subsequence (M^{n_k}) such that

$$P[\|M^{n_{k+1}} - M^{n_k}\|_{\text{sup}} \geq 2^{-k}] \leq 2^{-k} \quad \text{for all } k \in \mathbb{N}.$$

Hence, by the Borel-Cantelli Lemma,

$$P[\|M^{n_{k+1}} - M^{n_k}\|_{\text{sup}} < 2^{-k} \text{ eventually}] = 1,$$

and therefore $M_t^{n_k}$ converges almost surely uniformly in t as $k \rightarrow \infty$. The limit of the sequence (M^n) in $M^2([0, u])$ exists by (1), and the process M defined by

$$M_t := \begin{cases} \lim M_t^{n_k} & \text{if } (M^{n_k}) \text{ converges uniformly,} \\ 0 & \text{otherwise,} \end{cases} \quad (5.1.4)$$

is a continuous (respectively right continuous) representative of the limit. Indeed, by Fatou's Lemma,

$$\begin{aligned} \|M^{n_k} - M\|_{M^2([0, u])}^2 &\leq E[\|M^{n_k} - M\|_{\text{sup}}^2] = E[\lim_{l \rightarrow \infty} \|M^{n_k} - M^{n_l}\|_{\text{sup}}^2] \\ &\leq \liminf_{l \rightarrow \infty} E[\|M^{n_k} - M^{n_l}\|_{\text{sup}}^2], \end{aligned}$$

and the right hand side converges to 0 as $k \rightarrow \infty$. Finally, M is a martingale w.r.t. (\mathcal{F}_t^P) , and hence an element in $M_c^2([0, u])$ or in $M_d^2([0, u])$ respectively.

□

Remark. We point out that the (right) continuous representative (M_t) defined by (5.1.4) is a martingale w.r.t. the complete filtration (\mathcal{F}_t^P) , but it is not necessarily adapted w.r.t. (\mathcal{F}_t) .

Definition of Itô integral in M_c^2

Let $u \in \mathbb{R}^+$. For any bounded continuous (\mathcal{F}_t) adapted process (H_t) and any sequence (π_n) of partitions of \mathbb{R}_+ , the processes

$$I_t^n = \sum_{s \in \pi_n} H_s (B_{s' \wedge t} - B_{s \wedge t}), \quad t \in [0, u],$$

are continuous L^2 bounded martingales on $[0, u]$. We can therefore restate Theorem 5.1 in the following way:

Corollary 5.3 (Itô integrals for bounded continuous integrands, Variant 2). *Suppose that $(H_s)_{s \geq 0}$ is a bounded continuous (\mathcal{F}_s) adapted process. Then for any fixed $u \geq 0$, the Itô integral*

$$\int_0^\bullet H_s dB_s = \lim_{n \rightarrow \infty} (I_t^n)_{t \in [0, u]} \quad (5.1.5)$$

exists as a limit in $M_c^2([0, u])$. Moreover, the limit does not depend on the choice of a sequence of partitions (π_n) with $\text{mesh}(\pi_n) \rightarrow 0$.

Proof. The assertion is an immediate consequence of the definition of the M^2 norm, Theorem 5.1 and Lemma 5.2. □

Similar arguments as above apply if Brownian motion is replaced by a bounded martingale with continuous sample paths. In the rest of this chapter we will work out the construction of the Itô integral w.r.t. Brownian motion and more general continuous martingales more systematically and for a broader class of integrands.

5.2 Itô's isometry

Let $(M_t)_{t \geq 0}$ be a continuous (or, more generally, right continuous) martingale w.r.t. a filtration (\mathcal{F}_t) on a probability space (Ω, \mathcal{A}, P) . We now develop a more systematic approach for defining stochastic integrals $\int_0^t H_s dM_s$ of adapted processes (H_t) w.r.t. (M_t) .

Predictable step functions

In a first step, we define the integrals for predictable step functions (H_t) of type

$$H_t(\omega) = \sum_{i=0}^{n-1} A_i(\omega) I_{(t_i, t_{i+1}]}(t)$$

with $n \in \mathbb{N}$, $0 \leq t_0 < t_1 < t_2 < \dots < t_n$, and bounded \mathcal{F}_{t_i} -measurable random variables A_i , $i = 0, 1, \dots, n-1$. Let \mathcal{E} denote the vector space consisting of all stochastic processes of this form.

Definition (Itô integral for predictable step functions). For stochastic processes $H \in \mathcal{E}$ and $t \geq 0$ we define

$$\int_0^t H_s dM_s := \sum_{i=0}^{n-1} A_i \cdot (M_{t_{i+1} \wedge t} - M_{t_i \wedge t}) = \sum_{i: t_i < t} A_i \cdot (M_{t_{i+1} \wedge t} - M_{t_i}).$$

The stochastic process $H \bullet M$ given by

$$(H \bullet M)_t := \int_0^t H_s dM_s \quad \text{for } t \in [0, \infty]$$

is called the **Itô integral** of H w.r.t. M .

Note that the map $(H, M) \mapsto H \bullet M$ is bilinear. The process $H \bullet M$ is a continuous time **martingale transform** of M w.r.t. H . It models for example the net gain up to time t if we hold A_i units of an asset with price process (M_t) during each of the time intervals $(t_i, t_{i+1}]$.

Lemma 5.4. For any $H \in \mathcal{E}$, the process $H \bullet M$ is a continuous (\mathcal{F}_t) martingale up to time $t = \infty$.

Similarly to the discrete time case, the fact that A_i is predictable, i.e., \mathcal{F}_{t_i} -measurable, is essential for the martingale property:

Proof. By definition, $H \bullet M$ is continuous and (\mathcal{F}_t) adapted. It remains to verify that

$$E[(H \bullet M)_t | \mathcal{F}_s] = (H \bullet M)_s \quad \text{for any } 0 \leq s \leq t. \quad (5.2.1)$$

We do this in three steps:

- (1). At first we note that (5.2.1) holds for $s, t \in \{t_0, t_1, \dots, t_n\}$. Indeed, since A_i is \mathcal{F}_{t_i} -measurable, the process

$$(H \bullet M)_{t_j} = \sum_{i=0}^{j-1} A_i \cdot (M_{t_{i+1}} - M_{t_i}), \quad j = 0, 1, \dots, n,$$


is a martingale transform of the discrete time martingale (M_{t_i}) , and hence again a martingale.

(2). Secondly, suppose $s, t \in [t_j, t_{j+1}]$ for some $j \in \{0, 1, 2, \dots, n-1\}$. Then

$$E[(H \bullet M)_t - (H \bullet M)_s | \mathcal{F}_s] = E[A_j \cdot (M_t - M_s) | \mathcal{F}_s] = A_j \cdot E[M_t - M_s | \mathcal{F}_s] = 0$$

because A_j is \mathcal{F}_{t_j} -measurable and hence \mathcal{F}_s -measurable, and (M_t) is a martingale.

(3). Finally, suppose that $s \in [t_j, t_{j+1}]$ and $t \in [t_k, t_{k+1}]$ with $j < k$.



Then by the tower property for conditional expectations and by (1) and (2),

$$\begin{aligned} E[(H \bullet M)_t | \mathcal{F}_s] &= E[E[E[(H \bullet M)_t | \mathcal{F}_{t_k}] | \mathcal{F}_{t_{j+1}}] | \mathcal{F}_s] \\ &\stackrel{(2)}{=} E[E[(H \bullet M)_{t_k} | \mathcal{F}_{t_{j+1}}] | \mathcal{F}_s] \stackrel{(1)}{=} E[(H \bullet M)_{t_{j+1}} | \mathcal{F}_s] \\ &\stackrel{(2)}{=} (H \bullet M)_s. \end{aligned}$$

□

Remark (Riemann sum approximations). Non-anticipative Riemann sum approximations of stochastic integrals are Itô integrals of predictable step functions: If (H_t) is an adapted stochastic process and $\pi = \{t_0, t_1, \dots, t_n\}$ is a partition then

$$\sum_{i=0}^{n-1} H_{t_i} \cdot (M_{t_{i+1} \wedge t} - M_{t_i \wedge t}) = \int_0^t H_s^\pi dM_s \tag{5.2.2}$$

where $H^\pi := \sum_{i=0}^{n-1} H_{t_i} \cdot I_{(t_i, t_{i+1}]}$ is a process in \mathcal{E} .

Itô's isometry for Brownian motion

Recall that our goal is to prove that non-anticipative Riemann sum approximations for a stochastic integral converge. Let (π_n) be a sequence of partitions of $[0, t]$ with $\text{mesh}(\pi_n) \rightarrow 0$. By the remark above, the corresponding Riemann-Itô sums I^{π_n} defined by (5.2.2) are integrals of predictable step functions H^{π_n} . Hence in order to prove that the sequence (I^{π_n}) converges in the Hilbert space M_c^2 it suffices to show that

(1). (H^{π_n}) is a *Cauchy sequence w.r.t. an appropriate norm* on the vector space \mathcal{E} , and

(2). the “**Itô map**” $\mathcal{J} : \mathcal{E} \rightarrow M_c^2$ defined by

$$\mathcal{J}(H) = H \bullet M = \int_0^\bullet H_s dM_s$$

is *continuous w.r.t. this norm*.

It turns out that we can even identify explicitly a simple norm on \mathcal{E} such that the Itô map is an isometry. We first consider the case where (M_t) is a Brownian motion:

Theorem 5.5 (Itô's isometry for Brownian motion). *If (B_t) is an (\mathcal{F}_t) Brownian motion on (Ω, \mathcal{A}, P) then for any $u \in [0, \infty]$, and for any process $H \in \mathcal{E}$,*

$$\|H \bullet B\|_{M^2([0, u])}^2 = E \left[\left(\int_0^u H_s dB_s \right)^2 \right] = E \left[\int_0^u H_s^2 ds \right] = \|H\|_{L^2(P \otimes \lambda_{(0, u)})}^2 \quad (5.2.3)$$

Proof. Suppose that $H = \sum_{i=0}^{n-1} A_i \cdot I_{(t_i, t_{i+1}]}$ with $n \in \mathbb{N}$, $0 \leq t_0 < t_1 < \dots < t_n$ and A_i bounded and \mathcal{F}_{t_i} -measurable. With the notation $\delta_i B := B_{t_{i+1} \wedge u} - B_{t_i \wedge u}$, we obtain

$$E \left[\left(\int_0^u H_s dB_s \right)^2 \right] = E \left[\left(\sum_{i=0}^{n-1} A_i \delta_i B \right)^2 \right] = \sum_{i, k=0}^{n-1} E [A_i A_k \cdot \delta_i B \delta_k B]. \quad (5.2.4)$$

By the martingale property, the summands on the right hand side vanish for $i \neq k$. Indeed, if, for instance, $i < k$ then

$$E[A_i A_k \delta_i B \delta_k B] = E[A_i A_k \delta_i B \cdot E[\delta_k B \mid \mathcal{F}_{t_k}]] = 0.$$

Here we have used in an essential way, that A_k is \mathcal{F}_{t_k} -measurable. Similarly,

$$E[A_i^2 \cdot (\delta_i B)^2] = E[A_i^2 E[(\delta_i B)^2 \mid \mathcal{F}_{t_i}]] = E[A_i^2 \cdot \delta_i t]$$

by the independence of the increments of Brownian motion. Therefore, by (5.2.4) we obtain

$$E \left[\left(\int_0^u H_s dB_s \right)^2 \right] = \sum_{i=0}^{n-1} E[A_i^2 \cdot (t_{i+1} \wedge u - t_i \wedge u)] = E \left[\int_0^u H_s^2 ds \right].$$

The assertion now follows by definition of the M^2 norm. □

Theorem 5.5 shows that the linear map

$$\mathcal{J} : \mathcal{E} \rightarrow M_c^2([0, u]), \quad \mathcal{J}(H) = \left(\int_0^r H_s dB_s \right)_{r \in [0, u]},$$

is an isometry if the space \mathcal{E} of simple predictable processes $(s, \omega) \mapsto H_s(\omega)$ is endowed with the L^2 norm

$$\|H\|_{L^2(P \otimes \lambda_{(0, u)})} = E \left[\int_0^u H_s^2 ds \right]^{1/2}$$

on the product space $\Omega \times (0, u)$. In particular, \mathcal{J} respects $P \otimes \lambda$ classes, i.e., if $H_s(\omega) = \tilde{H}_s(\omega)$ for $P \otimes \lambda$ -almost every (ω, s) then $\int_0^\bullet H dB = \int_0^\bullet \tilde{H} dB$ P -almost surely. Hence \mathcal{J} also induces a linear map between the corresponding spaces of equivalence classes. As usual, we do not always differentiate between equivalence classes and functions, and so we denote the linear map on equivalence classes again by \mathcal{J} :

$$\begin{aligned} \mathcal{J} : \mathcal{E} \subset L^2(P \otimes \lambda_{(0, u)}) &\rightarrow M_c^2([0, u]), \\ \|\mathcal{J}(H)\|_{M^2([0, u])} &= \|H\|_{L^2(P \otimes \lambda_{(0, u)})}. \end{aligned} \tag{5.2.5}$$

Itô's isometry for martingales

An Itô isometry also holds if Brownian motion is replaced by a continuous square-integrable martingale (M_t) . More generally, suppose that $(M_t)_{t \geq 0}$ is a right continuous square integrable (\mathcal{F}_t) martingale satisfying the following assumption:

Assumption A. There exists a non-decreasing adapted continuous process $t \mapsto \langle M \rangle_t$ such that $\langle M \rangle_0 = 0$ and $M_t^2 - \langle M \rangle_t$ is a martingale.

For continuous square integrable martingales, the assumption is always satisfied. Indeed, assuming continuity, the “angle bracket process” $\langle M \rangle_t$ coincides almost surely with the quadratic variation process $[M]_t$ of M , cf. Section 6.3 below. For Brownian motion, Assumption A holds with

$$\langle B \rangle_t = t.$$

Note that for any $0 \leq s \leq t$, Assumption A implies

$$E[(M_t - M_s)^2 | \mathcal{F}_s] = E[M_t^2 - M_s^2 | \mathcal{F}_s] = E[\langle M \rangle_t - \langle M \rangle_s | \mathcal{F}_s]. \quad (5.2.6)$$

Since $t \mapsto \langle M \rangle_t(\omega)$ is continuous and non-decreasing for a given ω , it is the distribution function of a unique positive measure $\langle M \rangle(\omega, dt)$ on \mathbb{R}_+ .

Theorem 5.6 (Itô's isometry for martingales). *Suppose that $(M_t)_{t \geq 0}$ is a right continuous (\mathcal{F}_t) martingale with angle bracket process $\langle M \rangle$ satisfying Assumption A. Then for any $u \in [0, \infty]$, and for any process $H \in \mathcal{E}$,*

$$\|H \bullet M\|_{M^2([0, u])}^2 = E \left[\left(\int_0^u H_s dM_s \right)^2 \right] = E \left[\int_0^u H_s^2 d\langle M \rangle_s \right] \quad (5.2.7)$$

where $d\langle M \rangle$ denotes integration w.r.t. the positive measure with distribution function $F(t) = \langle M \rangle_t$.

For Brownian motion $\langle B \rangle_t = t$, so (5.2.7) reduces to (5.2.3).

Proof. The proof is similar to the proof of Theorem 5.5 above. Suppose again that $H = \sum_{i=0}^{n-1} A_i \cdot I_{(t_i, t_{i+1}]}$ with $n \in \mathbb{N}$, $0 \leq t_0 < t_1 < \dots < t_n$ and A_i bounded and \mathcal{F}_{t_i} -measurable. With the same notation as in the proof above, we obtain by the martingale properties of M and $M^2 - \langle M \rangle$,

$$E[A_i A_k \delta_i M \delta_k M] = 0 \quad \text{for } i \neq k, \quad \text{and}$$

$$E[A_i^2 \cdot (\delta_i M)^2] = E[A_i^2 E[(\delta_i M)^2 | \mathcal{F}_{t_i}]] = E[A_i^2 E[\delta_i \langle M \rangle | \mathcal{F}_{t_i}]] = E[A_i^2 \cdot \delta_i \langle M \rangle].$$

cf. (5.2.6). Therefore,

$$\begin{aligned} E \left[\left(\int_0^u H_s dM_s \right)^2 \right] &= E \left[\left(\sum_{i=0}^{n-1} A_i \delta_i M \right)^2 \right] = \sum_{i,k=0}^{n-1} E[A_i A_k \delta_i M \delta_k M] \\ &= \sum_{i=0}^{n-1} E[A_i^2 \delta_i \langle M \rangle] = E \left[\int_0^u H_s^2 d\langle M \rangle_s \right]. \end{aligned}$$

□

For a continuous square integrable martingale, Theorem 5.6 implies that the linear map

$$\mathcal{J} : \mathcal{E} \rightarrow \mathcal{M}_c^2([0, u]), \quad \mathcal{J}(H) = \left(\int_0^r H_s dM_s \right)_{r \in [0, u]},$$

is an isometry if the space \mathcal{E} of simple predictable processes $(s, \omega) \mapsto H_s(\omega)$ is endowed with the L^2 norm

$$\|H\|_{L^2(\Omega \times (0, u), P_{\langle M \rangle})} = E \left[\int_0^u H_s^2 d\langle M \rangle_s \right]^{1/2}$$

on the product space $\Omega \times (0, u)$ endowed with the positive measure

$$P_{\langle M \rangle}(d\omega dt) = P(d\omega) \langle M \rangle(\omega, dt). \quad (5.2.8)$$

Again, we denote the corresponding linear map induced on equivalence classes by the same letter \mathcal{J} .

Definition of Itô integrals for square-integrable integrands

From now on we assume that (M_t) is a *continuous* square integrable (\mathcal{F}_t) martingale with angle bracket process $\langle M \rangle_t$. We fix $u \in [0, \infty]$ and consider the isometry

$$\begin{aligned} \mathcal{J} : \mathcal{E} \subset L^2(\Omega \times (0, u), P_{\langle M \rangle}) &\rightarrow M_c^2([0, u]), \\ H &\mapsto H \bullet M \end{aligned} \quad (5.2.9)$$

mapping an elementary predictable process H to the continuous martingale

$$(H \bullet M)_t = \int_0^t H_s dM_s.$$

More precisely, we consider the induced map on equivalence classes.

Let $\overline{\mathcal{E}}_u$ denote the closure of the space \mathcal{E} in $L^2(\Omega \times (0, u), P_{\langle M \rangle})$. Since \mathcal{J} is linear with

$$\|\mathcal{J}(H)\|_{M^2([0, u])} = \|H\|_{L^2(\Omega \times (0, u), P_{\langle M \rangle})} \quad \text{for any } H \in \mathcal{E},$$

there is a unique extension to a continuous (and even isometric) linear map

$$\overline{\mathcal{J}} : \overline{\mathcal{E}}_u \subseteq L^2(\Omega \times (0, u), P_{\langle M \rangle}) \rightarrow M_c^2([0, u]).$$

This can be used to define the Itô integral for any process in $\overline{\mathcal{E}}_u$, i.e., for any process that can be approximated by predictable step functions w.r.t. the $L^2(P_{\langle M \rangle})$ norm:

$$H \bullet B := \overline{\mathcal{J}}(H), \quad \int_0^t H_s dB_s := (H \bullet B)_t.$$

Explicitly, we obtain the following definition of stochastic integrals for integrands in $\overline{\mathcal{E}}_u$:

Definition (Itô integral). For $H \in \overline{\mathcal{E}}_u$ the process $H \bullet M = (\int_0^t H_s dM_s)_{t \in [0, u]}$ is the up to modifications unique continuous martingale on $[0, u]$ satisfying

$$(H \bullet M)_t = \lim_{n \rightarrow \infty} (H^n \bullet M)_t \quad \text{in } L^2(P) \quad \text{for any } t \in [0, u]$$

whenever (H^n) is a sequence of elementary predictable processes such that $H^n \rightarrow H$ in $L^2(\Omega \times (0, u), P_{\langle M \rangle})$.

- Remark.** (1). By construction, the map $H \mapsto H \bullet M$ is an isometry from $\overline{\mathcal{E}_u}$ endowed with the $L^2(P_{\langle M \rangle})$ norm to $M_c^2([0, u])$. If $t \mapsto \langle M \rangle_t$ is absolutely continuous, then the closure $\overline{\mathcal{E}_u}$ of the elementary processes actually contains any (\mathcal{F}_t^P) adapted process $(\omega, t) \mapsto H_t(\omega)$ that is square-integrable w.r.t. $P_{\langle M \rangle}$, see XXX below.
- (2). The definition above is consistent in the following sense: If $H \bullet M$ is the stochastic integral defined on the time interval $[0, v]$ and $u \leq v$, then the restriction of $H \bullet M$ to $[0, u]$ coincides with the stochastic integral on $[0, u]$.

For $0 \leq s \leq t$ we define

$$\int_s^t H_r dM_r := (H \bullet M)_t - (H \bullet M)_s.$$

Exercise. Verify that for any $H \in \overline{\mathcal{E}_t}$,

$$\int_s^t H_r dM_r = \int_0^t H_r dB_r - \int_0^t I_{(0,s)}(r) H_r dM_r = \int_0^t I_{(s,t)}(r) H_r dM_r.$$

Having defined the Itô integral, we now show that bounded adapted processes with left-continuous sample paths are contained in the closure of the simple predictable processes, and the corresponding stochastic integrals are limits of predictable Riemann sum approximations. As above, we consider a sequence (π_n) of partitions of \mathbb{R}_+ such that $\text{mesh}(\pi_n) \rightarrow 0$.

Theorem 5.7 (Approximation by Riemann-Itô sums). *Let $u \in (0, \infty)$, and suppose that $(H_t)_{t \in [0, u]}$ is an (\mathcal{F}_t^P) adapted stochastic process on (Ω, \mathcal{A}, P) such that $(t, \omega) \mapsto H_t(\omega)$ is product-measurable and bounded. If $t \mapsto H_t$ is P -almost surely left continuous then H is in $\overline{\mathcal{E}_u}$, and*

$$\int_0^t H_s dM_s = \lim_{n \rightarrow \infty} \sum_{s \in \pi_n} H_s (M_{s' \wedge t} - M_{s \wedge t}), \quad t \in [0, u], \quad (5.2.10)$$

w.r.t. convergence uniformly in t in the $L^2(P)$ sense.

Remark. (1). In particular, a subsequence of the predictable Riemann sum approximations converges uniformly in t with probability one.

(2). The assertion also holds if H is unbounded with $\sup_{s \leq u} |H_s| \in \mathcal{L}^2(P)$.

Proof. For any $t \in [0, u]$, the Riemann sums on the right hand side of (5.2.10) are the stochastic integrals $\int_0^t H_s^n dM_s$ of the predictable step functions

$$H_t^n := \sum_{s \in \pi_n, s < t} H_s \cdot I_{(s, s^n]}(t), \quad n \in \mathbb{N}.$$

By left-continuity, $H_t^n \rightarrow H_t$ as $n \rightarrow \infty$ for any $t \in [0, u]$, P -almost surely. Therefore, $H^n \rightarrow H$ $P_{\langle M \rangle}$ -almost surely, and, by dominated convergence,

$$H^n \rightarrow H \quad \text{in } L^2(P_{\langle M \rangle}).$$

Here we have used that the sequence (H^n) is uniformly bounded since H is bounded by assumption. Now, by Itô's isometry,

$$\int_0^\bullet H_s dM_s = \lim_{n \rightarrow \infty} \int_0^\bullet H_s^n dM_s \quad \text{in } M_c^2([0, u]).$$

□

Identification of admissible integrands

Let $u \in (0, \infty]$. We have already shown that if $u < \infty$ then any product-measurable adapted bounded process with left-continuous sample paths is in $\overline{\mathcal{E}_u}$. More generally, we will prove now that if $M_t = B_t$ is a Brownian motion then any adapted process in $\mathcal{L}^2(P \otimes \lambda_{[0, u]})$ is contained in $\overline{\mathcal{E}_u}$, and hence “integrable” w.r.t. (B_t) . Let $\mathcal{L}_a^2(0, u)$ denote the linear space of all product-measurable, (\mathcal{F}_t^P) adapted stochastic processes $(\omega, t) \mapsto H_t(\omega)$ defined on $\Omega \times (0, u)$ such that

$$E \left[\int_0^u H_t^2 dt \right] < \infty.$$

The corresponding space of equivalence classes of $P \otimes \lambda$ versions is denoted by $L_a^2(0, u)$.

Lemma 5.8. $L_a^2(0, u)$ is a closed linear subspace of $L^2(P \otimes \lambda_{(0, u)})$.

Proof. It only remains to show that an $L^2(P \otimes \lambda)$ limit of (\mathcal{F}_t^P) adapted processes again has an (\mathcal{F}_t^P) adapted $P \otimes \lambda$ -version. Hence consider a sequence $H^n \in \mathcal{L}_a^2(0, u)$ with $H^n \rightarrow H$ in $L^2(P \otimes \lambda)$. Then there exists a subsequence (H^{n_k}) such that $H_t^{n_k}(\omega) \rightarrow H_t(\omega)$ for $P \otimes \lambda$ -almost every $(\omega, t) \in \Omega \times (0, u)$. The process \tilde{H} defined by $\tilde{H}_t(\omega) := \lim H_t^{n_k}(\omega)$ if the limit exists, $\tilde{H}_t(\omega) := 0$ otherwise, is an (\mathcal{F}_t^P) adapted version of H . \square

We can now identify the class of integrands H for which the stochastic integral $H \bullet B$ is well-defined as a limit of integrals of predictable step functions in $M_c^2([0, u])$:

Theorem 5.9 (Admissible integrands for Brownian motion). *For any $u \in (0, \infty]$,*

$$\overline{\mathcal{E}_u} = L_a^2(0, u).$$

Proof. Since $\mathcal{E} \subseteq \mathcal{L}_a^2(0, u)$ it only remains to show the inclusion “ \supseteq ”. Hence fix a process $H \in \mathcal{L}_a^2(0, u)$. We will prove in several steps that H can be approximated by simple predictable processes w.r.t. the $L^2(P \otimes \lambda_{(0,u)})$ norm:

- (1). Suppose first that H is bounded and has almost surely continuous trajectories. Then for $u < \infty$, H is in $\overline{\mathcal{E}_u}$ by Theorem 5.7. For $u = \infty$, H is still in $\overline{\mathcal{E}_u}$ provided there exists $t_0 \in (0, \infty)$ such that H_t vanishes for $t \geq t_0$.
- (2). Now suppose that (H_t) is bounded and, if $u = \infty$, vanishes for $t \geq t_0$. To prove $H \in \overline{\mathcal{E}_u}$ we approximate H by continuous adapted processes. To this end let $\psi_n : \mathbb{R} \rightarrow [0, \infty)$, $n \in \mathbb{N}$, be continuous functions such that $\psi(s) = 0$ for $s \notin (0, 1/n)$ and $\int_{-\infty}^{\infty} \psi_n(s) ds = 1$. Let $H^n := H * \psi_n$, i.e.,

$$H_t^n(\omega) = \int_0^{1/n} H_{t-\varepsilon}(\omega) \psi_n(\varepsilon) d\varepsilon, \quad (5.2.11)$$

where we set $H_t := 0$ for $t \leq 0$. We prove that

- (a) $H^n \rightarrow H$ in $L^2(P \otimes \lambda_{(0,u)})$, and
- (b) $H^n \in \overline{\mathcal{E}_u}$ for any $n \in \mathbb{N}$.

Combining (a) and (b), we see that H is in $\overline{\mathcal{E}_u}$ as well.

(a) Since H is in $\mathcal{L}^2(P \otimes \lambda_{(0,u)})$, we have

$$\int_0^u H_t(\omega)^2 dt < \infty \quad (5.2.12)$$

for P -almost every ω . It is a standard fact from analysis that (5.2.12) implies

$$\int_0^u |H_t^n(\omega) - H_t(\omega)|^2 dt \longrightarrow 0 \quad \text{as } n \rightarrow \infty.$$

By dominated convergence, we obtain

$$E \left[\int_0^u |H_t^n - H_t|^2 dt \right] \longrightarrow 0 \quad \text{as } n \rightarrow \infty \quad (5.2.13)$$

because H is bounded, the sequence (H_n) is uniformly bounded, and H and H^n vanish for $t \geq t_0 + 1$.

(b) This is essentially a consequence of part (1) of the proof. We sketch how to verify that H^n satisfies the assumptions made there:

- The sample paths $t \mapsto H_t^n(\omega)$ are continuous for all ω .
- $|H_t^n|$ is bounded by $\sup |H|$.
- The map $(\omega, t) \mapsto H_t^n(\omega)$ is product measurable by (5.2.11) and Fubini's Theorem, because the map $(\omega, t, \varepsilon) \mapsto H_{t-\varepsilon}(\omega)\psi_\varepsilon(\omega)$ is product measurable.
- If the process (H_t) is *progressively measurable*, i.e., if the map $(s, \omega) \mapsto H_s(\omega)$ ($s \in (0, t), \omega \in \Omega$) is measurable w.r.t. the product σ -algebra $\mathcal{B}(0, t) \otimes \mathcal{F}_t^P$ for any $t \geq 0$, then (H_t^n) is (\mathcal{F}_t^P) adapted by (5.2.11) and Fubini's Theorem. This is for example the case if (H_t) is right continuous or left continuous.
- In general, one can prove that (H_t) has a progressively measurable modification, whence (H_t^n) has an (\mathcal{F}_t^P) adapted modification. We omit the details.

(3). We finally prove that general $H \in \mathcal{L}_a^2(0, u)$ are contained in $\overline{\mathcal{E}_u}$. This is a consequence of (2), because we can approximate H by the processes

$$H_t^n := ((H_t \wedge n) \vee (-n)) \cdot I_{(0,n)}(t), \quad n \in \mathbb{N}.$$

These processes are bounded, they vanish for $t \geq n$, and $H^n \rightarrow H$ in $L^2(P \otimes \lambda_{(0,u)})$. By (2), H^n is contained in $\overline{\mathcal{E}_u}$ for any n , so H is in $\overline{\mathcal{E}_u}$ as well.

□

Remark (Riemann sum approximations). For discontinuous integrands, the predictable Riemann sum approximations considered above do not converge to the stochastic integral in general. However, one can prove that for $u < \infty$ any process $H \in L^2_a(0, u)$ is the limit of the simple predictable processes

$$H_t^n = \sum_{i=1}^{2^n-1} 2^n \int_{(i-1)2^{-n}u}^{i2^{-n}u} H_s ds \cdot I_{(i2^{-n}u, (i+1)2^{-n}u]}(t)$$

w.r.t. the $L^2(P \otimes \lambda_{[0,u]})$ norm, cf. [XXXSteele: “Stochastic calculus and financial applications”, Sect 6.6]. Therefore, the stochastic integral $\int_0^t H dB$ can be approximated for $t \leq u$ by the correspondingly modified Riemann sums.

For continuous martingales, a similar statement as in Theorem 5.9 holds provided the angle bracket process is absolutely continuous. Let $\mathcal{L}_a^2(0, u; M)$ denote the linear space of all product-measurable, (\mathcal{F}_t^P) adapted stochastic processes $(\omega, t) \mapsto H_t(\omega)$ such that

$$E \left[\int_0^u H_t^2 d\langle M \rangle_t \right] < \infty.$$

The corresponding space of equivalence classes w.r.t. $P_{\langle M \rangle}$ is denoted by $L_a^2(0, u; M)$.

Exercise (Admissible integrands w.r.t. martingales). Suppose that (M_t) is a continuous square integrable (\mathcal{F}_t) martingale. Show that if almost surely, $t \mapsto \langle M \rangle_t$ is absolutely continuous, then the closure $\overline{\mathcal{E}_u}$ of the elementary processes w.r.t. the $L^2(P_{\langle M \rangle})$ norm is given by

$$\overline{\mathcal{E}_u} = L_a^2(0, u; M).$$

5.3 Localization

Square-integrability of the integrand is still an assumption that we would like to avoid, since it is not always easy to verify or may even fail to hold. The key to extending

the class of admissible integrands further is localization, which enables us to define a stochastic integral w.r.t. a continuous martingale for any continuous adapted process. The price we have to pay is that for integrands that are not square integrable, the Itô integral is in general not a martingale, but only a local martingale.

Throughout this section we assume that M_t is a continuous square integrable martingale with absolutely continuous angle bracket process $\langle M \rangle_t$.

Local dependence on integrand and integrator

The approximations considered in the last section imply that the stochastic integral depends locally both on the integrand and on the integrator in the following sense:

Corollary 5.10. *Suppose that $T : \Omega \rightarrow [0, \infty]$ is a random variable, M, \widetilde{M} are square integrable martingales with absolutely continuous angle bracket processes $\langle M \rangle, \langle \widetilde{M} \rangle$, and H, \widetilde{H} are processes in $\mathcal{L}_a^2(0, \infty; M), \mathcal{L}_a^2(0, \infty; \widetilde{M})$ respectively, such that almost surely, $H_t = \widetilde{H}_t$ for any $t \in [0, T)$ and $M_t = \widetilde{M}_t$ for any $t \in [0, T]$. Then almost surely,*

$$\int_0^t H_s dM_s = \int_0^t \widetilde{H}_s d\widetilde{M}_s \quad \text{for any } t \in [0, T]. \quad (5.3.1)$$

Proof. We go through the same approximations as in the proof of Theorem 5.9 above:

- (1). Suppose first that H_t and \widetilde{H}_t are almost surely continuous and bounded, and there exists $t_0 \in \mathbb{R}_+$ such that $H_t = \widetilde{H}_t = 0$ for $t \geq t_0$. Let (π_n) be a sequence of partitions with $\text{mesh}(\pi_n) \rightarrow 0$. Then by Theorem 5.7,

$$\begin{aligned} \int_0^t H dM &= \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} H_s \cdot (M_{s' \wedge t} - M_s), & \text{and} \\ \int_0^t \widetilde{H} d\widetilde{M} &= \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} \widetilde{H}_s \cdot (\widetilde{M}_{s' \wedge t} - \widetilde{M}_s) \end{aligned}$$

with P -almost sure uniform convergence on finite time-intervals along a common subsequence. For $t \leq T$ the right-hand sides coincide, and thus (5.3.1) holds true.

- (2). Now suppose that H and \tilde{H} are bounded and $H_t = \tilde{H}_t = 0$ for $t \geq t_0$. Then the approximations

$$H_t^n = \int_0^{1/n} H_{t-\varepsilon} \psi_n(\varepsilon) d\varepsilon, \quad \tilde{H}_t^n = \int_0^{1/n} \tilde{H}_{t-\varepsilon} \psi_n(\varepsilon)$$

(with ψ_n defined as in the proof of Theorem 5.9 and $H_t := \tilde{H}_t := 0$ for $t < 0$) coincide for $t \leq T$. Hence by (1), on $\{t \leq T\}$,

$$\int_0^t H dM = \lim \int_0^t H^n dM = = \lim \int_0^t \tilde{H}^n d\tilde{M} = \int_0^t \tilde{H} d\tilde{M},$$

where the convergence holds again almost surely uniformly in t along a subsequence.

- (3). Finally, in the general case the assertion follows by approximating H and \tilde{H} by the bounded processes

$$H_t^n = ((H_t \wedge n) \vee (-n)) \cdot I_{[0,n]}(t), \quad \tilde{H}_t^n = ((\tilde{H}_t \wedge n) \vee (-n)) \cdot I_{[0,n]}(t).$$

□

Itô integrals for locally square-integrable integrands

Let M be a continuous square integrable martingale with absolutely continuous angle bracket process $\langle M \rangle$, and let $T : \Omega \rightarrow [0, \infty]$ be an (\mathcal{F}_t^P) stopping time. We will also be interested in the case where $T = \infty$. Let $\mathcal{L}_{a,\text{loc}}^2(0, T; M)$ denote the linear space consisting of all stochastic processes $(t, \omega) \mapsto H_t(\omega)$ defined for $t \in [0, T(\omega))$ such that the trivially extended process

$$\tilde{H}_t := \begin{cases} H_t & \text{for } t < T, \\ 0 & \text{for } t \geq T, \end{cases}$$

is product measurable in (t, ω) , adapted w.r.t. the filtration (\mathcal{F}_t^P) , and

$$t \mapsto H_t(\omega) \quad \text{is in } \mathcal{L}_{\text{loc}}^2([0, T(\omega)), d\langle M \rangle(\omega)) \quad \text{for } P\text{-a.e. } \omega. \quad (5.3.2)$$

Here for $u \in (0, \infty]$, the space $\mathcal{L}_{\text{loc}}^2([0, u], d\langle M \rangle(\omega))$ consists of all measurable functions $f : [0, u) \rightarrow [-\infty, \infty]$ such that $\int_0^s f(t)^2 d\langle M \rangle_t(\omega) < \infty$ for any $s \in (0, u)$. In particular, it contains all continuous functions.

From now on, we use the notation $H_t \cdot I_{\{t < T\}}$ for the trivial extension $(\tilde{H}_t)_{0 \leq t < \infty}$ of a process $(H_t)_{0 \leq t < T}$ beyond the stopping time T . Locally square integrable adapted processes allow for a localization by stopping times:

Lemma 5.11 (Localization by stopping). *If $(H_t)_{0 \leq t < T}$ is a process in $\mathcal{L}_{a, \text{loc}}^2(0, T; M)$ then there exists an increasing sequence $(T_n)_{n \in \mathbb{N}}$ of (\mathcal{F}_t^P) stopping times such that $T = \sup T_n$ almost surely, and*

$$H_t \cdot I_{\{t < T_n\}} \in \mathcal{L}_a^2(0, \infty; M) \quad \text{for any } n \in \mathbb{N}.$$

Proof. One easily verifies that the random variables T_n defined by

$$T_n := \inf \left\{ 0 \leq t < T : \int_0^t H_s^2 d\langle M \rangle_s \geq n \right\} \wedge T, \quad n \in \mathbb{N}, \quad (5.3.3)$$

are (\mathcal{F}_t^P) stopping times. Moreover, for almost every ω , the function $t \mapsto H_t(\omega)$ is in $\mathcal{L}_{\text{loc}}^2([0, T], d\langle M \rangle(\omega))$. Hence the function $t \mapsto \int_0^t H_s(\omega)^2 d\langle M \rangle_s$ is increasing and finite on $[0, T(\omega))$, and therefore $T_n(\omega) \nearrow T(\omega)$ as $n \rightarrow \infty$. Since T_n is an (\mathcal{F}_t^P) stopping time, the process $H_t \cdot I_{\{t < T_n\}}$ is (\mathcal{F}_t^P) -adapted, and by (5.3.3),

$$E \left[\int_0^\infty (H_s \cdot I_{\{s < T_n\}})^2 d\langle M \rangle_s \right] = E \left[\int_0^{T_n} H_s^2 d\langle M \rangle_s \right] \leq n \quad \text{for any } n.$$

□

A sequence of stopping times as in the lemma will also be called a **localizing sequence**. We can now extend the definition of the Itô integral to locally square-integrable adapted integrands:

Definition (Itô integral with locally square integrable integrand). *For a process $H \in \mathcal{L}_{a, \text{loc}}^2(0, T; M)$, the Itô stochastic integral w.r.t. the martingale M is defined for $t \in [0, T)$ by*

$$\int_0^t H_s dM_s := \int_0^t H_s \cdot I_{\{s < \hat{T}\}} dM_s \quad \text{for any } t \in [0, \hat{T}] \quad (5.3.4)$$

whenever \hat{T} is an (\mathcal{F}_t^P) stopping time such that $H_t \cdot I_{\{t < \hat{T}\}} \in \mathcal{L}_a^2(0, \infty; M)$.

Theorem 5.12. For $H \in \mathcal{L}_{a,loc}^2(0, T; M)$ the Itô integral $t \mapsto \int_0^t H_s dM_s$ is almost surely well defined by (5.3.4) as a continuous process on $[0, T)$.

Proof. We have to verify that the definition does not depend on the choice of the localizing stopping times. This is a direct consequence of Corollary 5.10: Suppose that \hat{T} and \tilde{T} are stopping times such that $H_t \cdot I_{\{t < \hat{T}\}}$ and $H_t \cdot I_{\{t < \tilde{T}\}}$ are both in $\mathcal{L}_a^2(0, \infty; M)$. Since the two trivially extended processes agree on $[0, \hat{T} \wedge \tilde{T})$, Corollary 5.10 implies that almost surely,

$$\int_0^t H_s \cdot I_{\{s < \hat{T}\}} dM_s = \int_0^t H_s \cdot I_{\{s < \tilde{T}\}} dM_s \quad \text{for any } t \in [0, \hat{T} \wedge \tilde{T}).$$

Hence, by Lemma 5.11, the stochastic integral is well defined on $[0, T)$. □

Stochastic integrals as local martingales

Itô integrals w.r.t. square integrable martingales are not necessarily martingales if the integrands are not square integrable. However, they are still local martingales in the sense of the definition stated below.

Definition (Predictable stopping time). An (\mathcal{F}_t^P) stopping time T is called *predictable* iff there exists an increasing sequence $(T_k)_{k \in \mathbb{N}}$ consisting of (\mathcal{F}_t^P) stopping times such that $T_k < T$ on $\{T \neq 0\}$ for any k , and $T = \sup T_k$.

Example (Hitting time of a closed set). The hitting time T_A of a closed set A by a continuous adapted process is predictable, as it can be approximated from below by the hitting times T_{A_k} of the neighbourhoods $A_k = \{x : \text{dist}(x, A) < 1/k\}$. On the other hand, the hitting time of an open set is not predictable in general.

Definition (Local martingale). Suppose that $T : \Omega \rightarrow [0, \infty]$ is a predictable stopping time. A stochastic process $M_t(\omega)$ defined for $0 \leq t < T(\omega)$ is called a **local martingale up to time T** , if and only if there exists an increasing sequence (T_k) of stopping times with $T = \sup T_k$ such that for any $k \in \mathbb{N}$, $T_k < T$ on $\{T > 0\}$, and the stopped process $(M_{t \wedge T_k})$ is a martingale for $t \in [0, \infty)$.

Recall that by the Optional Stopping Theorem, a continuous martingale stopped at a stopping time is again a martingale. Therefore, any continuous martingale $(M_t)_{t \geq 0}$ is a local martingale up to $T = \infty$. Even if (M_t) is assumed to be uniformly integrable, the converse implication fails to hold:

Exercise (A uniformly integrable local martingale that is not a martingale). Let $x \in \mathbb{R}^3$ with $x \neq 0$, and suppose that (B_t) is a three-dimensional Brownian motion with initial value $B_0 = x$. Prove that the process $M_t = 1/|B_t|$ is a uniformly integrable local martingale up to $T = \infty$, but (M_t) is not a martingale.

On the other hand, note that if (M_t) is a continuous local martingale up to $T = \infty$, and the family $\{M_{t \wedge T_k} : k \in \mathbb{N}\}$ is uniformly integrable for each fixed $t \geq 0$, then (M_t) is a martingale, because for $0 \leq s \leq t$

$$E[M_t | \mathcal{F}_s] = \lim_{k \rightarrow \infty} E[M_{t \wedge T_k} | \mathcal{F}_s] = \lim_{k \rightarrow \infty} M_{s \wedge T_k} = M_s$$

with convergence in L^1 .

As a consequence of the definition of the Itô integral by localization, we immediately obtain:

Theorem 5.13 (Itô integrals as local martingales). Suppose that T is a predictable stopping time w.r.t. (\mathcal{F}_t^P) . Then for any $H \in \mathcal{L}_{a,loc}^2(0, T; M)$, the Itô integral process $t \mapsto \int_0^t H_s dM_s$ is a continuous local martingale up to time T .

Proof. We can choose an increasing sequence (T_k) of stopping times such that $T_k < T$ on $\{T > 0\}$ and $H_t \cdot I_{\{t < T_k\}} \in \mathcal{L}_a^2(0, \infty; M)$ for any k . Then, by definition of the Itô integral,

$$\int_0^{t \wedge T_k} H_s dM_s = \int_0^{t \wedge T_k} H_s \cdot I_{\{s < T_k\}} dM_s \quad \text{almost surely for any } k \in \mathbb{N},$$

and the right-hand side is a continuous martingale in $M_c^2([0, \infty))$. □

The theorem shows that for a predictable (\mathcal{F}_t^P) stopping time T , the Itô map $H \mapsto \int_0^\bullet H dM$ extends to a linear map

$$\mathcal{J} : L_{\text{loc}}^2(0, T; M) \longrightarrow M_{c, \text{loc}}([0, T]),$$

where $L_{\text{loc}}^2(0, T; M)$ is the space of equivalence classes of processes in $\mathcal{L}_{\text{loc}}^2(0, T; M)$ that coincide for $P_{\langle M \rangle}$ -a.e. (ω, t) , and $M_{c, \text{loc}}([0, T])$ denotes the space of equivalence classes of continuous local (\mathcal{F}_t^P) martingales up to time T w.r.t. P -almost sure coincidence. Note that different notions of equivalence are used for the integrands and the integrals.

We finally observe that continuous local martingales (and hence stochastic integrals w.r.t. continuous martingales) can always be localized by a sequence of *bounded* martingales in $M_c^2([0, \infty))$:

Exercise (Localization by bounded martingales). Suppose that (M_t) is a continuous local martingale up to time T , and (T_k) is a localizing sequence of stopping times.

(1). Show that

$$\tilde{T}_k = T_k \wedge \inf\{t \geq 0 : |M_t| \geq k\} \wedge k$$

is another localizing sequence, and for all k , the stopped processes $(M_{t \wedge \tilde{T}_k})_{t \in [0, \infty)}$ are bounded martingales in $M_c^2([0, \infty))$.

(2). Show that if $T = \infty$ then $\hat{T}_k := \inf\{t \geq 0 : |M_t| \geq k\}$ is also a localizing sequence for M .

Approximation by Riemann-Itô sums

If the integrand (H_t) of a stochastic integral $\int H dB$ has continuous sample paths then local square integrability always holds, and the stochastic integral is a limit of Riemann-Itô sums: Let (π_n) be a sequence of partition of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$.

Theorem 5.14. *Suppose that T is a predictable stopping time, and $(H_t)_{0 \leq t < T}$ is a stochastic process defined for $t < T$. If the sample paths $t \mapsto H_t(\omega)$ are continuous on $[0, T(\omega))$ for any ω , and the trivially extended process $H_t \cdot I_{\{t < T\}}$ is (\mathcal{F}_t^P) adapted, then H is in $\mathcal{L}_{a,loc}^2(0, T; M)$, and for any $t \geq 0$,*

$$\int_0^t H_s dM_s = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} H_s \cdot (M_{s' \wedge t} - M_s) \quad \text{on } \{t < T\} \quad (5.3.5)$$

with convergence in probability.

Proof. Let $\lfloor t \rfloor_n = \max\{s \in \pi_n : s \leq t\}$ denote the next partition point below t . By continuity,

$$H_t \cdot I_{\{t < T\}} = \lim_{n \rightarrow \infty} H_{\lfloor t \rfloor_n} \cdot I_{\{t < T\}}.$$

Hence $(H_t \cdot I_{\{t < T\}})$ is (\mathcal{F}_t^P) adapted. It is also product-measurable, because

$$H_{\lfloor t \rfloor_n} \cdot I_{\{t < T\}} = \sum_{s \in \pi_n} H_s \cdot I_{\{s < T\}} \cdot I_{[s, s')}(t) \cdot I_{(0, \infty)}(T - t).$$

By continuity, $t \mapsto H_t(\omega)$ is locally bounded for every ω , and thus H is in $\mathcal{L}_{a,loc}^2(0, T; M)$. Moreover, suppose that (T_k) is a sequence of stopping times approaching T from below in the sense of the definition of a predictable stopping time given above. Then

$$\tilde{T}_k := T_k \wedge \inf\{t \geq 0 : |H_t| \geq k\}, \quad k \in \mathbb{N},$$

is a localizing sequence of stopping times with $H_t \cdot I_{\{t < T_k\}}$ in $\mathcal{L}_a^2(0, T; M)$ for any k , and $\tilde{T}_k \nearrow T$. Therefore, by definition of the Itô integral and by Theorem 5.7,

$$\begin{aligned} \int_0^t H_s dM_s &= \int_0^t H_s \cdot I_{\{s < \tilde{T}_k\}} dM_s = \int_0^t H_s \cdot I_{\{s \leq \tilde{T}_k\}} dM_s \\ &= \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} H_s \cdot (M_{s' \wedge t} - M_s) \quad \text{on } \{t \leq \tilde{T}_k\} \end{aligned}$$

w.r.t. convergence in probability. Since

$$P \left[\{t < T\} \setminus \bigcup_k \{t \leq \tilde{T}_k\} \right] = 0,$$

we obtain (5.3.5). □

Chapter 6

Itô's formula and pathwise integrals

Our approach to Itô's formula in this chapter follows that of [Föllmer: Stochastic Analysis, Vorlesungsskript Uni Bonn WS91/92]. We start with a heuristic derivation of the formula that will be the central topic of this chapter.

Suppose that $s \mapsto X_s$ is a function from $[0, t]$ to \mathbb{R} , and F is a smooth function on \mathbb{R} . If (π_n) is a sequence of partitions of the interval $[0, t]$ with $\text{mesh}(\pi_n) \rightarrow 0$ then by Taylor's theorem

$$F(X_{s'}) - F(X_s) = F'(X_s) \cdot (X_{s'} - X_s) + \frac{1}{2} F''(X_s) \cdot (X_{s'} - X_s)^2 + \text{higher order terms.}$$

Summing over $s \in \pi_n$ we obtain

$$F(X_t) - F(X_0) = \sum_{s \in \pi_n} F'(X_s) \cdot (X_{s'} - X_s) + \frac{1}{2} F''(X_s) \cdot (X_{s'} - X_s)^2 + \dots \quad (6.0.1)$$

We are interested in the limit of this formula as $n \rightarrow \infty$.

(a) Classical case, e.g. X continuously differentiable For $X \in C^1$ we have

$$\begin{aligned} X_{s'} - X_s &= \frac{dX_s}{ds}(s' - s) + O(|s - s'|^2), & \text{and} \\ (X_{s'} - X_s)^2 &= O(|s - s'|^2). \end{aligned}$$

Therefore, the second order terms can be neglected in the limit of (6.0.1) as $\text{mesh}(\pi_n) \rightarrow 0$. Similarly, the higher order terms can be neglected, and we obtain the limit equation

$$F(X_t) - F(X_0) = \int_0^t F'(X_s) dX_s, \quad (6.0.2)$$

or, in differential notation,

$$dF(X_t) = F'(X_t) dX_t, \quad (6.0.3)$$

Of course, (6.0.3) is just the chain rule of classical analysis, and (6.0.2) is the equivalent chain rule for Stieltjes integrals, cf. Section 6.1 below.

(b) X_t Brownian motion If (X_t) is a Brownian motion then

$$E[(X_{s'} - X_s)^2] = s' - s.$$

Summing these expectations over $s \in \pi_n$, we obtain the value t independently of n . This shows that the sum of the second order terms in (6.0.1) can not be neglected anymore. Indeed, as $n \rightarrow \infty$, a law of large numbers type result implies that we can almost surely replace the squared increments $(X_{s'} - X_s)^2$ in (6.0.1) asymptotically by their expectation values. The higher order terms are on average $O(|s' - s|^{3/2})$ whence their sum can be neglected. Therefore, in the limit of (6.0.1) as $n \rightarrow \infty$ we obtain the modified chain rule

$$F(X_t) - F(X_0) = \int_0^t F'(X_s) dX_s + \frac{1}{2} \int_0^t F''(X_s) ds \quad (6.0.4)$$

with probability one. The equation (6.0.4) is the basic version of Itô's celebrated formula.

In Section 6.1, we will first introduce Stieltjes integrals and the chain rule from Stieltjes calculus systematically. In Section 6.2 we prove a general version of Itô's formula for continuous functions with finite quadratic variation in dimension one. Here the setup and the proof are still purely deterministic. As an aside we obtain a pathwise definition for stochastic integrals involving only a single one-dimensional process due to Föllmer. After computing the quadratic variation of Brownian motion in Section 6.3, we consider first consequences of Itô's formula for Brownian motions and continuous martingales. Section 6.4 contains extensions to the multivariate and time-dependent case, as well as further applications.

6.1 Stieltjes integrals and chain rule

In this section, we define Lebesgue-Stieltjes integrals w.r.t. deterministic functions of finite variation, and prove a corresponding chain rule. The resulting calculus can then be applied path by path to stochastic processes with sample paths of finite variation.

Lebesgue-Stieltjes integrals

Fix $u \in (0, \infty]$, and suppose that $t \mapsto A_t$ is a right-continuous and non-decreasing function on $[0, u)$. Then $A_t - A_0$ is the distribution function of the positive measure μ on $(0, u)$ determined uniquely by

$$\mu_A[(s, t]] = A_t - A_s \quad \text{for any } 0 \leq s \leq t < u.$$

Therefore, we can define integrals of type $\int_s^t H_s dA_s$ as Lebesgue integrals w.r.t. the measure μ_A . We extend μ trivially to the interval $[0, u)$, so $\mathcal{L}_{\text{loc}}^1([0, u), \mu_A)$ is the space of all functions $H : [0, u) \rightarrow \mathbb{R}$ that are integrable w.r.t. μ_A on any interval $(0, t)$ with $t < u$. Then for any $u \in [0, \infty]$ and any function $H \in \mathcal{L}_{\text{loc}}^1([0, u), \mu_A)$, the **Lebesgue-Stieltjes integral of H w.r.t. A** is defined by

$$\int_s^t H_r dA_r := \int H_r \cdot I_{(s, t]}(r) \mu_A(dr) \quad \text{for } 0 \leq s \leq t < u.$$

It is easy to verify that the definition is consistent, i.e., varying u does not change the definition of the integrals, and that $t \mapsto \int_0^t H_r dA_r$ is again a right-continuous function.

For an arbitrary right-continuous function $A : [0, u) \rightarrow \mathbb{R}$, the (first order) variation of A on an interval $[0, t)$ is defined by

$$V_t^{(1)}(A) := \sup_{\pi} \sum_{s \in \pi} |A_{s' \wedge t} - A_{s \wedge t}| \quad \text{for } 0 \leq t < u,$$

where the supremum is over all partitions π of \mathbb{R}_+ . The function $t \mapsto A_t$ is said to be **(locally) of finite variation** on the interval $[0, u)$ iff $V_t^{(1)}(A) < \infty$ for any $t \in [0, u)$.

Any right-continuous function of finite variation can be written as the difference of two non-decreasing right-continuous functions. In fact, we have

$$A_t = A_t^{\nearrow} - A_t^{\searrow} \quad (6.1.1)$$

with

$$A_t^{\nearrow} = \sup_{\pi} \sum_{s \in \pi} (A_{s' \wedge t} - A_{s \wedge t})^+ = \frac{1}{2}(V_t^{(1)}(A) + A_t), \quad (6.1.2)$$

$$A_t^{\searrow} = \sup_{\pi} \sum_{s \in \pi} (A_{s' \wedge t} - A_{s \wedge t})^- = \frac{1}{2}(V_t^{(1)}(A) - A_t). \quad (6.1.3)$$

Exercise. Prove that if A_t is right-continuous and is locally of finite variation on $[0, u)$ then the functions $V_t^{(1)}(A)$, A_t^{\nearrow} and A_t^{\searrow} are all right-continuous and non-decreasing for $t < u$.

Remark (Hahn-Jordan decomposition). The functions $A_t^{\nearrow} - A_0^{\nearrow}$ and $A_t^{\searrow} - A_0^{\searrow}$ are again distribution functions of positive measures μ_A^+ and μ_A^- on $(0, u)$. Correspondingly, $A_t - A_0$ is the distribution function of the signed measure

$$\mu_A[B] := \mu_A^+[B] - \mu_A^-[B], \quad B \in \mathcal{B}(0, u), \quad (6.1.4)$$

and $V_t^{(1)}$ is the distribution of the measure $|\mu_A| = \mu_A^+ + \mu_A^-$. It is a consequence of (6.1.5) and (6.1.6) that the measures μ_A^+ and μ_A^- are singular, i.e., the mass is concentrated on disjoint sets S^+ and S^- . The decomposition (6.1.7) is hence a particular case of the Hahn-Jordan decomposition of a signed measure μ of finite variation into a positive and a negative part, and the measure $|\mu|$ is the total variation measure of μ , cf. e.g. [Alt: Lineare Funktionalanalysis].

We can now apply (6.1.1) to define Lebesgue-Stieltjes integrals w.r.t. functions of finite variation. A function is integrable w.r.t. a signed measure μ if and only if it is integrable w.r.t. both the positive part μ^+ and the negative part μ^- . The Lebesgue integral w.r.t. μ is then defined as the difference of the Lebesgue integrals w.r.t. μ^+ and μ^- . Correspondingly, we define the Lebesgue-Stieltjes integral w.r.t. a function A_t of finite variation as the Lebesgue integral w.r.t. the associated signed measure μ_A :

Definition. Suppose that $t \mapsto A_t$ is right-continuous and locally of finite variation on $[0, u)$. Then the **Lebesgue-Stieltjes integral w.r.t. A** is defined by

$$\int_s^t H_r dA_r := \int H_r \cdot I_{(s,t]}(r) dA_r^{\nearrow} - \int H_r \cdot I_{(s,t]}(r) dA_r^{\searrow}, \quad 0 \leq s \leq t < u,$$

for any function $H \in \mathcal{L}_{loc}^1((0, u), |dA|)$ where

$$\mathcal{L}_{loc}^1((0, u), |dA|) := \mathcal{L}_{loc}^1((0, u), dA^{\nearrow}) \cap \mathcal{L}_{loc}^1((0, u), dA^{\searrow})$$

is the intersection of the local \mathcal{L}^1 spaces w.r.t. the positive measures $dA^{\nearrow} = \mu_A^+$ and $dA^{\searrow} = \mu_A^-$ on $[0, u)$, or, equivalently, the local \mathcal{L}^1 space w.r.t. the total variation measure $|dA| = |\mu_A|$.

Remark. (1). *Simple integrands:* If $H_t = \sum_{i=0}^{n-1} c_i \cdot I_{(t_i, t_{i+1}]}$ is a step function with $0 \leq t_0 < t_1 < \dots < t_n < u$ and $c_0, c_1, \dots, c_{n-1} \in \mathbb{R}$ then

$$\int_0^t H_s dA_s = \sum_{i=0}^{n-1} c_i \cdot (A_{t_{i+1} \wedge t} - A_{t_i \wedge t}).$$

(2). *Continuous integrands; Riemann-Stieltjes integral:* If $H : [0, u) \rightarrow \mathbb{R}$ is a continuous function then the Stieltjes integral can be approximated by Riemann sums:

$$\int_0^t H_s dA_s = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} H_s \cdot (A_{s' \wedge t} - A_s), \quad t \in [0, u),$$

for any sequence (π_n) of partitions of \mathbb{R}_+ such that $\text{mesh}(\pi_n) \rightarrow 0$. For the proof note that the step functions

$$H_r^n = \sum_{\substack{s \in \pi_n \\ s < r}} H_s \cdot I_{(s, s']}(r), \quad r \in [0, u),$$

converge to H_r pointwise on $(0, u)$ by continuity. Moreover, again by continuity, H_r is locally bounded on $[0, u)$, and hence the sequence H_r^n is locally uniformly bounded. Therefore,

$$\int H_r^n I_{(0,t]}(r) dA_r \longrightarrow \int H_r I_{(0,t]}(r) dA_r$$

for any $t < u$ by the dominated convergence theorem.

- (3). *Absolutely continuous integrators:* If A_t is an absolutely continuous function on $[0, u)$ then A_t has locally finite variation

$$V_t^{(1)}(A) = \int_0^t |A'_s| ds < \infty \quad \text{for } t \in [0, u).$$

The signed measure μ_A with distribution function $A_t - A_0$ is then absolutely continuous w.r.t. Lebesgue measure with Radon-Nikodym density

$$\frac{d\mu_A}{dt}(t) = A'_t \quad \text{for almost every } t \in [0, u).$$

Therefore,

$$\mathcal{L}_{\text{loc}}^1([0, u), |dA|) = \mathcal{L}_{\text{loc}}^1([0, u), |A'|dt),$$

and the Lebesgue-Stieltjes integral of a locally integrable function H is given by

$$\int_0^t H_s dA_s = \int_0^t H_s A'_s ds \quad \text{for } t \in [0, u).$$

In the applications that we are interested in, the integrand will mostly be continuous, and the integrator absolutely continuous. Hence Remarks (2) and (3) above apply.

The chain rule in Stieltjes calculus

We are now able to prove Itô's formula in the special situation where the integrator has finite variation. In this case, the second order correction disappears, and Itô's formula reduces to the classical chain rule from Stieltjes calculus:

Theorem 6.1 (Fundamental Theorem of Stieltjes Calculus). *Suppose that $A : [0, u) \rightarrow \mathbb{R}$ is a continuous function of locally finite variation. Then for any $F \in C^2(\mathbb{R})$,*

$$F(A_t) - F(A_0) = \int_0^t F'(A_s) dA_s \quad \forall t \in [0, u). \quad (6.1.5)$$

Proof. Let $t \in [0, u)$ be given. Choose a sequence of partitions (π_n) of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$, and let

$$\delta A_s := A_{s' \wedge t} - A_{s \wedge t} \quad \text{for } s \in \pi_n,$$

where, as usual, s' denotes the next partition point. By Taylor's formula, for $s \in \pi_n$ with $s < t$ we have

$$F(A_{s' \wedge t}) - F(A_s) = F'(A_s) \delta A_s + \frac{1}{2} F''(Z_s) \cdot (\delta A_s)^2,$$

where Z_s is an intermediate value between A_s and $A_{s' \wedge t}$. Summing over $s \in \pi_n$, we obtain

$$F(A_t) - F(A_0) = \sum_{\substack{s \in \pi_n \\ s < t}} F'(A_s) \delta A_s + \frac{1}{2} \sum_{\substack{s \in \pi_n \\ s < t}} F''(Z_s) (\delta A_s)^2. \quad (6.1.6)$$

As $n \rightarrow \infty$, the first (Riemann) sum converges to the Stieltjes integral $\int_0^t F'(A_s) dA_s$ by continuity of $F'(A_s)$, cf. Remark (2) above.

To see that the second sum converges to zero, note that the range of the continuous function A restricted to $[0, t]$ is a bounded interval. Since F'' is continuous by assumption, F'' is bounded on this range by a finite constant c . As Z_s is an intermediate value between A_s and $A_{s' \wedge t}$, we obtain

$$\left| \sum_{\substack{s \in \pi_n \\ s < t}} F''(Z_s) (\delta A_s)^2 \right| \leq c \cdot \sum_{\substack{s \in \pi_n \\ s < t}} (\delta A_s)^2 \leq c \cdot V_t^{(1)}(A) \cdot \sup_{\substack{s \in \pi_n \\ s < t}} |\delta A_s|.$$

Since $V_t^{(1)}(A) < \infty$, and A is a uniformly continuous function on $[0, t]$, the right hand side converges to 0 as $n \rightarrow \infty$. Hence we obtain (6.1.5) in the limit of (6.1.6) as $n \rightarrow \infty$. \square

To see that (6.1.5) can be interpreted as a chain rule, we write the equation in differential form:

$$dF(A) = F'(A)dA. \quad (6.1.7)$$

In general, the equation (6.1.7) is to be understood mathematically only as an abbreviation for the integral equation (6.1.5). For intuitive arguments, the differential notation is obviously much more attractive than the integral form of the equation. However, for the differential form to be useful at all, we should be able to multiply the equation (6.1.7) by another function, and still obtain a valid equation. This is indeed possible due to the next result, which states briefly that if $dI = H dA$ then also $G dI = GH dA$:

Theorem 6.2 (Stieltjes integrals w.r.t. Stieltjes integrals). *Suppose that $I_s = \int_0^s H_r dA_r$ where $A : [0, u) \rightarrow \mathbb{R}$ is a function of locally finite variation, and $H \in \mathcal{L}_{loc}^1([0, u), |dA|)$. Then the function $s \mapsto I_s$ is again right continuous with locally finite variation $V_t^{(1)}(I) \leq \int_0^t |H| |dA| < \infty$, and, for any function $G \in \mathcal{L}_{loc}^1([0, u), |dI|)$,*

$$\int_0^t G_s dI_s = \int_0^t G_s H_s dA_s \quad \text{for } t \in [0, u). \quad (6.1.8)$$

Proof. Right continuity of I_t and the upper bound for the variation are left as an exercise.

We now use Riemann sum approximations to prove (6.1.8) if G is continuous. For a partition $0 = t_0 < t_1 < \dots < t_k = t$, we have

$$\sum_{i=0}^{n-1} G_{t_i} (I_{t_{i+1}} - I_{t_i}) = \sum_{i=0}^{n-1} G_{t_i} \cdot \int_{t_i}^{t_{i+1}} H_s dA_s = \int_0^t G_{[s]} H_s dA_s$$

where $\lfloor s \rfloor$ denotes the largest partition point $\leq s$. Choosing a sequence (π_n) of partitions with $\text{mesh}(\pi_n) \rightarrow 0$, the integral on the right hand side converges to the Lebesgue-Stieltjes integral $\int_0^t G_s H_s dA_s$ by continuity of G and the dominated convergence theorem, whereas the Riemann sum on the left hand side converges to $\int_0^t G_s dI_s$. Hence (6.1.8) holds for continuous G . The equation for general $G \in \mathcal{L}_{\text{loc}}^1([0, u], |dI|)$ follows then by standard arguments. \square

6.2 Quadratic variation and Itô's formula

Our next goal is to derive a generalization of the chain rule from Stieltjes calculus to continuous functions that are not of finite variation. Examples of such functions are typical sample paths of Brownian motion. As pointed out above, an additional term will appear in the chain rule in this case.

Quadratic variation

Consider once more the approximation (6.1.6) that we have used to prove the fundamental theorem of Stieltjes calculus. We would like to identify the limit of the last sum $\sum_{s \in \pi_n} F''(Z_s)(\delta A_s)^2$ when A has unfinite variation on finite intervals. For $F'' = 1$ this limit is called the quadratic variation of A if it exists:

Definition. Let $u \in (0, \infty]$ and let (π_n) be a sequence of partitions of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$. The **quadratic variation** $[X]_t$ of a continuous function $X : [0, u) \rightarrow \mathbb{R}$ w.r.t. the sequence (π_n) is defined by

$$[X]_t = \lim_{n \rightarrow \infty} \sum_{s \in \pi_n} (X_{s' \wedge t} - X_{s \wedge t})^2 \quad \text{for } t \in [0, u)$$

whenever the limit exists.

WARNINGS (Dependence on partition, classical 2-variation)

- (1). The quadratic variation should not be confused with the classical 2-variation defined by

$$V_t^{(2)}(X) := \sup_{\pi} \sum_{s \in \pi} |X_{s' \wedge t} - X_{s \wedge t}|^2$$

where the supremum is over all partitions π . The classical 2-variation $V_t^{(2)}(X)$ is strictly positive for any function X that is not constant on $[0, t]$ whereas $[X]_t$ vanishes in many cases, cf. Example (1) below.

- (2). In general, the quadratic variation may depend on the sequence of partitions considered. See however Examples (1) and (3) below.

Example. (1). *Functions of finite variation:* For any continuous function $A : [0, u) \rightarrow \mathbb{R}$ of locally finite variation, the quadratic variation along (π_n) vanishes:

$$[A]_t = 0 \quad \text{for any } t \in [0, u).$$

In fact, for $\delta A_s = A_{s' \wedge t} - A_{s \wedge t}$ we have

$$\sum_{s \in \pi_n} |\delta A_s|^2 \leq V_t^{(1)}(A) \cdot \sup_{\substack{s \in \pi_n \\ s < t}} |\delta A_s| \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

by uniform continuity and since $V_t^{(1)}(A) < \infty$.

- (2). *Perturbations by functions of finite variation:* If the quadratic variation $[X]_t$ of X w.r.t. (π_n) exists, then $[X + A]_t$ also exists, and

$$[X + A]_t = [X]_t.$$

This holds since

$$\sum |\delta(X + A)|^2 = \sum (\delta X)^2 + 2 \sum \delta X \delta A + \sum (\delta A)^2,$$

and the last two sums converge to 0 as $\text{mesh}(\pi_n) \rightarrow 0$ by Example (1) and the Cauchy-Schwarz inequality.

- (3). *Brownian motion:* If $(B_t)_{t \geq 0}$ is a one-dimensional Brownian motion then P -almost surely,

$$[B]_t = t \quad \text{for all } t \geq 0$$

w.r.t. any *fixed* sequence (π_n) of partitions such that $\text{mesh}(\pi_n) \rightarrow 0$, cf. Theorem 6.6 below.

- (4). *Itô processes*: If $I_t = \int_0^t H_s dB_s$ is the stochastic integral of a process $H \in \mathcal{L}_{a,\text{loc}}^2(0, \infty)$ w.r.t. Brownian motion then almost surely, the quadratic variation w.r.t. a fixed sequence of partitions is

$$[I]_t = \int_0^t H_s^2 ds \quad \text{for all } t \geq 0.$$

- (5). *Continuous martingales*: $[M]$ exists and is almost surely finite, see below.

Note that in Examples (3), (4) and (5), the exceptional sets may depend on the sequence (π_n) . If it exists, the quadratic variation $[X]_t$ is a non-decreasing function in t . In particular, Stieltjes integrals w.r.t. $[X]$ are well-defined provided $[X]$ is right continuous.

Lemma 6.3. *Suppose that $X : [0, u) \rightarrow \mathbb{R}$ is a continuous function. If the quadratic variation $[X]_t$ along (π_n) exists for $t \in [0, u)$, and $t \mapsto [X]_t$ is continuous then*

$$\sum_{\substack{s \in \pi_n \\ s < t}} H_s \cdot (X_{s' \wedge t} - X_s)^2 \quad \longrightarrow \quad \int_0^t H_s d[X]_s \quad \text{as } n \rightarrow \infty \quad (6.2.1)$$

for any continuous function $H : [0, u) \rightarrow \mathbb{R}$ and any $t \geq 0$.

Remark. Heuristically, the assertion of the lemma says that

$$\left\langle \int H d[X] = \int H (dX)^2 \right\rangle,$$

i.e., the infinitesimal increments of the quadratic variation are something like squared infinitesimal increments of X . This observation is crucial for controlling the second order terms in the Taylor expansion used for proving the Itô-Doeblin formula.

Proof. The sum on the left-hand side of (6.2.1) is the integral of H w.r.t. the finite positive measure

$$\mu_n := \sum_{\substack{s \in \pi_n \\ s < t}} (X_{s' \wedge t} - X_s)^2 \cdot \delta_s$$

on the interval $[0, t)$. The distribution function of μ_n is

$$F_n(u) = : \sum_{\substack{s \in \pi_n \\ s \leq t}} (X_{s' \wedge t} - X_s)^2, \quad u \in [0, t].$$

As $n \rightarrow \infty$, $F_n(u) \rightarrow [X]_u$ for any $u \in [0, t]$ by continuity of X . Since $[X]_u$ is a continuous function of u , convergence of the distribution functions implies weak convergence of the measures μ_n to the measure $d[X]$ on $[0, t)$ with distribution function $[X]$. Hence,

$$\int H_s \mu_n(ds) \longrightarrow \int H_s d[X]_s \quad \text{as } n \rightarrow \infty$$

for any continuous function $H : [0, t] \rightarrow \mathbb{R}$. □

Itô's formula and pathwise integrals in \mathbb{R}^1

We are now able to complete the proof of the following purely deterministic (pathwise) version of the one-dimensional Itô formula going back to [Föllmer: Calcul d'Itô sans probabilités, Sémin. Prob XV, LNM850XXX]:

Theorem 6.4 (Itô's formula without probability). *Suppose that $X : [0, u) \rightarrow \mathbb{R}$ is a continuous function with continuous quadratic variation $[X]$ w.r.t. (π_n) . Then for any function F that is C^2 in a neighbourhood of $X([0, u))$, and for any $t \in [0, u)$, the Itô integral*

$$\int_0^t F'(X_s) dX_s = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} F'(X_s) \cdot (X_{s' \wedge t} - X_s) \quad (6.2.2)$$

exists, and Itô's formula

$$F(X_t) - F(X_0) = \int_0^t F'(X_s) dX_s + \frac{1}{2} \int_0^t F''(X_s) d[X]_s \quad (6.2.3)$$

holds. In particular, if the quadratic variation $[X]$ does not depend on (π_n) then the Itô integral (6.2.2) does not depend on (π_n) either.

Note that the theorem **implies the existence** of $\int_0^t f(X_s) dX_s$ for any function $f \in C^1(\mathbb{R})$! Hence this type of Itô integrals can be defined in a purely deterministic way without relying on the Itô isometry. Unfortunately, the situation is more complicated in higher dimensions, cf. ?? below.

Proof. Fix $t \in [0, u]$ and $n \in \mathbb{N}$. As before, for $s \in \pi_n$ we set $\delta X_s = X_{s' \wedge t} - X_{s \wedge t}$ where s' denotes the next partition point. Then as above we have

$$F(X_t) - F(X_0) = \sum_{\substack{s \in \pi_n \\ s < t}} F'(X_s) \delta X_s + \frac{1}{2} \sum_{\substack{s \in \pi_n \\ s < t}} F''(Z_s^{(n)}) (\delta X_s)^2 \quad (6.2.4)$$

$$= \sum_{\substack{s \in \pi_n \\ s < t}} F'(X_s) \delta X_s + \frac{1}{2} \sum_{\substack{s \in \pi_n \\ s < t}} F''(X_s) (\delta X_s)^2 + \sum_{\substack{s \in \pi_n \\ s < t}} R_s^{(n)}, \quad (6.2.5)$$

where $Z_s^{(n)}$ is an intermediate point between X_s and $X_{s' \wedge t}$, and $R_s^{(n)} := \frac{1}{2} (F''(Z_s^{(n)}) - F''(X_s)) \cdot (\delta X_s)^2$. As $n \rightarrow \infty$, the second sum on the right hand side of (6.2.4) converges to $\int_0^t F''(X_s) d[X]_s$ by Lemma 6.3. We claim that the sum of the remainders $R_s^{(n)}$ converges to 0. To see this note that $Z_s^{(n)} = X_r$ for some $r \in [s, s' \wedge t]$, whence

$$|R_s^{(n)}| = |F''(Z_s^{(n)}) - F''(X_s)| \cdot (\delta X_s)^2 \leq \frac{1}{2} \varepsilon_n (\delta X_s)^2,$$

where

$$\varepsilon_n := \sup_{\substack{a, b \in [0, t] \\ |a-b| \leq \text{mesh}(\pi_n)}} |F''(X_a) - F''(X_b)|.$$

As $n \rightarrow \infty$, ε_n converges to 0 by uniform continuity of $F'' \circ X$ on the interval $[0, t]$.

Thus

$$\sum |R_s^{(n)}| \leq \frac{1}{2} \varepsilon_n \sum_{\substack{s \in \pi_n \\ s < t}} (\delta X_s)^2 \rightarrow 0 \quad \text{as well,}$$

because the sum of the squared increments converges to the finite quadratic variation $[X]_t$.

We have shown that all the terms on the right hand side of (6.2.4) except the first

Riemann-Itô sum converge as $n \rightarrow \infty$. Hence, by (6.2.4), the limit $\int_0^t F'(X_s) dX_s$ of the Riemann Itô sums also exists, and the limit equation (6.2.2) holds. \square

Remark. (1). In differential notation, we obtain the Itô chain rule

$$dF(X) = F'(X) dX + \frac{1}{2}F''(X) d[X]$$

which includes a second order correction term due to the quadratic variation. A justification for the differential notation is given in Section ??.

(2). For functions X with $[X] = 0$ we recover the classical chain rule $dF(X) = F'(X) dX$ from Stieltjes calculus as a particular case of Itô's formula.

Example. (1). *Exponentials:* Choosing $F(x) = e^x$ in Itô's formula, we obtain

$$e^{X_t} - e^{X_0} = \int_0^t e^{X_s} dX_s + \frac{1}{2} \int_0^t e^{X_s} d[X]_s,$$

or, in differential notation,

$$de^X = e^X dX + \frac{1}{2}e^X d[X].$$

Thus e^X does *not* solve the Itô differential equation

$$dZ = Z dX \tag{6.2.6}$$

if $[X] \neq 0$. An appropriate renormalization is required instead. We will see below that the correct solution of (6.2.6) is given by

$$Z_t = \exp(X_t - [X]/2),$$

cf. the first example below Theorem 6.18.

(2). *Polynomials:* Similarly, choosing $F(x) = x^n$ for some $n \in \mathbb{N}$, we obtain

$$dX^n = nX^{n-1} dX + \frac{n(n-1)}{2}X^{n-2} [X].$$

Again, X^n does not solve the equation $dX^n = nX^{n-1} dX$. Here, the appropriate renormalization leads to the Hermite polynomials : $X :^n$, cf. the second example below Theorem 6.18.

The chain rule for anticipative integrals

The form of the second order correction term appearing in Itô's formula depends crucially on choosing non-anticipative Riemann sum approximations. For limits of anticipative Riemann sums, we obtain different correction terms, and hence also different notions of integrals.

Theorem 6.5. *Suppose that $X : [0, u) \rightarrow \mathbb{R}$ is continuous with continuous quadratic variation $[X]$ along (π_n) . Then for any function F that is C^2 in a neighbourhood of $X([0, u))$ and for any $t \geq 0$, the **backward Itô integral***

$$\int_0^t F'(X_s) \hat{d}X_s := \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} F'(X_{s' \wedge t}) \cdot (X_{s' \wedge t} - X_s),$$

and the **Stratonovich integral**

$$\int_0^t F'(X_s) \circ dX_s := \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} \frac{1}{2} (F'(X_s) + F'(X_{s' \wedge t})) \cdot (X_{s' \wedge t} - X_s)$$

exist, and

$$F(X_t) - F(X_0) = \int_0^t F'(X_s) \hat{d}X_s - \frac{1}{2} \int_0^t F''(X_s) d[X]_s \quad (6.2.7)$$

$$= \int_0^t F'(X_s) \circ dX_s. \quad (6.2.8)$$

Proof. The proof of the backward Itô formula (6.2.7) is completely analogous to that of Itô's formula. The Stratonovich formula (6.2.8) follows by averaging the Riemann sum approximations to the forward and backward Itô rule. \square

Note that Stratonovich integrals satisfy the classical chain rule

$$\circ dF(X) = F'(X) \circ dX.$$

This makes them very attractive for various applications. For example, in stochastic differential geometry, the chain rule is of fundamental importance to construct stochastic processes that stay on a given manifold. Therefore, it is common to use Stratonovich instead of Itô calculus in this context, cf. the corresponding example in the next section. On the other hand, Stratonovich calculus has a significant disadvantage against Itô calculus: The Stratonovich integrals

$$\int_0^t H_s \circ dB_s = \lim_{n \rightarrow \infty} \sum \frac{1}{2} (H_s + H_{s' \wedge t}) (B_{s' \wedge t} - B_s)$$

w.r.t. Brownian motion typically are not martingales, because the coefficients $\frac{1}{2}(H_s + H_{s' \wedge t})$ are not predictable.

6.3 Itô's formula for Brownian motion and martingales

Our next aim is to compute the quadratic variation and to state Itô's formula for typical sample paths of Brownian motion. More generally, we will show that the quadratic variation exists almost surely for continuous local martingales.

Let (π_n) be a sequence of partitions of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$. We note first that for any function $t \mapsto X_t$ the identity

$$X_t^2 - X_0^2 = \sum_{\substack{s \in \pi_n \\ s < t}} (X_{s' \wedge t}^2 - X_s^2) = V_t^n + 2I_t^n \quad (6.3.1)$$

with

$$V_t^n = \sum_{\substack{s \in \pi_n \\ s < t}} (X_{s' \wedge t} - X_s)^2 \quad \text{and} \quad I_t^n = \sum_{\substack{s \in \pi_n \\ s < t}} X_s \cdot (X_{s' \wedge t} - X_s)$$

holds. The equation (6.3.1) is a discrete approximation of Itô's formula for the function $F(x) = x^2$. The remainder terms in the approximation vanish in this particular case.

Note that by (6.3.1), the quadratic variation $[X]_t = \lim_{n \rightarrow \infty} V_t^n$ exists if and only if the Riemann sum approximations I_t^n to the Itô integral $\int_0^t X_s dX_s$ converge:

$$\exists [X]_t = \lim_{n \rightarrow \infty} V_t^n \quad \iff \quad \exists \int_0^t X_s dX_s = \lim_{n \rightarrow \infty} I_t^n.$$

Now suppose that (X_t) is a continuous martingale with $E[X_t^2] < \infty$ for any $t \geq 0$. Then the Riemann sum approximations (I_t^n) are continuous martingales for any $n \in \mathbb{N}$. Therefore, by the maximal inequality, for a given $u > 0$, the processes (I_t^n) and (V_t^n) converge uniformly for $t \in [0, u]$ in $L^2(P)$ if and only if the random variables I_u^n or V_u^n respectively converge in $L^2(P)$.

Quadratic variation of Brownian motion

For the sample paths of a Brownian motion B , the quadratic variation $[B]$ exists almost surely along any *fixed* sequence of partitions (π_n) with $\text{mesh}(\pi_n) \rightarrow 0$, and $[B]_t = t$ a.s. In particular, $[B]$ is a *deterministic* function that does not depend on (π_n) . The reason is a law of large numbers type effect when taking the limit of the sum of squared increments as $n \rightarrow \infty$.

Theorem 6.6 (P. Lévy). *If (B_t) is a one-dimensional Brownian motion on (Ω, \mathcal{A}, P) then as $n \rightarrow \infty$*

$$\sup_{t \in [0, u]} \left| \sum_{\substack{s \in \pi_n \\ s < t}} (B_{s' \wedge t} - B_s)^2 - t \right| \longrightarrow 0 \quad P\text{-a.s. and in } L^2(\Omega, \mathcal{A}, P) \quad (6.3.2)$$

for any $u \in (0, \infty)$, and for each sequence (π_n) of partitions of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$.

Warning. (1). Although the almost sure limit in (6.3.2) does not depend on the sequence (π_n) , the exceptional set may depend on the chosen sequence!

(2). The classical quadratic variation $V_t^{(2)}(B) = \sup_{\pi} \sum_{s \in \pi} (\delta B_s)^2$ is almost surely infinite for any $t \geq 0$. The classical p -variation is almost surely finite if and only if $p > 2$.

Proof. (1). L^2 -convergence for fixed t : As usual, the proof of L^2 convergence is comparatively simple. For $V_t^n = \sum_{s \in \pi_n} (\delta B_s)^2$ with $\delta B_s = B_{s' \wedge t} - B_{s \wedge t}$, we have

$$\begin{aligned} E[V_t^n] &= \sum_{s \in \pi_n} E[(\delta B_s)^2] = \sum_{s \in \pi_n} \delta s = t, \quad \text{and} \\ \text{Var}[V_t^n] &= \sum_{s \in \pi_n} \text{Var}[(\delta B_s)^2] = \sum_{s \in \pi_n} E[(\delta B_s)^2 - \delta s]^2 \\ &= E[(Z^2 - 1)^2] \cdot \sum_{s \in \pi_n} (\delta s)^2 \leq \text{const.} \cdot t \cdot \text{mesh}(\pi_n) \end{aligned}$$

where Z is a standard normal random variable. Hence, as $n \rightarrow \infty$,

$$V_t^n - t = V_t^n - E[V_t^n] \rightarrow 0 \quad \text{in } L^2(\Omega, \mathcal{A}, P).$$

Moreover, by (6.3.1), $V_t^n - V_t^m = I_t^n - I_t^m$ is a continuous martingale for any $n, m \in \mathbb{N}$. Therefore, the maximal inequality yields uniform convergence of V_t^n to t for t in a finite interval in the $L^2(P)$ sense.

(2). *Almost sure convergence if $\sum \text{mesh}(\pi_n) < \infty$* : Similarly, by applying the maximal inequality to the process $V_t^n - V_t^m$ and taking the limit as $m \rightarrow \infty$, we obtain

$$P \left[\sup_{t \in [0, u]} |V_t^n - t| > \varepsilon \right] \leq \frac{2}{\varepsilon^2} E[(V_t^n - t)^2] \leq \text{const.} \cdot t \cdot \text{mesh}(\pi_n)$$

for any given $\varepsilon > 0$ and $u \in (0, \infty)$. If $\sum \text{mesh}(\pi_n) < \infty$ then the sum of the probabilities is finite, and hence $\sup_{t \in [0, u]} |V_t^n - t| \rightarrow 0$ almost surely by the Borel-Cantelli Lemma.

(3). *Almost sure convergence if $\sum \text{mesh}(\pi_n) = \infty$* : In this case, almost sure convergence can be shown by the backward martingale convergence theorem. We refer to Proposition 2.12 in [Revuz, YorXXX], because for our purposes almost sure convergence w.r.t arbitrary sequences of partitions is not essential.

□

Itô's formula for Brownian motion

By Theorem 6.6, we can apply Theorem 6.7 to almost every sample path of a one-dimensional Brownian motion (B_t) :

Theorem 6.7 (Itô's formula for Brownian motion). *Suppose that $F \in C^2(I)$ where $I \subseteq \mathbb{R}$ be an open interval. Then almost surely,*

$$F(B_t) - F(B_0) = \int_0^t F'(B_s) dB_s + \frac{1}{2} \int_0^t F''(B_s) ds \quad \text{for all } t < T, \quad (6.3.3)$$

where $T = \inf\{t \geq 0 : B_t \notin I\}$ is the first exit time from I .

Proof. For almost every ω , the quadratic variation of $t \mapsto B_t(\omega)$ along a fixed sequence of partitions is t . Moreover, for any $r < T(\omega)$, the function F is C^2 on a neighbourhood of $\{B_t(\omega) : t \in [0, r]\}$. The assertion now follows from Theorem 6.7 by noting that the pathwise integral and the Itô integral as defined in Section 5 coincide almost surely since both are limits of Riemann-Itô sums w.r.t. uniform convergence for t in a finite interval, almost surely along a common (sub)sequence of partitions. \square

Consequences

- (1). *Doob decomposition in continuous time:* The Itô integral $M_t^F = \int_0^t F'(B_s) dB_s$ is a local martingale up to T , and M_t^F is a square integrable martingale if $I = \mathbb{R}$ and F' is bounded. Therefore, (6.3.3) can be interpreted as a *continuous time Doob decomposition* of the process $(F(B_t))$ into the (local) martingale part M^F and an adapted process of finite variation. This process takes over the role of the predictable part in discrete time.

In particular, we obtain:

Corollary 6.8 (Martingale problem for Brownian motion). *Brownian motion is a solution of the martingale problem for the operator $\mathcal{L} = \frac{1}{2} \frac{d^2}{dx^2}$ with domain $\text{Dom}(\mathcal{L}) = \{F \in C^2(\mathbb{R}) : \frac{dF}{dx} \text{ is bounded}\}$, i.e., the process*

$$M_t^F = F(B_t) - F(B_0) - \int_0^t (\mathcal{L}f)(B_s) ds$$

is a martingale for any $F \in \text{Dom}(\mathcal{L})$.

The corollary demonstrates how Itô's formula can be applied to obtain solutions of martingale problems, cf./ below for generalizations.

- (2). *Kolmogorov's forward equation:* Taking expectation values in (6.3.3), we recover Kolmogorov's equation

$$E[F(B_t)] = E[F(B_0)] + \int_0^t E[(\mathcal{L}F)(B_s)] ds \quad \forall t \geq 0$$

for any $F \in C_b^2(\mathbb{R})$. In differential form,

$$\frac{d}{dt} E[F(B_t)] = \frac{1}{2} E[(\mathcal{F}'')(B_t)].$$

- (3). *Computation of expected values:* The Itô formula can be applied in many ways to compute expectation values:

Example. (a) For any $n \in \mathbb{N}$, the process

$$B_t^n - \frac{n(n-1)}{2} \int_0^t B_s^{n-2} ds = n \cdot \int_0^t B_s^{n-1} dB_s$$

is a martingale. By taking expectation values for $t = 1$ we obtain the recursion

$$\begin{aligned} E[B_1^n] &= \frac{n(n-1)}{2} \int_0^1 E[B_s^{n-2}] ds = \frac{n(n-1)}{2} \int_0^1 s^{n-2/2} ds \cdot E[B_1^{n-2}] \\ &= (n-1) \cdot E[B_1^{n-2}] \end{aligned}$$

for the moments of the standard normally distributed random variable B_1 . Of course this identity can be obtained directly by integration by parts in the Gaussian integral $\int x^n \cdot e^{-x^2/2} dx$.

(b) For $\alpha \in \mathbb{R}$, the process

$$\exp(\alpha B_t) - \frac{\alpha^2}{2} \int_0^t \exp(\alpha B_s) ds = \alpha \int_0^t \exp(\alpha B_s) dB_s$$

is a martingale because $E[\int_0^t \exp(2\alpha B_s) ds] < \infty$. Denoting by $T_b = \min\{t \geq 0 : B_t = b\}$ the first passage time to a level $b > 0$, we obtain the identity

$$E \left[\int_0^{T_b} \exp(\alpha B_s) ds \right] = \frac{2}{\alpha^2} (e^{\alpha b} - 1) \quad \text{for any } \alpha > 0$$

by optional stopping and dominated convergence.


Itô's formula is also the key tool to derive or solve stochastic differential equations for various stochastic processes of interest:

Example (Brownian motion on S^1). Brownian motion on the unit circle $S^1 = \{z \in \mathbb{C} : |z| = 1\}$ is the process given by

$$Z_t = \exp(iB_t) = \cos B_t + i \cdot \sin B_t$$

where (B_t) is a standard Brownian motion on \mathbb{R}^1 . Itô's formula yields the stochastic differential equation

$$dZ_t = \mathfrak{t}(Z_t) dB_t - \frac{1}{2} \mathfrak{n}(Z_t) dt, \tag{6.3.4}$$



where $\mathbf{t}(z) = iz$ is the unit tangent vector to S^1 at the point z , and $\mathbf{n}(z) = z$ is the outer normal vector. If we would omit the correction term $-\frac{1}{2}\mathbf{n}(Z_t) dt$ in (6.3.4), the solution to the s.d.e. would not stay on the circle. This is contrary to classical o.d.e. where the correction term is not required. For Stratonovich integrals, we obtain the modified equation

$$\circ dZ_t = \mathbf{t}(Z_t) \circ dB_t,$$

which does not involve a correction term!

Quadratic variation of continuous martingales

Next, we will show that the sample paths of continuous local martingales almost surely have finite quadratic variation. Let (M_t) be a continuous local martingale, and fix a sequence (π_n) of partitions of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$. Let

$$V_t^n = \sum_{s \in \pi_n} (M_{s' \wedge t} - M_{s \wedge t})^2$$

denote the quadratic variation of M along π_n . Recall the crucial identity

$$M_t^2 - M_0^2 = \sum_{s \in \pi_n} (M_{s' \wedge t}^2 - M_{s \wedge t}^2) = V_t^n + 2I_t^n \quad (6.3.5)$$

where $I_t^n = \sum_{s \in \pi_n} M_s (M_{s' \wedge t} - M_{s \wedge t})$ are the Riemann sum approximations to the Itô integral $\int_0^t M dM$. The identity shows that V_t^n converges (uniformly) as $n \rightarrow \infty$ if and only if the same holds for I_t^n . Moreover, in this case, we obtain the limit equation

$$M_t^2 - M_0^2 = [M]_t + 2 \int_0^t M_s dM_s \quad (6.3.6)$$

which is exactly Itô's equation for $F(x) = x^2$.

Theorem 6.9 (Existence of quadratic variation). *Suppose that (M_t) is a continuous local martingale on (Ω, \mathcal{A}, P) . Then there exist a continuous non-decreasing process $t \mapsto [M]_t$ and a continuous local martingale $t \mapsto \int_0^t M dM$ such that as $n \rightarrow \infty$,*

$$\sup_{s \in [0, t]} |V_s^n - [M]_s| \rightarrow 0 \quad \text{and} \quad \sup_{s \in [0, t]} \left| I_s^n - \int_0^s M dM \right| \rightarrow 0$$

in probability for any $t \geq 0$, and in $L^2(P)$ respectively if M is bounded. Moreover, the identity (6.3.6) holds.

Notice that in the theorem, we do not assume the existence of an angle bracket process $\langle M \rangle$. Indeed, the theorem proves that for continuous local martingales, the angle bracket process always exists and it coincides almost surely with the quadratic variation process $[M]$! We point out that for discontinuous martingales, $\langle M \rangle$ and $[M]$ do not coincide.

Proof. We first assume that M is a bounded martingale: $|M_t| \leq C$ for some finite constant C . We then show that (I_n) is a Cauchy sequence in the Hilbert space $M_c^2([0, t])$ for any given $t \in \mathbb{R}_+$. To this end let $n, m \in \mathbb{N}$. We assume without loss of generality that $\pi_m \subseteq \pi_n$, otherwise we compare to a common refinement of both partitions. For $s \in \pi_n$, we denote the next partition point in π_n by s' , and the previous partition point in π_m by $[s]_m$. Fix $t \geq 0$. Then

$$\begin{aligned} I_t^n - I_t^m &= \sum_{\substack{s \in \pi_n \\ s < t}} (M_s - M_{[s]_m}) (M_{s' \wedge t} - M_s), & \text{and hence} \\ \|I^n - I^m\|_{M^2([0, t])}^2 &= E [(I_t^n - I_t^m)^2] \\ &= \sum_{\substack{s \in \pi_n \\ s < t}} E [(M_s - M_{[s]_m})^2 (M_{s' \wedge t} - M_s)^2] \\ &\leq E [\delta_m^2]^{1/2} E \left[\left(\sum (\delta M_s)^2 \right)^2 \right]^{1/2}, \end{aligned} \tag{6.3.7}$$

where $\delta_m := \sup\{|M_s - M_r|^2 : |s - r| \leq \text{mesh}(\pi_m)\}$. Here we have used that the non-diagonal summands cancel because M is a martingale.

Since M is bounded and continuous, dominated convergence shows that $E[\delta_m^2] \rightarrow 0$ as $m \rightarrow \infty$. Furthermore,

$$\begin{aligned} E \left[\left(\sum_s (\delta M_s)^2 \right)^2 \right] &= E \left[\sum_s (\delta M_s)^4 \right] + E \left[\sum_{r, s: r < s} (\delta M_r)^2 (\delta M_s)^2 \right] \\ &\leq 4C^2 E \left[\sum_s (\delta M_s)^2 \right] + 2 E \left[\sum_r (\delta M_r)^2 E \left[\sum_{s > r} (\delta M_s)^2 | \mathcal{F}_r \right] \right] \\ &\leq 6C^2 E[M_t^2 - M_0^2] \leq 6C^4 < \infty. \end{aligned}$$

Here we have used that by the martingale property,

$$\begin{aligned} E \left[\sum_s (\delta M_s)^2 \right] &= E[M_t^2 - M_0^2] \leq C^2, \quad \text{and} \\ E \left[\sum_{s>r} (\delta M_s)^2 | \mathcal{F}_r \right] &= E[M_t^2 - M_r^2 | \mathcal{F}_r] \leq C^2. \end{aligned}$$

By (6.3.7), $\|I^n - I^m\|_{M^2([0,t])}^2 \rightarrow 0$ as $n, m \rightarrow \infty$. Hence $(I_s^n)_{s \in [0,t]}$ converges uniformly as $n \rightarrow \infty$ in the $L^2(P)$ sense. By (6.3.5), $(V_s^n)_{s \in [0,t]}$ converges uniformly as $n \rightarrow \infty$ in the $L^2(P)$ sense as well. Hence the limits $\int_0^\bullet M dM$ and $[M]$ exist, the stochastic integral is in $M_c^2([0,t])$, and the identity (6.3.6) holds.

It remains to extend the result from bounded martingales to local martingales. If M is a continuous local martingale then there exists a sequence of stopping times $T_k \uparrow \infty$ such that the stopped processes $(M_{T_k \wedge t})_{t \geq 0}$ are continuous bounded martingales. Hence the corresponding quadratic variations $[M_{T_k \wedge \bullet}]$ converge uniformly in the $L^2(P)$ sense for any finite t and k . Therefore, the approximations V_t^n for the quadratic variation of M converge uniformly in the $L^2(P)$ sense on each of the random intervals $[0, T_k \wedge t]$, and thus for any $\varepsilon, \delta > 0$,

$$P \left[\sup_{s \leq t} |V_s^n - [M]_s| > \varepsilon \right] \leq P[t > T_k] + P \left[\sup_{s \leq T_k} |V_s^n - [M]_s| > \varepsilon \right] \leq \delta$$

for k, n sufficiently large. □

Having shown the existence of the quadratic variation $[M]$ for continuous local martingales, we observe next that $[M]$ is always non-trivial if M is not constant:

Theorem 6.10 (Non-constant continuous martingales have non-trivial quadratic variation). *Suppose that (M_t) is a continuous local martingale. If $[M]_t = 0$ almost surely for some $t \geq 0$, then M is almost surely constant on the interval $[0, t]$.*

Proof. Again, we assume at first that M is a bounded martingale. Then the Itô integral $\int_0^\bullet M dM$ is a martingale as well. Therefore, by (6.3.6),

$$\|M - M_0\|_{M^2([0,t])}^2 = E[(M_t - M_0)^2] = E[M_t^2 - M_0^2] = E[[M]_t] = 0,$$

i.e., $M_s = M_0$ for any $s \in [0, t]$. In the general case, the assertion follows once more by localization. \square

The theorem shows in particular that every local martingale with continuous finite variation paths is almost surely constant, i.e., *the Doob type decomposition of a continuous stochastic process into a local martingale and a continuous finite variation process starting at 0 is unique up to equivalence*. As a consequence we observe that the quadratic variation is the *unique* angle bracket process of M . In particular, up to modification on measure zero sets, $[M]$ does not depend on the chosen partition sequence (π_n) :

Corollary 6.11 (Quadratic variation as unique angle bracket process). *Suppose that (M_t) is a continuous local martingale. Then $[M]$ is the up to equivalence unique continuous process of finite variation such that $[M]_0 = 0$ and $M_t^2 - [M]_t$ is a local martingale.*

Proof. By (6.3.6), $M_t^2 - [M]_t$ is a continuous local martingale. To prove uniqueness, suppose that (A_t) and (\tilde{A}_t) are continuous finite variation processes with $A_0 = \tilde{A}_0 = 0$ such that both $M_t^2 - A_t$ and $M_t^2 - \tilde{A}_t$ are local martingales. Then $A_t - \tilde{A}_t$ is a continuous local martingale as well. Since the paths have finite variation, the quadratic variation of $A - \tilde{A}$ vanishes. Hence almost surely, $A_t - \tilde{A}_t = A_0 - \tilde{A}_0 = 0$ for all t . \square

From continuous martingales to Brownian motion

A remarkable consequence of Itô's formula for martingales is that any continuous local martingale (M_t) (up to $T = \infty$) with quadratic variation given by $[M]_t = t$ for any $t \geq 0$ is a Brownian motion ! In fact, for $0 \leq s \leq t$ and $p \in \mathbb{R}$, Itô's formula yields

$$e^{ipM_t} - e^{ipM_s} = ip \int_s^t e^{ipM_r} dM_r - \frac{p^2}{2} \int_s^t e^{ipM_r} dr$$

where the stochastic integral can be identified as a local martingale. From this identity it is not difficult to conclude that the increment $M_t - M_s$ is conditionally independent of \mathcal{F}_s^M with characteristic function

$$E[e^{ip(M_t - M_s)}] = e^{-p^2(t-s)/2} \quad \text{for any } p \in \mathbb{R},$$

i.e., (M_t) has independent increments with distribution $M_t - M_s \sim N(0, t - s)$.

Theorem 6.12 (P. Lévy 1948). *A continuous local martingale $(M_t)_{t \in [0, \infty)}$ is a Brownian motion if and only if almost surely,*

$$[M]_t = t \quad \text{for any } t \geq 0.$$

Exercise (Lévy's characterization of Brownian motion). Extend the sketch above to a proof of Theorem 6.12.

Lévy's Theorem is the basis for many important developments in stochastic analysis including transformations and weak solutions for stochastic differential equations. An extension to the multi-dimensional case with a detailed proof, as well as several applications, are contained in Section 11.1 below.

One remarkable consequence of Lévy's characterization of Brownian motion is that every continuous local martingale can be represented as a time-changed Brownian motion (in general possibly on an extended probability space):

Exercise (Continuous local martingales as time-changed Brownian motions). Let $(M_t)_{t \in [0, \infty)}$ be a continuous local martingale, and assume for simplicity that $t \mapsto [M]_t$ is almost surely strictly increasing with $\lim_{t \rightarrow \infty} [M]_t = \infty$. Prove that there exists a Brownian motion $(B_t)_{t \in [0, \infty)}$ such that

$$M_t = B_{[M]_t} \quad \text{for } t \in [0, \infty). \quad (6.3.8)$$

Hint: Set $B_a = M_{T_a}$ where $T_a = [M]^{-1}(a) = \inf\{t \geq 0 : [M]_t = a\}$, and verify by Lévy's characterization that B is a Brownian motion.

In a more general form, the representation of continuous local martingales as time-changed Brownian motions is due to Dambis and Dubins-Schwarz (1965), cf. [37] or Section 11.2 below for details. Remarkably, even before Itô, Wolfgang Doeblin, the son of Alfred Doeblin, had developed an alternative approach to stochastic calculus where stochastic integrals are defined as time changes of Brownian motion. Doeblin died when fighting as a French soldier at the German front in World War II, and his results that were hidden in a closed envelope at the Académie de Sciences have become known and been published only recently, more than fifty years after their discovery, cf. [Doeblin, Sur l'équation de Kolmogoroff, 1940/2000], [Yor: Présentation du pli cacheté, C.R.Acad.Sci. Paris 2000].

6.4 Multivariate and time-dependent Itô formula

We now extend Itô's formula to \mathbb{R}^d -valued functions and stochastic processes. Let $u \in (0, \infty]$ and suppose that $X : [0, u) \rightarrow D, X_t = (X_t^{(1)}, \dots, X_t^{(d)})$, is a continuous function taking values in an open set $D \subseteq \mathbb{R}^d$. As before, we fix a sequence (π_n) of partitions of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$. For a function $F \in C^2(D)$, we have similarly as in the one-dimensional case:

$$F(X_{s' \wedge t}) - F(X_s) = \nabla F(X_s) \cdot (X_{s' \wedge t} - X_s) + \frac{1}{2} \sum_{i,j=1}^d \frac{\partial^2 F}{\partial x_i \partial x_j}(X_s) (X_{s' \wedge t}^{(i)} - X_s^{(i)}) (X_{s' \wedge t}^{(j)} - X_s^{(j)}) + R_s^{(n)} \quad (6.4.1)$$

for any $s \in \pi_n$ with $s < t$ where the dot denotes the Euclidean inner product $R_s^{(n)}$ is the remainder term in Taylor's formula. We would like to obtain a multivariate Itô formula by summing over $s \in \pi_n$ with $s < t$ and taking the limit as $n \rightarrow \infty$. A first problem that arises in this context is the identification of the limit of the sums

$$\sum_{\substack{s \in \pi_n \\ s < t}} g(X_s) \delta X_s^{(i)} \delta X_s^{(j)}$$

for a continuous function $g : D \rightarrow \mathbb{R}$ as $n \rightarrow \infty$.

Covariation

Suppose that $X, Y : [0, u) \rightarrow \mathbb{R}$ are continuous functions with continuous quadratic variations $[X]_t$ and $[Y]_t$ w.r.t. (π_n) .

Definition. *The function*

$$[X, Y]_t = \lim_{n \rightarrow \infty} \sum_{s \in \pi_n} (X_{s' \wedge t} - X_{s \wedge t})(Y_{s' \wedge t} - Y_{s \wedge t}), \quad t \in [0, u),$$

*is called the **covariation of X and Y w.r.t. (π_n)** if the limit exists.*

The covariation $[X, Y]_t$ is the bilinear form corresponding to the quadratic form $[X]_t$. In particular, $[X, X] = [X]$. Furthermore:

Lemma 6.13 (Polarization identity). *The covariation $[X, Y]_t$ exists and is a continuous function in t if and only if the quadratic variation $[X + Y]_t$ exists and is continuous respectively. In this case,*

$$[X, Y]_t = \frac{1}{2}([X + Y]_t - [X]_t - [Y]_t).$$

Proof. For $n \in \mathbb{N}$ we have

$$2 \sum_{s \in \pi_n} \delta X_s \delta Y_s = \sum_{s \in \pi_n} (\delta X_s + \delta Y_s)^2 - \sum_{s \in \pi_n} (\delta X_s)^2 - \sum_{s \in \pi_n} (\delta Y_s)^2.$$

The assertion follows as $n \rightarrow \infty$ because the limits $[X]_t$ and $[Y]_t$ of the last two terms are continuous functions by assumption. \square

Remark. Note that by the polarization identity, the covariation $[X, Y]_t$ is the difference of two increasing functions, i.e., $t \mapsto [X, Y]_t$ has finite variation.

Example. (1). *Functions and processes of finite variation:* If Y has finite variation then $[X, Y]_t = 0$ for any $t \geq 0$. Indeed,

$$\left| \sum_{s \in \pi_n} \delta X_s \delta Y_s \right| \leq \sup_{s \in \pi_n} |\delta X_s| \cdot \sum_{s \in \pi_n} |\delta Y_s|$$

and the right hand side converges to 0 by uniform continuity of X on $[0, t]$. In particular, we obtain again

$$[X + Y] = [X] + [Y] + 2[X, Y] = [X].$$

(2). *Independent Brownian motions:* If (B_t) and (\tilde{B}_t) are independent Brownian motions on a probability space (Ω, \mathcal{A}, P) then for any given sequence (π_n) ,

$$[B, \tilde{B}]_t = \lim_{n \rightarrow \infty} \sum_{s \in \pi_n} \delta B_s \delta \tilde{B}_s = 0 \quad \text{for any } t \geq 0$$

P -almost surely. For the proof note that $(B_t + \tilde{B}_t)/\sqrt{2}$ is again a Brownian motion, whence

$$[B, \tilde{B}]_t = [(B + \tilde{B})/\sqrt{2}]_t - \frac{1}{2}[B]_t - \frac{1}{2}[\tilde{B}]_t = t - \frac{t}{2} - \frac{t}{2} = 0 \quad \text{almost surely.}$$

(3). *Itô processes:* If $I_t = \int_0^t G_s dB_s$ and $J_t = \int_0^t H_s d\tilde{B}_s$ with continuous adapted processes (G_t) and (H_t) and Brownian motions (B_t) and (\tilde{B}_t) then

$$[I, J]_t = 0 \quad \text{if } B \text{ and } \tilde{B} \text{ are independent, and} \quad (6.4.2)$$

$$[I, J]_t = \int_0^t G_s H_s ds \quad \text{if } B = \tilde{B}, \quad (6.4.3)$$

cf. Theorem ?? below.

More generally, under appropriate assumptions on G, H, X and Y , the identity

$$[I, J]_t = \int_0^t G_s H_s d[X, Y]_s$$

holds for Itô integrals $I_t = \int_0^t G_s dX_s$ and $J_t = \int_0^t H_s dY_s$, cf. e.g. Corollary ??.

Itô to Stratonovich conversion

The covariation also occurs as the correction term in Itô compared to Stratonovich integrals:

Theorem 6.14. *If the Itô integral*

$$\int_0^t X_s Y_s = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} X_s \delta Y_s$$

and the covariation $[X, Y]_t$ exists along a sequence (π_n) of partitions with $\text{mesh}(\pi_n) \rightarrow 0$ then the corresponding backward Itô integral $\int_0^t X_s \hat{d}Y_s$ and the Stratonovich integral $\int_0^t X_s \circ dY_s$ also exist, and

$$\begin{aligned} \int_0^t X_s \hat{d}Y_s &= \int_0^t X_s Y_s + [X, Y]_t, & \text{and} \\ \int_0^t X_s \circ dY_s &= \int_0^t X_s Y_s + \frac{1}{2}[X, Y]_t. \end{aligned}$$

Proof. This follows from the identities

$$\begin{aligned} \sum X_{S' \wedge t} \delta Y_s &= \sum X_s \delta Y_s + \sum \delta X_s \delta Y_s, & \text{and} \\ \sum \frac{1}{2}(X_s + X_{S' \wedge t}) \delta Y_s &= \sum X_s \delta Y_s + \frac{1}{2} \sum \delta X_s \delta Y_s. \end{aligned}$$

□

Itô's formula in \mathbb{R}^d

By the polarization identity, if $[X]_t$, $[Y]_t$ and $[X + Y]_t$ exist and are continuous then $[X, Y]_t$ is a continuous function of finite variation.

Lemma 6.15. *Suppose that X, Y and $X + Y$ are continuous function on $[0, u)$ with continuous quadratic variations w.r.t. (π_n) . Then*

$$\sum_{\substack{s \in \pi_n \\ s < t}} H_s (X_{s' \wedge t} - X_s)(Y_{s' \wedge t} - Y_s) \longrightarrow \int_0^t H_s d[X, Y]_s \quad \text{as } n \rightarrow \infty$$

for any continuous function $H : [0, u) \rightarrow \mathbb{R}$ and any $t \geq 0$.

Proof. The assertion follows from Lemma 6.3 by polarization. \square

By Lemma 6.15, we can take the limit as $\text{mesh}(\pi_n) \rightarrow 0$ in the equation derived by summing (6.4.2) over all $s \in \pi_n$ with $s < t$. In analogy to the one-dimensional case, this yields the following multivariate version of the pathwise Itô formula:

Theorem 6.16 (Multivariate Itô formula without probability). *Suppose that $X : [0, u) \rightarrow D \subseteq \mathbb{R}^d$ is a continuous function with continuous covariations $[X^{(i)}, X^{(j)}]_t, 1 \leq i, j \leq d$, w.r.t. (π_n) . Then for any $F \in C^2(D)$ and $t \in [0, u)$,*

$$F(X_t) = F(X_0) + \int_0^t \nabla F(X_s) \cdot dX_s + \frac{1}{2} \sum_{i,j=1}^d \int_0^t \frac{\partial^2 F}{\partial x_i \partial x_j}(X_s) d[X^{(i)}, X^{(j)}]_s,$$

where the Itô integral is the limit of Riemann sums along (π_n) :

$$\int_0^t \nabla F(X_s) \cdot dX_s = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} \nabla F(X_s) \cdot (X_{s' \wedge t} - X_s). \quad (6.4.4)$$

The details of the proof are similar to the one-dimensional case and left as an exercise to the reader. Note that the theorem shows in particular that the Itô integral in (6.4.4) is independent of the sequence (π_n) if the same holds for the covariations $[X^{(i)}, X^{(j)}]$.

Remark (Existence of pathwise Itô integrals). The theorem implies the existence of the Itô integral $\int_0^t b(X_s) \cdot dX_s$ if $b = \nabla F$ is the gradient of a C^2 function $F : D \subseteq \mathbb{R}^d \rightarrow \mathbb{R}$. In contrast to the one-dimensional case, not every C^1 vector field $b : D \rightarrow \mathbb{R}^d$ is a gradient. Therefore, for $d \geq 2$ we do **not** obtain existence of $\int_0^t b(X_s) \cdot dX_s$ for any $b \in C^1(D, \mathbb{R}^d)$ from Itô's formula. In particular, we do not know in general if the integrals $\int_0^t \frac{\partial F}{\partial x_i}(X_s) dX_s^{(i)}, 1 \leq i \leq d$, exists and if

$$\int_0^t \nabla F(X_s) \cdot dX_s = \sum_{i=1}^d \int_0^t \frac{\partial F}{\partial x_i}(X_s) dX_s^{(i)}.$$

If (X_t) is a Brownian motion this is almost surely the case by the existence proof for Itô integrals w.r.t. Brownian motion from Section 5.

Example (Itô's formula for Brownian motion in \mathbb{R}^d). Suppose that $B_t = (B_t^{(1)}, \dots, B_t^{(d)})$ is a d -dimensional Brownian motion defined on a probability space (Ω, \mathcal{A}, P) . Then the component processes $B_t^{(1)}, \dots, B_t^{(d)}$ are independent one-dimensional Brownian motions. Hence for a given sequence of partitions (π_n) with $\text{mesh}(\pi_n) \rightarrow 0$, the covariations $[B^{(i)}, B^{(j)}], 1 \leq i, j \leq d$, exists almost surely by Theorem 6.6 and the example above, and

$$[B^{(i)}, B^{(j)}] = t \cdot \delta_{ij} \quad \forall t \geq 0$$

P -almost surely. Therefore, we can apply Itô's formula to almost every trajectory. For an open subset $D \subseteq \mathbb{R}^d$ and a function $F \in C^2(D)$ we obtain:

$$F(B_t) = F(B_0) + \int_0^t \nabla F(B_s) \cdot dB_s + \frac{1}{2} \int_0^t \Delta F(B_s) ds \quad \forall t < T_{DC} \quad P\text{-a.s.} \quad (6.4.5)$$

where $T_{DC} := \inf\{t \geq 0 : B_t \notin D\}$ denotes the first exit time from D . As in the one-dimensional case, (6.4.5) yields a decomposition of the process $F(B_t)$ into a continuous local martingale and a continuous process of finite variation, cf. Section ?? for applications.

Product rule, integration by parts

As a special case of the multivariate Itô formula, we obtain the following integration by parts identity for Itô integrals:

Corollary 6.17. *Suppose that $X, Y : [0, u) \rightarrow \mathbb{R}$ are continuous functions with continuous quadratic variations $[X]$ and $[Y]$, and continuous covariation $[X, Y]$. Then*

$$X_t Y_t - X_0 Y_0 = \int_0^t \begin{pmatrix} Y_s \\ X_s \end{pmatrix} \cdot d \begin{pmatrix} X_s \\ Y_s \end{pmatrix} + [X, Y]_t \quad \text{for any } t \in [0, u). \quad (6.4.6)$$

If one, or, equivalently, both of the Itô integrals $\int_0^t Y_s dX_s$ and $\int_0^t X_s dY_s$ exist then (6.4.6) yields

$$X_t Y_t - X_0 Y_0 = \int_0^t Y_s dX_s + \int_0^t X_s dY_s + [X, Y]_t. \quad (6.4.7)$$

Proof. The identity (6.4.6) follows by applying Itô's formula in \mathbb{R}^2 to the process (X_t, Y_t) and the function $F(x, y) = xy$. If one of the integrals $\int_0^t Y dX$ or $\int_0^t X dY$ exists, then the other exists as well, and

$$\int_0^t \begin{pmatrix} Y_s \\ X_s \end{pmatrix} \cdot d \begin{pmatrix} X_s \\ Y_s \end{pmatrix} = \int_0^t Y_s dX_s + \int_0^t X_s dY_s.$$

□

As it stands, (6.4.7) is an integration by parts formula for Itô integrals which involves the correction term $[X, Y]_t$. In differential notation, it is a product rule for Itô differentials:

$$d(XY) = X dY + Y dX + [X, Y].$$

Again, in Stratonovich calculus a corresponding product rule holds without the correction term $[X, Y]$:

$$\circ d(XY) = X \circ dY + Y \circ dX.$$

Remark / Warning (Existence of $\int X dY$, Lévy area). Under the conditions of the theorem, the Itô integrals $\int_0^t X dY$ and $\int_0^t Y dX$ do not necessarily exist! The following statements are equivalent:

- (1). The Itô integral $\int_0^t Y_s dX_s$ exists (along (π_n)).
- (2). The Itô integral $\int_0^t X_s dY_s$ exists.

(3). The **Lévy area** $A_t(X, Y)$ defined by

$$A_t(X, Y) = \int_0^t (Y dX - X dY) = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} (Y_s \Delta X_s - X_s \Delta Y_s)$$

exists.

Hence, if the Lévy area $A_t(X, Y)$ is given, the stochastic integrals $\int X dY$ and $\int Y dX$ can be constructed pathwise. Pushing these ideas further leads to the rough paths theory developed by T. Lyons and others, cf. [Lyons, St. Flour], [Friz: Rough paths theory].

Example (Integrating finite variation processes w.r.t. Brownian motion). If (H_t) is an adapted process with continuous sample paths of finite variation and (B_t) is a one-dimensional Brownian motion then $[H, B] = 0$, and hence

$$H_t B_t - H_0 B_0 = \int_0^t H_s dB_s + \int_0^t B_s dH_s.$$

This integration by parts identity can be used as an alternative definition of the stochastic integral $\int_0^t H dB$ for integrands of finite variation, which can then again be extended to general integrands in $\mathcal{L}_a^2(0, t)$ by the Itô isometry.

Time-dependent Itô formula

The multi-dimensional Itô formula can be applied to functions that depend explicitly on the time variable t or on the quadratic variation $[X]_t$. For this purpose we simply add t or $[X]_t$ respectively as an additional component to the function, i.e., we apply the multi-dimensional Itô formula to $Y_t = (t, X_t)$ or $Y_t = (t, [X]_t)$ respectively.

Theorem 6.18. *Suppose that $X : [0, u) \rightarrow \mathbb{R}^d$ is a continuous function with continuous covariations $[X^{(i)}, X^{(j)}]_t$, along (π_n) , and let $F \in C^2(A([0, u)) \times \mathbb{R}^d)$. If $A : [0, u) \rightarrow \mathbb{R}$ is a continuous function of finite variation then the integral*

$$\int_0^t \nabla_x F(A_s, X_s) \cdot dX_s = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} \nabla_x F(A_s, X_s) \cdot (X_{s' \wedge t} - X_s)$$

exists, and the Itô formula

$$F(A_t, X_t) = F(0, X_0) + \int_0^t \nabla_x F(A_s, X_s) \cdot dX_s + \int_0^t \frac{\partial F}{\partial a}(A_s, X_s) dA_s \quad (6.4.8)$$

$$\frac{1}{2} \sum_{i,j=1}^d \int_0^t \frac{\partial^2 F}{\partial x_i \partial x_j}(A_s, X_s) d[X^{(i)}, X^{(j)}]_s \quad (6.4.9)$$

holds for any $t \geq 0$. Here $\partial F/\partial a$ denotes the derivative of $F(a, x)$ w.r.t. the first component, and $\nabla_x F$ and $\partial^2 F/\partial x_i \partial x_j$ are the gradient and the second partial derivatives w.r.t. the other components. The most important application of the theorem is for $A_t = t$. Here we obtain the time-dependent Itô formula

$$dF(t, X_t) = \nabla_x F(t, X_t) \cdot dX_t + \frac{\partial F}{\partial t}(t, X_t) dt + \frac{1}{2} \sum_{i,j=1}^d \frac{\partial^2 F}{\partial x_i \partial x_j}(t, X_t) d[X^{(i)}, X^{(j)}]_t. \quad (6.4.10)$$

Similarly, if $d = 1$ and $A_t = [X]_t$ then we obtain

$$dF([X]_t, X_t) = \frac{\partial F}{\partial t}([X]_t, X_t) dt + \left(\frac{\partial F}{\partial a} + \frac{1}{2} \frac{\partial^2 F}{\partial x^2} \right) ([X]_t, X_t) d[X]_t. \quad (6.4.11)$$

If $(X)_t$ is a Brownian motion and $d = 1$ then both formulas coincide.

Proof. Let $Y_t = (Y_t^{(0)}, Y_t^{(1)}, \dots, Y_t^{(d)}) := (A_t, X_t)$. Then $[Y^{(0)}, Y^{(i)}]_t = 0$ for any $t \geq 0$ and $0 \leq i \leq d$ because $Y_t^{(0)} = A_t$ has finite variation. Therefore, by Itô's formula in \mathbb{R}^{d+1} ,

$$F(A_t, X_t) = F(A_0, X_0) + I_t + \frac{1}{2} \sum_{i,j=1}^d \frac{\partial^2 F}{\partial x_i \partial x_j}(A_s, X_s) d[X^{(i)}, X^{(j)}]_s$$

where

$$\begin{aligned} I_t &= \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} \nabla^{\mathbb{R}^{d+1}} F(A_s, X_s) \cdot \begin{pmatrix} A_{s' \wedge t} - A_s \\ X_{s' \wedge t} - X_s \end{pmatrix} \\ &= \lim_{n \rightarrow \infty} \left(\sum \frac{\partial F}{\partial a}(A_s, X_s) (A_{s' \wedge t} - A_s) + \sum \nabla_x F(A_s, X_s) \cdot (X_{s' \wedge t} - X_s) \right). \end{aligned}$$

The first sum on the right hand side converges to the Stieltjes integral $\int_0^t \frac{\partial F}{\partial a}(A_s, X_s) dA_s$ as $n \rightarrow \infty$. Hence, the second sum also converges, and we obtain (6.4.7) in the limit as $n \rightarrow \infty$. \square

Note that if $h(t, x)$ is a solution of the dual heat equation

$$\frac{\partial h}{\partial t} + \frac{1}{2} \frac{\partial^2 h}{\partial x^2} = 0 \quad \text{for } t \geq 0, x \in \mathbb{R}, \quad (6.4.12)$$

then by (6.4.11),

$$h([X]_t, X_t) = h(0, X_0) + \int_0^t \frac{\partial h}{\partial x}([X]_s, X_s) dX_s.$$

In particular, if (X_t) is a Brownian motion, or more generally a local martingale, then $h([X]_t, X_t)$ is also a local martingale. The next example considers two situations where this is particularly interesting:

Example. (1). *Itô exponentials:* For any $\alpha \in \mathbb{R}$, the function

$$h(t, x) = \exp(\alpha x - \alpha^2 t/2)$$

satisfies (6.4.12) and $\partial h / \partial x = \alpha h$. Hence the function

$$Z_t^{(\alpha)} := \exp\left(\alpha X_t - \frac{1}{2} \alpha^2 [X]_t\right)$$

is a solution of the Itô differential equation

$$dZ_t^{(\alpha)} = \alpha Z_t^{(\alpha)} dX_t$$

with initial condition $Z_0^{(\alpha)} = 1$. This shows that in Itô calculus, the functions $Z_t^{(\alpha)}$ are the correct replacements for the exponential functions. The additional factor $\exp(-\alpha^2 [X]_t/2)$ should be thought of as an appropriate renormalization in the continuous time limit.

For a Brownian motion (X_t) , we obtain the exponential martingales as generalized exponentials.

(2). *Hermite polynomials*: For $n = 0, 1, 2, \dots$, the Hermite polynomials

$$h_n(t, x) = \frac{\partial^n}{\partial \alpha^n} \exp(\alpha x - \frac{1}{2} \alpha^2 t) \Big|_{\alpha=0}$$

also satisfy (6.4.12). The first Hermite polynomials are $1, x, x^2 - t, x^3 - 3tx, \dots$

Note also that

$$\exp(\alpha x - \alpha^2 t/2) = \sum_{n=0}^{\infty} \frac{\alpha^n}{n!} h_n(t, x)$$

by Taylor's theorem. Moreover, the following properties can be easily verified:

$$h_n(1, x) = e^{x^2/2} (-1)^n \frac{d^n}{dx^n} e^{-x^2/2} \quad \text{for any } x \in \mathbb{R}, \quad (6.4.13)$$

$$h_n(t, x) = t^{n/2} h_n(1, x/\sqrt{t}) \quad \text{for any } t \geq 0, x \in \mathbb{R}, \quad (6.4.14)$$

$$\frac{\partial h_n}{\partial x} = n h_{n-1}, \quad \frac{\partial h_n}{\partial t} + \frac{1}{2} \frac{\partial^2 h_n}{\partial x^2} = 0. \quad (6.4.15)$$

For example, (6.4.13) holds since

$$\exp(\alpha x - \alpha^2/2) = \exp(-(x - \alpha)^2/2) \exp(x^2/2)$$

yields

$$h_N(1, x) = \exp(x^2/2) (-1)^n \frac{d^n}{d\beta^n} \exp(-\beta^2/2) \Big|_{\beta=x},$$

and (6.4.14) follows from

$$\begin{aligned} \exp(\alpha x - \alpha^2 t/2) &= \exp(\alpha \sqrt{t} \cdot (x/\sqrt{t}) - (\alpha \sqrt{t})^2/2) \\ &= \sum_{n=0}^{\infty} \frac{\alpha^n}{n!} t^{n/2} h_n(1, x/\sqrt{t}). \end{aligned}$$

By (6.4.13) and (6.4.14), h_n is a polynomial of degree n . For any $n \geq 0$, the function

$$H_t^{(n)} := h_n([X]_t, X_t)$$

is a solution of the Itô equation

$$dH_t^{(n)} = n H_t^{(n-1)} dX_t. \quad (6.4.16)$$

Therefore, the Hermite polynomials are appropriate replacements for the ordinary monomials x^n in Itô calculus. If $X_0 = 0$ then $H_0^{(n)} = 0$ for $n \geq 1$, and we obtain inductively

$$H_t^{(0)} = 1, \quad H_t^{(1)} = \int_0^t dX_s, \quad H_t^{(2)} = \int H_s^{(1)} dX_s = \int_0^t \int_0^s dX_r dX_s,$$

and so on.

Corollary 6.19 (Itô 1951). *If $X : [0, u) \rightarrow \mathbb{R}$ is continuous with $X_0 = 0$ and continuous quadratic variation then for $t \in [0, u)$,*

$$\int_0^t \int_0^{s_n} \cdots \int_0^{s_2} dX_{s_1} \cdots dX_{s_{n-1}} dX_{s_n} = \frac{1}{n!} h_n([X]_t, X_t).$$

Proof. The equation follows from (6.4.16) by induction on n . □

Iterated Itô integrals occur naturally in Taylor expansions of Itô calculus. Therefore, the explicit expression from the corollary is valuable for numerical methods for stochastic differential equations, cf. Section ?? below.

Chapter 7

Brownian Motion and Partial Differential Equations

The stationary and time-dependent Itô formula enable us to work out the connection of Brownian motion to several partial differential equations involving the Laplace operator in detail. One of the many consequences is the evaluation of probabilities and expectation values for Brownian motion by p.d.e. methods. More generally, Itô's formula establishes a link between stochastic processes and analysis that is extremely fruitful in both directions.

Suppose that (B_t) is a d -dimensional Brownian motion defined on a probability space (Ω, \mathcal{A}, P) such that every sample path $t \mapsto B_t(\omega)$ is continuous. We first note that Itô's formula shows that Brownian motion solves the martingale problem for the operator $\mathcal{L} = \frac{1}{2}\Delta$ in the following sense:

Corollary 7.1 (Time-dependent martingale problem). *The process*

$$M_t^F = F(t, B_t) - F(0, B_0) - \int_0^t \left(\frac{\partial F}{\partial s} + \frac{1}{2} \Delta F \right) (s, B_s) ds$$

is a continuous (\mathcal{F}_t^B) martingale for any C^2 function $F : [0, \infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ with bounded first derivatives. Moreover, M^F is a continuous local martingale up to $T_{D^c} = \inf\{t \geq 0 : B_t \notin D\}$ for any $F \in C^2([0, \infty) \times D)$, $D \subseteq \mathbb{R}^d$ open.

Proof. By the continuity assumptions one easily verifies that M^F is (\mathcal{F}_t^B) adapted. Moreover, by the time-dependent Itô formula (6.4.10),

$$M_t^F = \int_0^t \nabla_x F(s, B_s) \cdot dB_s \quad \text{for } t < T_{D^c},$$

which implies the claim. \square

Choosing a function F that does not explicitly depend on t , we obtain in particular that

$$M_t^F = F(B_t) - F(B_0) - \int_0^t \frac{1}{2} \Delta F(B_s) ds$$

is a martingale for any $f \in C_b^2(\mathbb{R}^d)$, and a local martingale up to T_{D^c} for any $F \in C^2(D)$.

7.1 Dirichlet problem, recurrence and transience

As a first consequence of Corollary 7.1 we can now complete the proof of the stochastic representation for solutions of the Dirichlet problem that has been already mentioned in Section 3.2 above. By solving the Dirichlet problem for balls explicitly, we will then study recurrence, transience and polar sets for multi-dimensional Brownian motion.

The Dirichlet problem revisited

Suppose that $h \in C^2(D) \cap C(\overline{D})$ is a solution of the Dirichlet problem

$$\Delta h = 0 \quad \text{on } D, \quad h = f \quad \text{on } \partial D, \quad (7.1.1)$$

for a bounded open set $D \subset \mathbb{R}^d$ and a continuous function $f : \partial D \rightarrow \mathbb{R}$. If (B_t) is under P_x a continuous Brownian motion with $B_0 = x$ P_x -almost surely, then by Corollary 7.1, the process $h(B_t)$ is a local (\mathcal{F}_t^B) martingale up to T_{D^c} . By applying the optional

stopping theorem with a localizing sequence of bounded stopping times $S_n \nearrow T_{D^c}$, we obtain

$$h(x) = E_x[h(B_0)] = E_x[h(B_{S_n})] \quad \text{for any } n \in \mathbb{N}.$$

Since $P_x[T_{D^c} < \infty] = 1$ and h is bounded on \overline{D} , dominated convergence then yields the stochastic representation

$$h(x) = E_x[h(B_{T_{D^c}})] = E_x[f(B_{T_{D^c}})] \quad \text{for any } x \in \mathbb{R}^d.$$

We thus have shown:

Theorem 7.2 (Stochastic representation for solutions of the Dirichlet problem).

Suppose that D is a bounded open subset of \mathbb{R}^d , f is a continuous function on the boundary ∂D , and $h \in C^2(D) \cap C(\overline{D})$ is a solution of the Dirichlet problem (7.1.1). Then

$$h(x) = E_x[f(B_T)] \quad \text{for any } x \in \overline{D}.$$

We will generalize this result substantially in Theorem 7.5 below. Before, we apply the Dirichlet problem to study recurrence and transience of Brownian motions:

Recurrence and transience of Brownian motion in \mathbb{R}^d

Let (B_t) be a d -dimensional Brownian motion on (Ω, \mathcal{A}, P) with initial value $B_0 = x_0, x_0 \neq 0$. For $r \geq 0$ let


$$T_r = \inf\{t > 0 : |B_t| = r\}.$$

We now compute the probabilities $P[T_a < T_b]$ for $a < |x_0| < b$. Note that this is a multi-dimensional analogue of the *classical ruin problem*. To compute the probability for given a, b we consider the domain

$$D = \{x \in \mathbb{R}^d : a < |x| < b\}.$$

For $b < \infty$, the first exit time T_{D^c} is almost surely finite,

$$T_{D^c} = \min(T_a, T_b), \quad \text{and} \quad P[T_a < T_b] = P[|B_{T_{D^c}}| = a].$$



Suppose that $h \in C(\bar{U}) \cap C^2(U)$ is a solution of the Dirichlet problem

$$\Delta h(x) = 0 \quad \text{for all } x \in D, \quad h(x) = \begin{cases} 1 & \text{if } |x| = a, \\ 0 & \text{if } |x| = b. \end{cases} \quad (7.1.2)$$

Then $h(B_t)$ is a bounded local martingale up to T_{D^c} and optional stopping yields

$$P[T_a < T_b] = E[h(B_{T_{D^c}})] = h(x_0). \quad (7.1.3)$$

By rotational symmetry, the solution of the Dirichlet problem (7.1.2) can be computed explicitly. The Ansatz $h(x) = f(|x|)$ leads us to the boundary value problem

$$\frac{d^2 f}{dr^2}(|x|) + \frac{d-1}{|x|} \frac{df}{dr}(|x|) = 0, \quad f(a) = 1, f(b) = 0,$$

for a second order ordinary differential equation. Solutions of the o.d.e. are linear combinations of the constant function 1 and the function

$$\phi(s) := \begin{cases} s & \text{for } d = 1, \\ \log s & \text{for } d = 2, \\ s^{2-d} & \text{for } d \geq 3. \end{cases}$$




Figure 7.1: The function $\phi(s)$ for different values of d : red ($d = 1$), blue ($d = 2$) and purple ($d = 3$)

Hence, the unique solution f with boundary conditions $f(a) = 1$ and $f(b) = 0$ is

$$f(r) = \frac{\phi(b) - \phi(r)}{\phi(b) - \phi(a)}.$$

Summarizing, we have shown:

Theorem 7.3 (Ruin problem in \mathbb{R}^d). For $a, b > 0$ with $a < |x_0| < b$,

$$P[T_a < T_b] = \frac{\phi(b) - \phi(|x_0|)}{\phi(b) - \phi(a)}, \quad \text{and}$$

$$P[T_b < \infty] = \begin{cases} 1 & \text{for } d \leq 2 \\ (a/|x_0|)^{d-2} & \text{for } d > 2. \end{cases}$$

Proof. The first equation follows by 6.4.12. Moreover,

$$P[T_a < \infty] = \lim_{b \rightarrow \infty} P[T_a < T_b] = \begin{cases} 1 & \text{for } d \leq 2 \\ \phi(|x_0|)/\phi(a) & \text{for } d \geq 3. \end{cases}$$

□

Corollary 7.4. For a Brownian motion in \mathbb{R}^d the following statements hold for any initial value $x_0 \in \mathbb{R}^d$:

- (1). If $d \leq 2$ then every non-empty ball $D \subseteq \mathbb{R}^d$ is **recurrent**, i.e., the last visit time of D is almost surely infinite:

$$L_d = \sup\{t \geq 0 : B_t \in D\} = \infty \quad P\text{-a.s.}$$


- (2). If $d \geq 3$ then every ball D is **transient**, i.e.,

$$L_d < \infty \quad P\text{-a.s.}$$

- (3). If $d \geq 2$ then every point $x \in \mathbb{R}^d$ is **polar**, i.e.,

$$P[\exists t > 0 : B_t = x] = 0.$$

We point out that the last statement holds even if the starting point x_0 coincides with x . The first statement implies that a typical Brownian sample path is dense in \mathbb{R}^2 , whereas by the second statement, $\lim_{t \rightarrow \infty} |B_t| = \infty$ almost surely for $d \geq 3$.



Proof.

- (1),(2) The first two statements follow from Theorem 7.3 and the Markov property.

(3). For the third statement we assume w.l.o.g. $x = 0$. If $x_0 \neq 0$ then

$$P[T_0 < \infty] = \lim_{b \rightarrow \infty} P[T_0 < T_b]$$

for any $a > 0$. By Theorem 7.3,

$$P[T_0 < T_b] \leq \inf_{a > 0} P[T_a < T_b] = 0 \quad \text{for } d \geq 2,$$

whence $T_0 = \infty$ almost surely. If $x_0 = 0$ then by the Markov property,

$$P[\exists t > \varepsilon : B_t = 0] = E[P_{B_\varepsilon}[T_0 < \infty]] = 0$$

for any $\varepsilon > 0$. thus we again obtain

$$P[T_0 < \infty] = \lim_{\varepsilon \searrow 0} P[\exists t > \varepsilon : B_t = 0] = 0.$$

□

Remark (Polarity of linear subspaces). For $d \geq 2$, any $(d - 2)$ dimensional subspace $V \subseteq \mathbb{R}^d$ is polar for Brownian motion. For the proof note that the orthogonal projection of a one-dimensional Brownian motion onto the orthogonal complement V^\perp is a 2-dimensional Brownian motion.

7.2 Boundary value problems, exit and occupation times

The connection of Brownian motion to boundary value problems for partial differential equations involving the Laplace operator can be extended substantially:

The stationary Feynman-Kac-Poisson formula

Suppose that $f : \partial D \rightarrow \mathbb{R}$, $V : D \rightarrow \mathbb{R}$ and $g : D \rightarrow [0, \infty)$ are continuous functions defined on an open bounded domain $D \subset \mathbb{R}^d$, or on its boundary respectively. We assume that under P_x , (B_t) is Brownian motion with $P_x[B_0 = x] = 1$, and that

$$E_x \left[\exp \int_0^T V^-(B_s) ds \right] < \infty \quad \text{for any } x \in D, \quad (7.2.1)$$

where $T = T_{D^c}$ is the first exit time from D .

Note that (7.2.1) always holds if V is non-negative.

Theorem 7.5. *If $u \in C^2(D) \cap C(\bar{D})$ is a solution of the boundary problem*

$$\frac{1}{2}\Delta u(x) = V(x)u(x) - g(x) \quad \text{for } x \in D \quad (7.2.2)$$

$$u(x) = f(x) \quad \text{for } x \in \partial D, \quad (7.2.3)$$

and (7.2.1) holds then

$$u(x) = E_x \left[\exp \left(- \int_0^T V(B_s) ds \right) \cdot f(B_T) \right] + E_x \left[\int_0^T \exp \left(- \int_0^t V(B_s) ds \right) \cdot g(B_t) dt \right] \quad (7.2.4)$$

for any $x \in D$.

Remark. Note that we *assume* the existence of a smooth solution of the boundary value problem (7.2.2). Proving that the function u defined by (7.2.4) is a solution of the b.v.p. without assuming existence is much more demanding.

Proof. By continuity of V and (B_s) , the sample paths of the process

$$A_t = \int_0^t V(B_s) ds$$

are C^1 and hence of finite variation for $t < T$. Let

$$X_t = e^{-A_t} u(B_t), \quad t < T.$$

Applying Itô's formula with $F(a, b) = e^{-a}u(b)$ yields the decomposition

$$\begin{aligned} dX_t &= e^{-A_t} \nabla u(B_t) \cdot dB_t - e^{-A_t} u(B_t) dA_t + \frac{1}{2} e^{-A_t} \Delta u(B_t) dt \\ &= e^{-A_t} \nabla u(B_t) \cdot dB_t + e^{-A_t} \left(\frac{1}{2} \Delta u - V \cdot u \right) (B_t) dt \end{aligned}$$

of X_t into a local martingale up to time T and an absolutely continuous part. Since u is a solution of (7.2.2), we have $\frac{1}{2} \Delta u - Vu = -g$ on D . By applying the optional stopping theorem with a localizing sequence $T_n \nearrow T$ of stopping times, we obtain the representation

$$\begin{aligned} u(x) = E_x[X_0] &= E_x[X_{T_n}] + E_x \left[\int_0^{T_n} e^{-A_t} g(B_t) dt \right] \\ &= E_x[e^{-A_{T_n}} u(B_{T_n})] + E_x \left[\int_0^{T_n} e^{-A_t} g(B_t) dt \right] \end{aligned}$$

for $x \in D$. The assertion (7.2.4) now follows provided we can interchange the limit as $n \rightarrow \infty$ and the expectation values. For the second expectation on the right hand side this is possible by the monotone convergence theorem, because $g \geq 0$. For the first expectation value, we can apply the dominated convergence theorem, because

$$|e^{-A_{T_n}} u(B_{T_n})| \leq \exp \left(\int_0^T V^-(B_s) ds \right) \cdot \sup_{y \in \bar{D}} |u(y)| \quad \forall n \in \mathbb{N},$$

and the majorant is integrable w.r.t. each P_x by Assumption 7.2.1. □

Remark (Extension to diffusion processes). A corresponding result holds under appropriate assumptions if the Brownian motion (B_t) is replaced by a diffusion process (X_t) solving a stochastic differential equation of the type $dX_t = \sigma(X_t) dB_t + b(X_t) dt$, and the operator $\frac{1}{2} \Delta$ in (7.2.2) is replaced by the generator

$$\mathcal{L} = \frac{1}{2} \sum_{i,j=1}^d a_{i,j}(x) \frac{\partial^2}{\partial x_i \partial x_j} + b(x) \cdot \nabla, \quad a(x) = \sigma(x) \sigma(x)^\top,$$

of the diffusion process, cf. ???. The theorem hence establishes a general connection between Itô diffusions and boundary value problems for linear second order elliptic partial differential equations.

By Theorem 7.5 we can compute many interesting expectation values for Brownian motion by solving appropriate p.d.e. We now consider various corresponding applications.

Let us first recall the Dirichlet problem where $V \equiv 0$ and $g \equiv 0$. In this case, $u(x) = E_x[f(B_t)]$. We have already pointed out in the last section that this can be used to compute exit distributions and to study recurrence, transience and polarity of linear subspaces for Brownian motion in \mathbb{R}^d . A second interesting case of Theorem 7.5 is the stochastic representation for solutions of the Poisson equation:

Poisson problem and mean exit time

If V and f vanish in Theorem 7.3, the boundary value problem (7.2.2) reduces to the boundary value problem

$$\frac{1}{2}\Delta u = -g \quad \text{on } D, \quad u = 0 \quad \text{on } D,$$

for the Poisson equation. The solution has the stochastic representation

$$u(x) = E_x \left[\int_0^T g(B_t) dt \right], \quad x \in D, \quad (7.2.5)$$

which can be interpreted as an average cost accumulated by the Brownian path before exit from the domain D . In particular, choosing $g \equiv 1$, we can compute the mean exit time

$$u(x) = E_x[T]$$

from D for Brownian motion starting at x by solving the corresponding Poisson problem.

Example. If $D = \{x \in \mathbb{R}^d : |x| < r\}$ is a ball around 0 of radius $r > 0$, then the solution $u(x)$ of the Poisson problem

$$\frac{1}{2}\Delta u(x) = \begin{cases} -1 & \text{for } |x| < r \\ 0 & \text{for } |x| = r \end{cases}$$

can be computed explicitly. We obtain

$$E_x[T] = u(x) = \frac{r^2 - |x|^2}{d} \quad \text{for any } x \in D.$$

Occupation time density and Green function

If (B_t) is a Brownian motion in \mathbb{R}^d then the corresponding Brownian motion with absorption at the first exit time from the domain D is the Markov process (X_t) with state space $D \cup \{\Delta\}$ defined by

$$X_t = \begin{cases} B_t & \text{for } t < T \\ \Delta & \text{for } t \geq T \end{cases},$$

where Δ is an extra state added to the state space. By setting $g(\Delta) = 0$, the stochastic representation (7.2.5) for a solution of the Poisson problem can be written in the form

$$u(x) = E_x \left[\int_0^\infty g(X_t) dt \right] = \int_0^\infty (p_t^D g)(x) dt, \tag{7.2.6}$$

where

$$p_t^D(x, A) = P_x[X_t \in A], \quad A \subseteq \mathbb{R}^d \text{ measurable},$$

is the transition function for the absorbed process (X_t) . Note that for $A \subset \mathbb{R}^d$,

$$p_t^D(x, A) = P_x[B_t \in A \text{ and } t < T] \leq p_t(x, A) \tag{7.2.7}$$

where p_t is the transition function of Brownian motion on \mathbb{R}^d . For $t > 0$ and $x \in \mathbb{R}^d$, the transition function $p_t(x, \bullet)$ of Brownian motion is absolutely continuous. Therefore, by (7.2.7), the sub-probability measure $p_t^D(x, \bullet)$ restricted to \mathbb{R}^d is also absolutely continuous with non-negative density

$$p_t^D(x, y) \leq p_t(x, y) = (2\pi t)^{-d/2} \exp\left(-\frac{|x - y|^2}{2t}\right).$$

The function p_t^D is called the **heat kernel on the domain** D w.r.t. absorption on the boundary. Note that

$$G^D(x, y) = \int_0^\infty p_t^D(x, y) dt$$

is an **occupation time density**, i.e., it measures the average time a Brownian motion started in x spends in a small neighbourhood of y before it exits from the Domain

D . By (7.2.6), a solution u of the Poisson problem $\frac{1}{2}\Delta u = -g$ on D , $u = 0$ on ∂D , can be represented as

$$u(x) = \int_D G^D(x, y)g(y) dy \quad \text{for } x \in D.$$

This shows that the occupation time density $G^D(x, y)$ is the **Green function** (i.e., the fundamental solution of the Poisson equation) **for the operator $\frac{1}{2}$ with Dirichlet boundary conditions on the domain D .**

Note that although for domains with irregular boundary, the Green's function might not exist in the classical sense, the function $G^D(x, y)$ is always well-defined!

Stationary Feynman-Kac formula and exit time distributions

Next, we consider the case where g vanishes and $f \equiv 1$ in Theorem 7.5. Then the boundary value problem (7.2.4) takes the form

$$\frac{1}{2}\Delta u = Vu \quad \text{on } D, \quad u = 1 \quad \text{on } \partial D. \quad (7.2.8)$$

The p.d.e. $\frac{1}{2}\Delta u = Vu$ is a stationary Schrödinger equation. We will comment on the relation between the Feynman-Kac formula and Feynman's path integral formulation of quantum mechanics below. For the moment, we only note that for the solution of (??), the stochastic representation

$$u(x) = E_x \left[\exp \left(- \int_0^T V(B_t) dt \right) \right]$$

holds for $x \in D$.

As an application, we can, at least in principle, compute the full distribution of the exit time T . In fact, choosing $V \equiv \alpha$ for some constant $\alpha > 0$, the corresponding solution u_α of (7.2.8) yields the Laplace transform

$$u_\alpha(x) = E_x[e^{-\alpha T}] = \int_0^\infty e^{-\alpha t} \mu_x(dt) \quad (7.2.9)$$

of $\mu_x = P_x \circ T^{-1}$.

Example (Exit times in \mathbb{R}^1). Suppose $d = 1$ and $D = (-1, 1)$. Then (7.2.8) with $V = \alpha$ reads

$$\frac{1}{2}u''_{\alpha}(x) = \alpha u_{\alpha}(x) \quad \text{for } x \in (-1, 1), \quad u_{\alpha}(1) = u_{\alpha}(-1) = 1.$$

This boundary value problem has the unique solution

$$u_{\alpha}(x) = \frac{\cosh(x \cdot \sqrt{2\alpha})}{\cosh(\sqrt{2\alpha})} \quad \text{for } x \in [-1, 1].$$

By inverting the Laplace transform (7.2.9), one can now compute the distribution μ_x of the first exit time T from $(-1, 1)$. It turns out that μ_x is absolutely continuous with density

$$f_T(t) = \frac{1}{\sqrt{2\pi t^3}} \sum_{n=-\infty}^{\infty} \left((4n+1+x)e^{-\frac{(4n+1+x)^2}{2t}} + (4n+1-x)e^{-\frac{(4n+1-x)^2}{2t}} \right), \quad t \geq 0.$$




Figure 7.2: The density of the first exit time T depending on the starting point $x \in [-1, 1]$ and the time $t \in (0, 2]$.

Boundary value problems in \mathbb{R}^d and total occupation time

Suppose we would like to compute the distribution of the total occupation time

$$\int_0^\infty I_A(B_s) ds$$

of a bounded domain $A \subset \mathbb{R}^d$ for Brownian motion. This only makes sense for $d \geq 3$, since for $d \leq 2$, the total occupation time of any non-empty open set is almost surely infinite by recurrence of Brownian motion in \mathbb{R}^1 and \mathbb{R}^2 . The total occupation time is of the form $\int_0^\infty V(B_s) ds$ with $V = I_A$. Therefore, we should in principle be able to apply Theorem 7.3, but we have to replace the exit time T by $+\infty$ and hence the underlying bounded domain D by \mathbb{R}^d .

Corollary 7.6. *Suppose $d \geq 3$ and let $V : \mathbb{R}^d \rightarrow [0, \infty)$ be continuous. If $u \in C^2(\mathbb{R}^d)$ is a solution of the boundary value problem*

$$\frac{1}{2}\Delta u = Vu \quad \text{on } \mathbb{R}^d, \quad \lim_{|x| \rightarrow \infty} u(x) = 1 \quad (7.2.10)$$

then

$$u(x) = E_x \left[\exp \left(- \int_0^\infty V(B_t) dt \right) \right] \quad \text{for any } x \in \mathbb{R}^d.$$

Proof. Applying the stationary Feynman-Kac formula on an open bounded subset $D \subset \mathbb{R}^d$, we obtain the representation

$$u(x) = E_x \left[u(B_{T_{D^c}}) \exp \left(- \int_0^{T_{D^c}} V(B_t) dt \right) \right] \quad (7.2.11)$$

by Theorem 7.3. Now let $D_n = \{x \in \mathbb{R}^d : |x| < n\}$. Then $T_{D_n^c} \nearrow \infty$ as $n \rightarrow \infty$. Since $d \geq 3$, Brownian motion is transient, i.e., $\lim_{t \rightarrow \infty} |B_t| = \infty$, and therefore by (7.2.10)

$$\lim_{n \rightarrow \infty} u(B_{T_{D_n^c}}) = 1 \quad P_x\text{-almost surely for any } x.$$

Since u is bounded and V is non-negative, we can apply dominated convergence in (7.2.11) to conclude

$$u(x) = E_x \left[\exp \left(- \int_0^\infty V(B_t) dt \right) \right].$$

□

Let us now return to the computation of occupation time distributions. consider a bounded subset $A \subset \mathbb{R}^d, d \geq 3$, and let

$$v_\alpha(x) = E_x \left[\exp \left(-\alpha \int_0^\infty I_A(B_s) ds \right) \right], \quad \alpha > 0,$$

denote the Laplace transform of the total occupation time of A . Although $V = \alpha I_A$ is not a continuous function, a representation of v_α as a solution of a boundary problem holds:

Exercise. Prove that if $A \subset \mathbb{R}^d$ is a bounded domain with smooth boundary ∂A and $u_\alpha \in C^1(\mathbb{R}^d) \cap C^2(\mathbb{R}^d \setminus \partial A)$ satisfies

$$\frac{1}{2} \Delta u_\alpha = \alpha I_A u_\alpha \quad \text{on } \mathbb{R}^d \setminus \partial A, \quad \lim_{|x| \rightarrow \infty} u_\alpha(x) = 1, \quad (7.2.12)$$

then $v_\alpha = u_\alpha$.

Remark. The condition $u_\alpha \in C^1(\mathbb{R}^d)$ guarantees that u_α is a weak solution of the p.d.e. (7.2.10) on all of \mathbb{R}^d including the boundary ∂U .

Example (Total occupation time of the unit ball in \mathbb{R}^3). Suppose $A = \{x \in \mathbb{R}^3 : |x| < 1\}$. In this case the boundary value problem (7.2.10) is rotationally symmetric. The ansatz $u_\alpha(x) = f_\alpha(|x|)$, yields a Bessel equation for f_α on each of the intervals $(0, 1)$ and $(1, \infty)$:

$$\frac{1}{2} f_\alpha''(r) + r^{-1} f_\alpha'(r) = \alpha f_\alpha(r) \quad \text{for } r < 1, \quad \frac{1}{2} f_\alpha''(r) + r^{-1} f_\alpha(r) = 0 \quad \text{for } r > 1.$$

Taking into account the boundary condition and the condition $u_\alpha \in C^1(\mathbb{R}^d)$, one obtains the rotationally symmetric solution

$$u_\alpha(x) = \begin{cases} 1 + \left(\frac{\tanh(\sqrt{2\alpha})}{\sqrt{2\alpha}} - 1 \right) \cdot r^{-1} & \text{for } r \in [1, \infty), \\ \frac{\sinh(\sqrt{2\alpha}r)}{\sqrt{2\alpha} \cosh \sqrt{2\alpha}} \cdot r^{-1} & \text{for } r \in (0, 1) \\ \frac{1}{\cosh(\sqrt{2\alpha})} & \text{for } r = 0 \end{cases}.$$

of (7.2.10), and hence an explicit formula for v_α . In particular, for $x = 0$ we obtain the simple formula

$$E_0 \left[\exp \left(-\alpha \int_0^\infty I_A(B_t) dt \right) \right] = u_\alpha(0) = \frac{1}{\cosh(\sqrt{2\alpha})}.$$

The right hand side has already appeared in the example above as the Laplace transform of the exit time distribution of a one-dimensional Brownian motion starting at 0 from the interval $(-1, 1)$. Since the distribution is uniquely determined by its Laplace transform, we have proven the remarkable fact that the total occupation time of the unit ball for a standard Brownian motion in \mathbb{R}^3 has the same distribution as the first exit time from the unit ball for a standard one-dimensional Brownian motion:

$$\int_0^\infty I_{\{|B_t^{\mathbb{R}^3}| < 1\}} dt \sim \inf\{t > 0 : |B_t^{\mathbb{R}^3}| > 1\}.$$

This is a particular case of a theorem of Ciesielski and Taylor who proved a corresponding relation between Brownian motion in \mathbb{R}^{d+2} and \mathbb{R}^d for arbitrary d .

7.3 Heat Equation and Time-Dependent Feynman-Kac Formula

Itô's formula also yields a connection between Brownian motion (or, more generally, solutions of stochastic differential equations) and parabolic partial differential equations. The parabolic p.d.e. are Kolmogorov forward or backward equations for the corresponding Markov processes. In particular, the time-dependent Feynman-Kac formula shows

that the backward equation for Brownian motion with absorption is a heat equation with dissipation.

Brownian Motion with Absorption

Suppose we would like to describe the evolution of a Brownian motion that is absorbed during the evolution of a Brownian motion that is absorbed during an infinitesimal time interval $[t, t + dt]$ with probability $V(t, x)dt$ where x is the current position of the process. We assume that the *absorption rate* $V(t, x)$ is given by a measurable locally-bounded function

$$V : [0, \infty) \times \mathbb{R}^d \rightarrow [0, \infty).$$

Then the accumulated absorption rate up to time t is given by the increasing process

$$A_t = \int_0^t V(s, B_s) ds, \quad t \geq 0.$$

We can think of the process A_t as an internal clock for the Brownian motion determining the absorption time. More precisely, we define:

Definition. Suppose that $(B_t)_{t \geq 0}$ is a d -dimensional Brownian motion and T is a with parameter 1 exponentially distributed random variable independent of (B_t) . Let Δ be a separate state added to the state space \mathbb{R}^d . Then the process (X_t) defined by

$$X_t := \begin{cases} B_t & \text{for } A_t < T, \\ \Delta & \text{for } A_t \geq T, \end{cases}$$

is called a **Brownian motion with absorption rate** $V(t, x)$, and the random variable

$$\zeta := \inf\{t \geq 0 : A_t \geq T\}$$

is called the **absorption time**.

A justification for the construction is given by the following informal computation: For an infinitesimal time interval $[t, t + dt]$ and almost every ω ,

$$\begin{aligned} P[\zeta \leq t + dt \mid (B_s)_{s \geq 0}, \zeta > t](\omega) &= P[A_{t+dt}(\omega) \geq T \mid A_t(\omega) < T] \\ &= P[A_{t+dt}(\omega) - A_t(\omega) \geq T] \\ &= P[V(t, B_t(\omega))dt \geq T] \\ &= V(t, B_t(\omega))dt \end{aligned}$$

by the memoryless property of the exponential distribution, i.e., $V(t, x)$ is indeed the infinitesimal absorption rate.

Rigorously, it is not difficult to verify that (X_t) is a Markov process with state space $\mathbb{R}^d \cup \{\Delta\}$ where Δ is an absorbing state. The Markov process is time-homogeneous if $V(t, x)$ is independent of t .

For a measurable subset $D \subseteq \mathbb{R}^d$ and $t \geq 0$ the distribution μ_t of X_t is given by

$$\begin{aligned} \mu_t[D] &= P[X_t \in D] = P[B_t \in D \text{ and } A_t < T] \\ &= E[P[A_t < T \mid (B_t)]; B_t \in D] \\ &= E \left[\exp \left(- \int_0^t V(s, B_s) ds \right) ; B_t \in D \right]. \end{aligned} \tag{7.3.1}$$

Itô's formula can be used to prove a Kolmogorov type forward equation:

Theorem 7.7 (Forward equation for Brownian motion with absorption). *The sub-probability measures μ_t on \mathbb{R}^d solve the heat equation*

$$\frac{\partial \mu_t}{\partial t} = \frac{1}{2} \Delta \mu_t - V(t, \bullet) \mu_t \tag{7.3.2}$$

in the following distributional sense:

$$\int f(x) \mu_t(dx) - \int f(x) \mu_0(dx) = \int_0^t \int \left(\frac{1}{2} \Delta f(x) - V(s, x) f(x) \right) \mu_s(dx) ds$$

for any function $f \in C_0^2(\mathbb{R}^d)$.

Here $C_0^2(\mathbb{R}^d)$ denotes the space of C^2 -functions with compact support. Under additional regularity assumptions it can be shown that μ_t has a smooth density that solves (7.3.1) in the classical sense. The equation (7.3.1) describes heat flow with cooling when the heat at x at time t dissipates with rate $V(t, x)$.

Proof. By (7.3.1),

$$\int f d\mu_t = E[\exp(-A_t); f(B_t)] \quad (7.3.3)$$

for any bounded measurable function $f : \mathbb{R}^d \rightarrow \mathbb{R}$. For $f \in C_0^2(\mathbb{R}^d)$, an application of Itô's formula yields

$$e^{-A_t} f(B_t) = f(B_0) + M_t + \int_0^t e^{-A_s} f(B_s) V(s, B_s) ds + \frac{1}{2} \int_0^t e^{-A_s} \Delta f(B_s) ds,$$

for $t \geq 0$, where (M_t) is a local martingale. Taking expectation values for a localizing sequence of stopping times and applying the dominated convergence theorem subsequently, we obtain

$$E[e^{-A_t} f(B_t)] = E[f(B_0)] + \int_0^t E[e^{-A_s} (\frac{1}{2} \Delta f - V(s, \bullet) f)(B_s)] ds.$$

Here we have used that $\frac{1}{2} \Delta f(x) - V(s, x) f(x)$ is uniformly bounded for $(s, x) \in [0, t] \times \mathbb{R}^d$, because f has compact support and V is locally bounded. The assertion now follows by (7.3.3). \square

Exercise (Heat kernel and Green's function). The transition kernel for Brownian motion with time-homogeneous absorption rate $V(x)$ restricted to \mathbb{R}^d is given by

$$p_t^V(x, D) = E_x \left[\exp \left(- \int_0^t V(B_s) ds \right) ; B_t \in D \right].$$

- (1). Prove that for any $t > 0$ and $x \in \mathbb{R}^d$, the sub-probability measure $p_t^V(x, \bullet)$ is absolutely continuous on \mathbb{R}^d with density satisfying

$$0 \leq p_t^V(x, y) \leq (2\pi t)^{-d/2} \exp(-|x - y|^2/(2t)).$$

(2). Identify the occupation time density

$$G^V(x, y) = \int_0^\infty p_t^V(x, y) dt$$

as a fundamental solution of an appropriate boundary value problem. Adequate regularity may be assumed.

Time-dependent Feynman-Kac formula

In Theorem 7.7 we have applied Itô's formula to prove a Kolmogorov type forward equation for Brownian motion with absorption. To obtain a corresponding backward equation, we have to reverse time:

Theorem 7.8 (Feynman-Kac). Fix $t > 0$, and let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ and $V, g : [0, t] \times \mathbb{R}^d \rightarrow \mathbb{R}$ be continuous functions. Suppose that f is bounded, g is non-negative, and V satisfies

$$E_x \left[\exp \int_0^t V(t-s, B_s) ds \right] < \infty \quad \text{for all } x \in \mathbb{R}^d. \quad (7.3.4)$$

If $u \in C^{1,2}((0, t] \times \mathbb{R}^d) \cap C([0, t] \times \mathbb{R}^d)$ is a bounded solution of the heat equation

$$\frac{\partial u}{\partial s}(s, x) = \frac{1}{2} \Delta u(s, x) - V(s, x)u(s, x) + g(s, x) \quad \text{for } s \in (0, t], x \in \mathbb{R}^d, \quad (7.3.5)$$

$$u(0, x) = f(x),$$

then u has the stochastic representation

$$u(t, x) = E_x \left[f(B_t) \exp \left(- \int_0^t V(t-s, B_s) ds \right) \right] + E_x \left[\int_0^t g(t-r, B_r) \exp \left(- \int_0^r V(t-s, B_s) ds \right) dr \right].$$

Remark. The equation (7.3.5) describes heat flow with sinks and dissipation.

Proof. We first reverse time on the interval $[0, t]$. The function

$$\hat{u}(s, x) = u(t - s, x)$$

solves the p.d.e.

$$\begin{aligned} \frac{\partial \hat{u}}{\partial s}(s, x) &= -\frac{\partial u}{\partial t}(t - s, x) = -\left(\frac{1}{2}\Delta u - Vu + g\right)(t - s, x) \\ &= -\left(\frac{1}{2}\Delta \hat{u} - \hat{V}\hat{u} + \hat{g}\right)(s, x) \end{aligned}$$

on $[0, t]$ with terminal condition $\hat{u}(t, x) = f(x)$. Now let $X_r = \exp(-A_r)\hat{u}(r, B_r)$ for $r \in [0, t]$, where

$$A_r := \int_0^r \hat{V}(s, B_s) ds = \int_0^r V(t - s, B_s) ds.$$

By Itô's formula, we obtain for $\tau \in [0, t]$,

$$\begin{aligned} X_\tau - X_0 &= M_\tau - \int_0^\tau e^{-A_r} \hat{u}(r, B_r) dA_r + \int_0^\tau e^{-A_r} \left(\frac{\partial \hat{u}}{\partial s} + \frac{1}{2}\Delta \hat{u}\right)(r, B_r) dr \\ &= M_\tau + \int_0^\tau e^{-A_r} \left(\frac{\partial \hat{u}}{\partial s} + \frac{1}{2}\Delta \hat{u} - \hat{V}\hat{u}\right)(r, B_r) dr \\ &= M_\tau - \int_0^\tau e^{-A_r} \hat{g}(r, B_r) dr \end{aligned}$$

with a local martingale $(M_\tau)_{\tau \in [0, t]}$ vanishing at 0. Choosing a corresponding localizing sequence of stopping times T_n with $T_n \nearrow t$, we obtain by the optional stopping theorem and by dominated convergence,

$$\begin{aligned} u(t, x) &= \hat{u}(0, x) = E_x[X_0] \\ &= E_x[X_t] + E_x \left[\int_0^t e^{-A_r} \hat{g}(r, B_r) dr \right] \\ &= E_x[e^{-A_t} u(0, B_t)] + E_x \left[\int_0^t e^{-A_r} g(t - r, B_r) dr \right]. \end{aligned}$$

□

Remark (Extension to diffusion processes). Again a similar result holds under a appropriate regularity assumptions for Brownian motion replaced by a solution of a s.d.e. $dX_t = \sigma(X_t)dB_t + b(X_t)dt$ and $\frac{1}{2}\Delta$ replaced by the corresponding generator, cf. ??.

Occupation times and arc-sine law

The Feynman-Kac formula can be used to study the distribution of occupation times of Brownian motion. We consider an example where the distribution can be computed explicitly: The proportion of time during the interval $[0, t]$ spent by a one-dimensional standard Brownian motion (B_t) in the interval $(0, \infty)$. Let

$$A_t = \lambda(\{s \in [0, t] : B_s > 0\}) = \int_0^t I_{(0, \infty)}(B_s) ds.$$

Theorem 7.9 (Arc-sine law of P.Lévy). For any $t > 0$ and $\theta \in [0, 1]$,

$$P_0[A_t/t \leq \theta] = \frac{2}{\pi} \arcsin \sqrt{\theta} = \frac{1}{\pi} \int_0^\theta \frac{ds}{\sqrt{s(1-s)}}.$$




Figure 7.3: Density of A_t/t .

Note that the theorem shows in particular that a law of large numbers does **not** hold! Indeed, for each $\varepsilon > 0$,

$$P_0 \left[\left| \frac{1}{t} \int_0^t I_{(0,\infty)}(B_s) ds - \frac{1}{2} \right| > \varepsilon \right] \not\rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Even for large times, values of A_t/t close to 0 or 1 are the most probable. By the functional central limit theorem, the proportion of time that one player is ahead in a long coin tossing game or a counting of election results is also close to the arcsine law. In particular, it is more than 20 times more likely that one player is ahead for more than 98% of the time than it is that each player is ahead between 49% and 51% of the time [Steele].

Before proving the arc-sine law, we give an informal derivation based on the time-dependent Feynman-Kac formula.

The idea for determining the distribution of A_t is again to consider the Laplace transforms

$$u(t, x) = E_x[\exp(-\beta A_t)], \quad \beta > 0.$$

By the Feynman-Kac formula, we could expect that u solves the equation

$$\frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial x^2} \tag{7.3.6}$$

with initial condition $u(0, x) = 1$. To solve the parabolic p.d.e. (7.3.6), we consider another Laplace transform: The Laplace transform

$$v_\alpha(x) = \int_0^\infty e^{-\alpha t} u(t, x) dt = E_x \left[\int_0^\infty e^{-\alpha t - \beta A_t} dt \right], \quad \alpha > 0,$$

of a solution $u(t, x)$ of (7.3.6) w.r.t. t . An informal computation shows that v_α should satisfy the o.d.e.

$$\begin{aligned} \frac{1}{2} v_\alpha'' - \beta I_{(0,\infty)} v_\alpha &= \int_0^\infty e^{-\alpha t} \left(\frac{1}{2} \frac{\partial^2 u}{\partial x^2} - \beta I_{(0,\infty)} u \right) (t, \bullet) dt \\ &= \int_0^\infty e^{-\alpha t} \frac{\partial u}{\partial t} (t, \bullet) dt = e^{-\alpha t} u(t, \bullet) \Big|_0^\infty - \alpha \int_0^\infty e^{-\alpha t} u(t, \bullet) dt \\ &= 1 - \alpha v_\alpha, \end{aligned}$$

i.e., v_α should be a bounded solution of

$$\alpha v_\alpha - \frac{1}{2}v_\alpha'' + \beta I_{0,\infty}v_\alpha = g \quad (7.3.7)$$

where $g(x) = 1$ for all x . The solution of (7.3.7) can then be computed explicitly, and yield the arc-sine law by Laplace inversion.

Remark. The method of transforming a parabolic p.d.e. by the Laplace transform into an elliptic equation is standard and used frequently. In particular, the Laplace transform of a transition semigroup $(p_t)_{t \geq 0}$ is the corresponding resolvent $(g_\alpha)_{\alpha \geq 0}$, $g_\alpha = \int_0^\infty e^{-\alpha t} p_t dt$, which is crucial for potential theory.

Instead of trying to make the informal argument above rigorous, one can directly prove the arc-sine law by applying the stationary Feynman-Kac formula:

Exercise. Prove Lévy's arc-sine law by proceeding in the following way:

- (1). Let $g \in C_b(\mathbb{R})$. Show that if v_α is a bounded solution of (7.3.7) on $\mathbb{R} \setminus \{0\}$ with $v_\alpha \in C^1(\mathbb{R}) \cap C^2(\mathbb{R} \setminus \{0\})$ then

$$v_\alpha(x) = E_x \left[\int_0^\infty g(B_t) e^{-\alpha t - \beta A_t} dt \right] \quad \text{for any } x \in \mathbb{R}.$$

- (2). Compute a corresponding solution v_α for $g \equiv 1$, and conclude that

$$\int_0^\infty e^{-\alpha t} E_0[e^{-\beta A_t}] dt = \frac{1}{\sqrt{\alpha(\alpha + \beta)}}.$$

- (3). Now use the uniqueness of the Laplace inversion to show that the distribution μ_t of A_t/t under P_\bullet is absolutely continuous with density

$$f_{A_t/t}(s) = \frac{1}{\pi \sqrt{s \cdot (1 - s)}}.$$

Chapter 8

Stochastic Differential Equations: Explicit Computations

Suppose that $(B_t)_{t \geq 0}$ is a given Brownian motion defined on a probability space (Ω, \mathcal{A}, P) . We will now study solutions of stochastic differential equations (SDE) of type

$$dX_t = b(t, X_t) dt + \sigma(t, X_t) dB_t \quad (8.0.1)$$

where b and σ are continuous functions defined on $\mathbb{R}_+ \times \mathbb{R}^d$ or an appropriate subset.

Recall that $\mathcal{F}_t^{B,P}$ denotes the completion of the filtration $\mathcal{F}_t^B = \sigma(B_s \mid 0 \leq s \leq t)$ generated by the Brownian motion. Let T be an $(\mathcal{F}_t^{B,P})$ stopping time. We call a process $(t, \omega) \mapsto X_t(\omega)$ defined for $t < T(\omega)$ **adapted w.r.t.** $(\mathcal{F}_t^{B,P})$, if the trivially extended process $\tilde{X}_t = X_t \cdot I_{\{t < T\}}$ defined by

$$\tilde{X}_t := \begin{cases} X_t & \text{for } t < T \\ 0 & \text{for } t \geq T \end{cases},$$

is $(\mathcal{F}_t^{B,P})$ -adapted.

Definition. An almost surely continuous stochastic process $(t, \omega) \mapsto X_t(\omega)$ defined for $t \in [0, T(\omega))$ is called a **strong solution** of the stochastic differential equation (8.0.1) if it is $(\mathcal{F}_t^{B,P})$ -adapted, and the equation

$$X_t = X_0 + \int_0^t b(s, X_s) ds + \int_0^t \sigma(s, X_s) dB_s \quad \text{for } t \in [0, T) \quad (8.0.2)$$

holds P -almost surely.

The terminology “strong” solution will be explained later when we introduce “weak” solutions. The point is that a strong solution is adapted w.r.t. the filtration $(\mathcal{F}_t^{B,P})$ generated by the Brownian motion. Therefore, a strong solution is essentially (up to modification on measure zero sets) a *measurable function of the given Brownian motion*! The concept of strong and weak solutions of SDE is not related to the analytic definition of strong and weak solutions for partial differential equations.

In this section we study properties of solutions and we compute explicit solutions for one-dimensional SDE. We start with an example:

Example (Asset price model in continuous time). A nearby model for an asset price process $(S_n)_{n=0,1,2,\dots}$ in discrete time is to define S_n recursively by

$$S_{n+1} - S_n = \alpha_n(S_0, \dots, S_n)S_n + \sigma_n(S_0, \dots, S_n)S_n\eta_{n+1}$$

with i.i.d. random variables $\eta_i, i \in \mathbb{N}$, and measurable functions $\alpha_n, \sigma_n : \mathbb{R}^n \rightarrow \mathbb{R}$. Trying to set up a corresponding model in continuous time, we arrive at the stochastic differential equation

$$dS_t = \alpha_t S_t dt + \sigma_t S_t dB_t \quad (8.0.3)$$

with an (\mathcal{F}_t) -Brownian motion (B_t) and (\mathcal{F}_t^P) adapted continuous stochastic processes $(\alpha_t)_{t \geq 0}$ and $(\sigma_t)_{t \geq 0}$, where (\mathcal{F}_t) is a given filtration on a probability space (Ω, \mathcal{A}, P) . The processes α_t and σ_t describe the *instantaneous mean rate of return* and the *volatility*. Both are allowed to be time dependent and random.

In order to compute a solution of (8.0.3), we assume $S_t > 0$ for any $t \geq 0$, and divide the equation by S_t :

$$\frac{1}{S_t} dS_t = \alpha_t dt + \sigma_t dB_t. \quad (8.0.4)$$

We will prove in Section 8.1 that if an SDE holds then the SDE multiplied by a continuous adapted process also holds, cf. Theorem 8.1. Hence (8.0.4) is equivalent to (8.0.3) if $S_t > 0$. If (8.0.4) would be a classical ordinary differential equation then we could use the identity $d \log S_t = \frac{1}{S_t} dS_t$ to solve the equation. In Itô calculus, however, the classical chain rule is violated. Nevertheless, it is still useful to compute $d \log S_t$ by Itô's formula. The process (S_t) has quadratic variation

$$[S]_t = \left[\int_0^\bullet \sigma_r S_r dB_r \right]_t = \int_0^t \sigma_r^2 S_r^2 dr \quad \text{for any } t \geq 0,$$

almost surely along an appropriate sequence (π_n) of partitions with $\text{mesh}(\pi_n) \rightarrow 0$. The first equation holds by (8.0.3), since $t \mapsto \int_0^t \alpha_r S_r dr$ has finite variation, and the second identity is proved in Theorem 8.1 below. Therefore, Itô's formula implies:

$$\begin{aligned} d \log S_t &= \frac{1}{S_t} dS_t - \frac{1}{2S_t^2} d[S]_t \\ &= \alpha_t dt + \sigma_t dB_t - \frac{1}{2} \sigma_t^2 dt \\ &= \mu_t dt + \sigma_t dB_t, \end{aligned}$$

where $\mu_t := \alpha_t - \sigma_t^2/2$, i.e.,

$$\log S_t - \log S_0 = \int_0^t \mu_s ds + \int_0^t \sigma_s dB_s,$$

or, equivalently,

$$S_t = S_0 \cdot \exp \left(\int_0^t \mu_s ds + \int_0^t \sigma_s dB_s \right). \quad (8.0.5)$$

Conversely, one can verify by Itô's formula that (S_t) defined by (8.0.5) is indeed a solution of (8.0.3). Thus we have proven existence, uniqueness and an explicit representation for a strong solution of (8.0.3). In the special case when $\alpha_t \equiv \alpha$ and $\sigma_t \equiv \sigma$ are constants in t and ω , the solution process

$$S_t = S_0 \exp(\sigma B_t + (\alpha - \sigma^2/2)t)$$

is called a **geometric Brownian motion with parameters α and σ** .




Figure 8.1: Three one dimensional geometric Brownian motions with $\alpha^2 = 1$ and $\sigma = 0.1$ (blue), $\sigma = 1.0$ (red) and $\sigma = 2.0$ (magenta).

8.1 Stochastic Calculus for Itô processes

By definition, any solution of an SDE of the form (8.0.1) is the sum of an absolutely continuous adapted process and an Itô stochastic integral w.r.t. the underlying Brownian motion, i.e.,

$$X_t = A_t + I_t \quad \text{for } t < T, \quad (8.1.1)$$

where

$$A_t = \int_0^t K_s ds \quad \text{and} \quad I_t = \int_0^t H_s dB_s \quad (8.1.2)$$

with $(H_t)_{t < T}$ and $(K_t)_{t < T}$ almost surely continuous and $(\mathcal{F}_t^{B,P})$ -adapted. A stochastic process of type (8.1.1) is called an **Itô process**. In order to compute and analyze

solutions of SDE we will apply Itô's formula to Itô processes. Since the absolutely continuous process (A_t) has finite variation, classical Stieltjes calculus applies to this part of an Itô process. It remains to consider the stochastic integral part (I_t) :

Stochastic integrals w.r.t. Itô processes

Let (π_n) be a sequence of partitions of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$. Recall that for $t \geq 0$,

$$I_t = \int_0^t H_s dB_s = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} H_s \cdot (B_{s' \wedge t} - B_s)$$

w.r.t. convergence in probability on $\{t < T\}$, cf. Theorem 5.14.

Theorem 8.1 (Composition rule and quadratic variation). *Suppose that T is a predictable stopping time and $(H_t)_{t < T}$ is almost surely continuous and adapted.*

(1). *For any almost surely continuous, adapted process $(G_t)_{0 \leq t < T}$, and for any $t \geq 0$,*

$$\lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} G_s (I_{s' \wedge t} - I_s) = \int_0^t G_s H_s dB_s \quad (8.1.3)$$

with convergence in probability on $\{t < T\}$. Moreover, if H is in $\mathcal{L}_a^2([0, a])$ and G is bounded on $[0, a] \times \Omega$ for some $a > 0$, then the convergence holds in $M_c^2([0, a])$ and thus uniformly for $t \in [0, a]$ in the $L^2(P)$ sense.

(2). *For any $t \geq 0$, the quadratic variation $[I]_t$ along (π_n) is given by*

$$[I]_t = \lim_{n \rightarrow \infty} \sum_{\substack{s \in \pi_n \\ s < t}} (I_{s' \wedge t} - I_s)^2 = \int_0^t H_s^2 ds \quad (8.1.4)$$

w.r.t. convergence in probability on $\{t < T\}$.

XXX gleich $[I, J]$ berechnen, Beweis analog

Remark (Uniform convergence). Similarly to the proof of Theorem 5.14 one can show that there is a sequence of bounded stopping times $T_k \nearrow T$ such that almost surely along a subsequence, the convergence in (8.1.3) and (8.1.4) holds uniformly on $[0, T_k]$ for any $k \in \mathbb{N}$.

Proof. (1). We first fix $a > 0$ and assume that H is in $\mathcal{L}_a^2([0, a])$ and G is bounded, left-continuous and adapted on $[0, \infty) \times \Omega$. Since $I_{s' \wedge t} - I_s = \int_s^{s' \wedge t} H_r dB_r$, we obtain

$$\sum_{\substack{s \in \pi_n \\ s < t}} G_s (I_{s' \wedge t} - I_s) = \int_0^t G_{[r]_n} H_r dB_r$$

where $[r]_n = \max\{s \in \pi_n : s \leq r\}$ is the next partition point below r . As $n \rightarrow \infty$, the right-hand side converges to $\int_0^t G_r H_r dB_r$ in $M_c^2([0, a])$ because $G_{[r]_n} H_r \rightarrow G_r H_r$ in $L^2(P \otimes \lambda_{[0, a]})$ by continuity of G and dominated convergence.

The assertion in the general case now follows by localization: Suppose (S_k) and (T_k) are increasing sequences of stopping times with $T_k \nearrow T$ and $H_t I_{\{t \leq S_k\}} \in \mathcal{L}_a^2([0, \infty))$, and let

$$\tilde{T}_k = S_k \wedge T_k \wedge \inf\{t \geq 0 : |G_t| > k\} \wedge k.$$

Then $\tilde{T}_k \nearrow T$, the process $H_t^{(k)} := H_t I_{\{t \leq T_k\}}$ is in $\mathcal{L}_a^2([0, \infty))$ the process $G_t^{(k)} := G_t I_{\{t \leq T_k\}}$ is bounded, left-continuous and adapted, and

$$I_s = \int_0^s H_r^{(k)} dB_r, \quad G_s = G_s^{(k)} \quad \text{for any } s \in [0, t]$$

holds almost surely on $\{t \leq \tilde{T}_k\}$. Therefore as $n \rightarrow \infty$,

$$\begin{aligned} \sum_{\substack{s \in \pi_n \\ s < t}} G_s (I_{s' \wedge t} - I_s) &= \sum_{\substack{s \in \pi_n \\ s < t}} G_s^{(k)} (I_{s' \wedge t} - I_s) \\ &\rightarrow \int_0^t G_r^{(k)} H_r^{(k)} dB_r = \int_0^t G_r H_r dB_r \end{aligned}$$

uniformly for $t \leq \tilde{T}_k$ in $L^2(P)$. The claim follows, since

$$P \left[\{t < T\} \setminus \bigcup_k \{t \leq \tilde{T}_k\} \right] = 0.$$

(2). We first assume that H is in $\mathcal{L}_a^2([0, \infty))$, continuous and bounded. Then for $s \in \pi_n$,

$$\delta I_s = I_{s' \wedge t} - I_s = \int_s^{s' \wedge t} H_r dB_r = H_s \delta B_s + R_s^{(n)}$$

where $R_s^{(n)} := \int_s^{s' \wedge t} (H_r - H_{[r]_n}) dB_r$. Therefore,

$$\sum_{\substack{s \in \pi_n \\ s < t}} (\delta I_s)^2 = \sum_{\substack{s \in \pi_n \\ s < t}} H_s^2 (\delta B_s)^2 + 2 \sum_{\substack{s \in \pi_n \\ s < t}} R_s^{(n)} H_s \delta B_s + \sum_{\substack{s \in \pi_n \\ s < t}} (R_s^{(n)})^2.$$

Since $[B]_t = t$ almost surely, the first term on the right-hand side converges to $\int_0^t H_s^2 ds$ with probability one. It remains to show that the remainder terms converge to 0 in probability as $n \rightarrow \infty$. This is the case, since

$$\begin{aligned} E \left[\sum (R_s^{(n)})^2 \right] &= \sum E[(R_s^{(n)})^2] = \sum \int_s^{s' \wedge t} E[(H_r - H_{[r]_n})^2] dr \\ &= \int_0^t E[(H_r - H_{[r]_n})^2] dr \longrightarrow 0 \end{aligned}$$

by the Itô isometry and continuity and boundedness of H , whence $\sum (R_s^{(n)})^2 \rightarrow 0$ in \mathcal{L}^1 and in probability, and $\sum R_s^{(n)} H_s \delta B_s \rightarrow 0$ in the same sense by the Schwarz inequality.

For H defined up to a stopping time T , the assertion now follows by a localization procedure similar to the one applied above. □

The theorem and the corresponding composition rule for Stieltjes integrals suggest that we may define stochastic integrals w.r.t. an Itô process

$$X_t = X_0 + \int_0^t H_s dB_s + \int_0^t K_s ds, \quad t < T,$$

in the following way:

Definition. Suppose that (B_t) is a Brownian motion on (Ω, \mathcal{A}, P) w.r.t. a filtration (\mathcal{F}_t) , X_0 is an (\mathcal{F}_0^P) -measurable random variable, T is a predictable (\mathcal{F}_t^P) -stopping time, and (G_t) , (H_t) and (K_t) are almost surely continuous, (\mathcal{F}_t^P) adapted processes defined for $t < T$. Then the stochastic integral of (G_t) w.r.t. (X_t) is the Itô process defined by

$$\int_0^t G_s dX_s = \int_0^t G_s H_s dB_s + \int_0^t G_s K_s ds, \quad t < T.$$

By Theorem 8.1, this definition is consistent with a definition by Riemann sum approximations. Moreover, the definition shows that the class of Itô processes w.r.t. a given Brownian motion is *closed under taking stochastic integrals!* In particular, strong solutions of SDE w.r.t. Itô processes are again Itô processes.

Calculus for Itô processes

We summarize calculus rules for Itô processes that are immediate consequences of the definition above and Theorem 8.1: Suppose that (X_t) and (Y_t) are Itô processes, and (G_t) , (\tilde{G}_t) and (H_t) are adapted continuous process that are all defined up to a stopping time T . Then the following calculus rules hold for Itô stochastic differentials:

Linearity:

$$\begin{aligned} d(X + cY) &= dX + c dY && \text{for any } c \in \mathbb{R}, \\ (G + cH) dX &= G dX + cH dX && \text{for any } c \in \mathbb{R}. \end{aligned}$$

Composition rule:

$$dY = G dX \quad \Rightarrow \quad \tilde{G} dY = \tilde{G} G dX,$$

Quadratic variation:

$$dY = G dX \quad \Rightarrow \quad d[Y] = G^2 d[X],$$

Itô rule: For any function $F \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R})$,

$$dF(t, X) = \frac{\partial F}{\partial x}(t, X) dX + \frac{\partial F}{\partial t}(t, X) dt + \frac{1}{2} \frac{\partial^2 F}{\partial x^2}(t, X) d[X]$$

All equations are to be understood in the sense that the corresponding stochastic integrals over any interval $[0, t]$, $t < T$, coincide almost surely.

The proofs are straightforward. For example, if

$$Y_t = Y_0 + \int_0^t G_s dX_s$$

and

$$X_t = X_0 + \int_0^t K_s ds + \int_0^t H_s dB_s$$

then, by the definition above, for $t < T$,

$$Y_t = Y_0 + \int_0^t G_s K_s ds + \int_0^t G_s H_s dB_s,$$

and hence

$$\int_0^t \tilde{G}_s dY_s = \int_0^t \tilde{G}_s G_s K_s ds + \int_0^t \tilde{G}_s G_s H_s dB_s = \int_0^t \tilde{G}_s G_s dX_s$$

and

$$[Y]_t = \left[\int_0^\bullet G_s H_s dB_s \right]_t = \int_0^t G_s^2 H_s^2 ds = \int_0^t G_s^2 d[X]_s.$$

Moreover, Theorem 8.1 guarantees that the stochastic integrals in Itô's formula (which are limits of Riemann-Itô sums) coincide with the stochastic integrals w.r.t. Itô processes defined above.

Example (Option Pricing in continuous time I). We again consider the continuous time asset price model introduced in the beginning of Chapter 8. Suppose an agent is holding ϕ_t units of a single asset with price process (S_t) at time t , and he invests the remainder $V_t - \phi_t S_t$ of his wealth V_t in the money market with interest rate R_t . We

assume that (ϕ_t) and (R_t) are continuous adapted processes. Then the change of wealth in a small time unit should be described by the Itô equation

$$dV_t = \phi_t dS_t + R_t(V_t - \phi_t S_t) dt.$$

Similarly to the discrete time case, we consider the discounted wealth process

$$\tilde{V}_t := \exp\left(-\int_0^t R_s ds\right) V_t.$$

Since $t \mapsto \int_0^t R_s ds$ has finite variation, the Itô rule and the composition rule for stochastic integrals imply:

$$\begin{aligned} d\tilde{V}_t &= \exp\left(-\int_0^t R_s ds\right) dV_t - \exp\left(-\int_0^t R_s ds\right) R_t V_t dt \\ &= \exp\left(-\int_0^t R_s ds\right) \phi_t dS_t - \exp\left(-\int_0^t R_s ds\right) R_t \phi_t S_t dt \\ &= \phi_t \cdot \left(\exp\left(-\int_0^t R_s ds\right) dS_t - \exp\left(-\int_0^t R_s ds\right) R_t S_t dt\right) \\ &= \phi_t d\tilde{S}_t, \end{aligned}$$

where \tilde{S}_t is the discounted asset price process. Therefore,

$$\tilde{V}_t - \tilde{V}_0 = \int_0^t \phi_s d\tilde{S}_s \quad \forall t \geq 0 \quad P\text{-almost surely.}$$

As a consequence, we observe that if (\tilde{S}_t) is a (local) martingale under a probability measure P_* that is equivalent to P then the discounted wealth process (\tilde{V}_t) is also a local martingale under P_* . A corresponding probability measure P_* is called an *equivalent martingale measure* or *risk neutral measure*, and can be identified by Girsanov's theorem, cf. Section 9.3 below. Once we have found P_* , option prices can be computed similarly as in discrete time under the additional assumption that the true measure P for the asset price process is equivalent to P_* , see Section 9.4.

The Itô-Doebelin formula in \mathbb{R}^1

We will now apply Itô's formula to solutions of stochastic differential equations. Let $b, \sigma \in C(\mathbb{R}_+ \times I)$ where $I \subseteq \mathbb{R}$ is an open interval. Suppose that (B_t) is an (\mathcal{F}_t) -Brownian motion on (Ω, \mathcal{A}, P) , and $(X_t)_{0 \leq t < T}$ is an (\mathcal{F}_t^P) -adapted process with values in I and defined up to an (\mathcal{F}_t^P) stopping time T such that the SDE

$$X_t - X_0 = \int_0^t b(s, X_s) ds + \int_0^t \sigma(s, X_s) dB_s \quad \text{for any } t < T \quad (8.1.5)$$

holds almost surely.

Corollary 8.2 (Doebelin 1941, Itô 1944). *Let $F \in C^{1,2}(\mathbb{R}_+ \times I)$. Then almost surely,*

$$\begin{aligned} F(t, X_t) - F(0, X_0) &= \int_0^t (\sigma F')(s, X_s) dB_s \\ &\quad + \int_0^t \left(\frac{\partial F}{\partial t} + \frac{1}{2} \sigma^2 F'' + b F' \right) (s, X_s) ds \quad \text{for any } t < T, \end{aligned} \quad (8.1.6)$$

where $F' = \partial F / \partial x$ denotes the partial derivative w.r.t. x .

Proof. Let (π_n) be a sequence of partitions with $\text{mesh}(\pi_n) \rightarrow 0$. Since the process $t \mapsto X_0 + \int_0^t b(s, X_s) ds$ has sample paths of locally finite variation, the quadratic variation of (X_t) is given by

$$[X]_t = \left[\int_0^\bullet \sigma(s, X_s) dB_s \right]_t = \int_0^t \sigma(s, X_s)^2 ds \quad \forall t < T$$

w.r.t. almost sure convergence along a subsequence of (π_n) . Hence Itô's formula can be applied to almost every sample path of (X_t) , and we obtain

$$\begin{aligned} F(t, X_t) - F(0, X_0) &= \int_0^t F'(s, X_s) dX_s + \int_0^t \frac{\partial F}{\partial t}(s, X_s) ds + \frac{1}{2} \int_0^t F''(s, X_s) d[X]_s \\ &= \int_0^t (\sigma F')(s, X_s) dB_s + \int_0^t (bF')(s, X_s) ds + \int_0^t \frac{\partial F}{\partial t}(s, X_s) ds + \frac{1}{2} \int_0^t (\sigma^2 F'')(s, X_s) ds \end{aligned}$$

for all $t < T$, P -almost surely. Here we have used (8.1.5) and the fact that the Itô integral w.r.t. X is an almost sure limit of Riemann-Itô sums after passing once more to an appropriate subsequence of (π_n) . \square

Exercise (Black Scholes partial differential equation). A stock price is modeled by a geometric Brownian Motion (S_t) with parameters $\alpha, \sigma > 0$. We assume that the interest rate is equal to a real constant r for all times. Let $c(t, x)$ be the value of an option at time t if the stock price at that time is $S_t = x$. Suppose that $c(t, S_t)$ is replicated by a hedging portfolio, i.e., there is a trading strategy holding ϕ_t shares of stock at time t and putting the remaining portfolio value $V_t - \phi_t S_t$ in the money market account with fixed interest rate r so that the total portfolio value V_t at each time t agrees with $c(t, S_t)$.

“Derive” the *Black-Scholes partial differential equation*

$$\frac{\partial c}{\partial t}(t, x) + rx \frac{\partial c}{\partial x}(t, x) + \frac{1}{2} \sigma^2 x^2 \frac{\partial^2 c}{\partial x^2}(t, x) = rc(t, x) \quad (8.1.7)$$

and the *delta-hedging rule*

$$\phi_t = \frac{\partial c}{\partial x}(t, S_t) \quad (=:\text{Delta}). \quad (8.1.8)$$

Hint: Consider the discounted portfolio value $\tilde{V}_t = e^{-rt}V_t$ and, correspondingly, the discounted option value $e^{-rt}c(t, S_t)$. Compute the Ito differentials, and conclude that both processes coincide if c is a solution to (8.1.7) and ϕ_t is given by (8.1.8).

Martingale problem for solutions of SDE

The Itô-Doeblin formula shows that if (X_t) is a solution of (8.1.5) then

$$M_t^F = F(t, X_t) - F(0, X_0) - \int_0^t (\mathcal{L}F)(s, X_s) ds$$

is a local martingale up to T for any $F \in C^{1,2}(\mathbb{R}_+ \times I)$ and

$$(\mathcal{L}F)(t, x) = \frac{1}{2}\sigma(t, x)^2 F''(t, x) + b(t, x)F'(t, x).$$

In particular, in the time-homogeneous case and for $T = \infty$, any solution of (8.1.5) solves the martingale problem for the operator $\mathcal{L}F = \frac{1}{2}\sigma^2 F'' + bF'$ with domain $C_0^2(I)$. Similarly as for Brownian motion, the martingales identified by the Itô-Doeblin formula can be used to compute various expectation values for the Itô diffusion (X_t) . In the next section we will look at first examples.

Remark (Uniqueness and Markov property of strong solutions). If the coefficients are, for example, Lipschitz continuous, then the strong solution of the SDE (8.1.5) is unique, and it has the strong Markov property, i.e., it is a diffusion process in the classical sense (a strong Markov process with continuous sample paths). By the Itô-Doeblin formula, the generator of this Markov process is an extension of the operator $(\mathcal{L}, C_0^2(I))$.

Although in general, uniqueness and the Markov property may not hold for solutions of the SDE (8.1.5), we call any solution of this equation an **Itô diffusion**.

8.2 Stochastic growth

In this section we consider time-homogeneous Itô diffusions taking values in the interval $I = (0, \infty)$. They provide natural models for stochastic growth processes, e.g. in mathematical biology, financial mathematics and many other application fields. Analogue results also hold if I is replaced by an arbitrary non-empty open interval.

Suppose that $(X_t)_{0 \leq t < T}$ is a strong solution of the SDE

$$\begin{aligned} dX_t &= b(X_t) dt + \sigma(X_t) dB_t & \text{for } t \in [0, T), \\ X_0 &= x_0, \end{aligned}$$

with a given Brownian motion (B_t) , $x_0 \in (0, \infty)$, and continuous time-homogeneous coefficients $b, \sigma : (0, \infty) \rightarrow \mathbb{R}$. We assume that the solution is defined up to the explosion time

$$T = \sup_{\varepsilon, r > 0} T_{\varepsilon, r}, \quad T_{\varepsilon, r} = \inf\{t \geq 0 \mid X_t \notin (\varepsilon, r)\}.$$

The corresponding generator is

$$\mathcal{L}F = bF' + \frac{1}{2}\sigma^2 F''.$$

Before studying some concrete models, we show in the general case how harmonic functions can be used to compute exit distributions (e.g. ruin probabilities) and to analyze the asymptotic behaviour of X_t as $t \nearrow T$.

Scale functions and exit distributions

To determine the exit distribution from a finite subinterval $(\varepsilon, r) \subset (0, \infty)$ we compute the harmonic functions of \mathcal{L} . For $h \in C^2(0, \infty)$ with $h' > 0$ we obtain:

$$\mathcal{L}h = 0 \quad \iff \quad h'' = -\frac{2b}{\sigma^2}h' \quad \iff \quad (\log h')' = -\frac{2b}{\sigma^2}.$$

Therefore, the two-dimensional vector space of harmonic functions is spanned by the constant function 1 and by the function

$$s(x) = \int_{x_0}^x \exp\left(-\int_{x_0}^z \frac{2b(y)}{\sigma(y)^2} dy\right) dz.$$

$s(x)$ is called a **scale function** of the process (X_t) . It is strictly increasing and harmonic on $(0, \infty)$. Hence we can think of $s : (0, \infty) \rightarrow (s(0), s(\infty))$ as a coordinate transformation, and the transformed process $s(X_t)$ is a local martingale up to the explosion time T . Applying the martingale convergence theorem and the optional stopping theorem to $s(X_t)$ one obtains:

Theorem 8.3. *For any $\varepsilon, r \in (0, \infty)$ with $\varepsilon < x_0 < r$ we have:*

(1). The exit time $T_{\varepsilon,r} = \inf\{t \in [0, T) : X_t \notin (\varepsilon, r)\}$ is almost surely less than T .

$$(2). P[T_\varepsilon < T_r] = P[X_{T_{\varepsilon,r}} = \varepsilon] = \frac{s(r) - s(x)}{s(r) - s(\varepsilon)}.$$

The proof of Theorem 8.3 is left as an exercise.

Remark. (1). Note that any affine transformation $\tilde{s}(x) = cs(x) + d$ with constants $c > 0$ and $d \in \mathbb{R}$ is also harmonic and strictly increasing, and hence a scale function. The ratio $(s(r) - s(x))/(s(r) - s(\varepsilon))$ is invariant under non-degenerate affine transformations of s .

(2). The scale function and the ruin probabilities depend only on the ratio $b(x)/\sigma(x)^2$.

Recurrence and asymptotics

We now apply the formula for the exit distributions in order to study the asymptotics of one-dimensional non-degenerate Itô diffusions as $t \nearrow T$. For $\varepsilon \in (0, x_0)$ we obtain

$$\begin{aligned} P[T_\varepsilon < T] &= P[T_\varepsilon < T_r \text{ for some } r \in (x_0, \infty)] \\ &= \lim_{r \rightarrow \infty} P[T_\varepsilon < T_r] = \lim_{r \rightarrow \infty} \frac{s(r) - s(x_0)}{s(r) - s(\varepsilon)}. \end{aligned}$$

In particular, we have

$$P[X_t = \varepsilon \text{ for some } t \in [0, T)] = P[T_\varepsilon < T] = 1$$

if and only if $s(\infty) = \lim_{r \nearrow \infty} s(r) = \infty$.

Similarly, one obtains for $r \in (x_0, \infty)$:

$$P[X_t = r \text{ for some } t \in [0, T)] = P[T_r < T] = 1$$

if and only if $s(0) = \lim_{\varepsilon \searrow 0} s(\varepsilon) = -\infty$.

Moreover,

$$P[X_t \rightarrow \infty \text{ as } t \nearrow T] = P\left[\bigcup_{\varepsilon > 0} \bigcap_{r < \infty} \{T_r < T_\varepsilon\}\right] = \lim_{\varepsilon \searrow 0} \lim_{r \nearrow \infty} \frac{s(x_0) - s(\varepsilon)}{s(r) - s(\varepsilon)},$$

and

$$P[X_t \rightarrow 0 \text{ as } t \nearrow T] = P\left[\bigcup_{r < \infty} \bigcap_{\varepsilon > 0} \{T_\varepsilon < T_r\}\right] = \lim_{r \nearrow \infty} \lim_{\varepsilon \searrow 0} \frac{s(x_0) - s(\varepsilon)}{s(r) - s(\varepsilon)}.$$

Summarizing, we have shown:

Corollary 8.4 (Asymptotics of one-dimensional Itô diffusions).

(1). If $s(0) = -\infty$ and $s(\infty) = \infty$, then the process (X_t) is recurrent, i.e.,

$$P[X_t = y \text{ for some } t \in [0, T)] = 1 \quad \text{for any } x_0, y \in (0, \infty).$$

(2). If $s(0) > -\infty$ and $s(\infty) = \infty$ then $\lim_{t \nearrow T} X_t = 0$ almost surely.

(3). If $s(0) = -\infty$ and $s(\infty) < \infty$ then $\lim_{t \nearrow T} X_t = \infty$ almost surely.

(4). If $s(0) > -\infty$ and $s(\infty) < \infty$ then

$$P\left[\lim_{t \nearrow T} X_t = 0\right] = \frac{s(\infty) - s(x_0)}{s(\infty) - s(0)}$$

and

$$P\left[\lim_{t \nearrow T} X_t = \infty\right] = \frac{s(x_0) - s(0)}{s(\infty) - s(0)}$$

Intuitively, if $s(0) = -\infty$, in the natural scale the boundary is transformed to $-\infty$, which is not a possible limit for the local martingale $s(X_t)$, whereas otherwise $s(0)$ is finite and approached by $s(X_t)$ with strictly positive probability.

Example. Suppose that $b(x)/\sigma(x)^2 \approx \gamma x^{-1}$ as $x \nearrow \infty$ and $b(x)/\sigma(x)^2 \approx \delta x^{-1}$ as $x \searrow 0$ holds for $\gamma, \delta \in \mathbb{R}$ in the sense that $b(x)/\sigma(x)^2 - \gamma x^{-1}$ is integrable at ∞ and $b(x)/\sigma(x)^2 - \delta x^{-1}$ is integrable at 0. Then $s'(x)$ is of order $x^{-2\gamma}$ as $x \nearrow \infty$ and of order $x^{-2\delta}$ as $x \searrow 0$. Hence

$$s(\infty) = \infty \iff \gamma \leq \frac{1}{2}, \quad s(0) = -\infty \iff \delta \geq \frac{1}{2}.$$

In particular, recurrence holds if and only if $\gamma \leq \frac{1}{2}$ and $\delta \geq \frac{1}{2}$.

More concrete examples will be studied below.

Remark (Explosion in finite time, Feller's test). Corollary 8.4 does not tell us whether the explosion time T is infinite with probability one. It can be shown that this is always the case if (X_t) is recurrent. In general, *Feller's test for explosions* provides a necessary and sufficient condition for the absence of explosion in finite time. The idea is to compute a function $g \in C(0, \infty)$ such that $e^{-t}g(X_t)$ is a local martingale and to apply the optional stopping theorem. The details are more involved than in the proof of corollary above, cf. e.g. Section 6.2 in [Durrett: Stochastic calculus].

Geometric Brownian motion

A geometric Brownian motion with parameters $\alpha \in \mathbb{R}$ and $\sigma > 0$ is a solution of the s.d.e.

$$dS_t = \alpha S_t dt + \sigma S_t dB_t. \quad (8.2.1)$$

We have already shown in the beginning of Section ?? that for $B_0 = 0$, the unique strong solution of (8.2.1) with initial condition $S_0 = x_0$ is

$$S_t = x_0 \cdot \exp(\sigma B_t + (\alpha - \sigma^2/2)t).$$

The distribution of S_t at time t is a **lognormal distribution**, i.e., the distribution of $c \cdot e^Y$ where c is a constant and Y is normally distributed. Moreover, one easily verifies that (S_t) is a time-homogeneous Markov process with log-normal transition densities

$$p_t(x, y) = \frac{1}{\sqrt{2\pi t\sigma^2}} \exp\left(-\frac{(\log(y/x) - \mu t)^2}{2t\sigma^2}\right), \quad t, x, y > 0,$$

where $\mu = \alpha - \sigma^2/2$. By the Law of Large Numbers for Brownian motion,

$$\lim_{t \rightarrow \infty} S_t = \begin{cases} +\infty & \text{if } \mu > 0 \\ 0 & \text{if } \mu < 0 \end{cases}.$$

If $\mu = 0$ then (S_t) is recurrent since the same holds for (B_t) .

We now convince ourselves that we obtain the same results via the scale function:

The ratio of the drift and diffusion coefficient is

$$\frac{b(x)}{\sigma(x)^2} = \frac{\alpha x}{(\sigma x)^2} = \frac{\alpha}{\sigma^2 x},$$

and hence

$$s'(x) = \text{const.} \cdot \exp\left(-\int_{x_0}^x \frac{2\alpha}{\sigma^2 y} dy\right) = \text{const.} \cdot x^{-2\alpha/\sigma^2}.$$

Therefore,

$$s(\infty) = \infty \iff 2\alpha/\sigma^2 \leq 1, \quad s(0) = \infty \iff 2\alpha/\sigma^2 \geq 1,$$

which again shows that $S_t \rightarrow \infty$ for $\alpha > \sigma^2/2$, $S_t \rightarrow 0$ for $\alpha < \sigma^2/2$, and S_t is recurrent for $\alpha = \sigma^2/2$.

Feller's branching diffusion

Our second growth model is described by the stochastic differential equation

$$dX_t = \beta X_t dt + \sigma \sqrt{X_t} dB_t, \quad X_0 = x_0, \quad (8.2.2)$$

with given constants $\beta \in \mathbb{R}, \sigma > 0$, and values in \mathbb{R}_+ . Note that in contrast to the equation of geometric Brownian motion, the multiplicative factor $\sqrt{X_t}$ in the noise term is not a linear function of X_t . As a consequence, there is no explicit formula for a solution of (8.2.2). Nevertheless, a general existence result guarantees the existence of a strong solution defined up to the explosion time

$$T = \sup_{\varepsilon, r > 0} T_{\mathbb{R} \setminus (\varepsilon, r)},$$

cf. ???. SDEs similar to (8.2.2) appear in various applications.

Example (Diffusion limits of branching processes). We consider a Galton-Watson branching process Z_t^h with time steps $t = 0, h, 2h, 3h, \dots$ of size $h > 0$, i.e., Z_0^h is a given initial population size, and

$$Z_{t+h}^h = \sum_{i=1}^{Z_t^h} N_i, t/h \quad \text{for } t = k \cdot h, k = 0, 1, 2, \dots,$$

with independent identically distributed random variables $N_{i,k}, i \geq 1, k \geq 0$. The random variable Z_{kh}^h describes the size of a population in the k -th generation when $N_{i,l}$

is the number of offspring of the i -th individual in the l -th generation. We assume that the mean and the variance of the offspring distribution are given by

$$E[N_{i,l}] = 1 + \beta h \quad \text{and} \quad \text{Var}[N_{i,l}] = \sigma^2$$

for finite constants $\beta, \sigma \in \mathbb{R}$.

We are interested in a scaling limit of the model as the size h of time steps goes to 0. To establish convergence to a limit process as $h \searrow 0$ we rescale the population size by h , i.e., we consider the process

$$X_t^h := h \cdot Z_{\lfloor t \rfloor}^h, \quad t \in [0, \infty).$$

The mean growth (“drift”) of this process in one time step is

$$E[X_{t+h}^h - X_t^h \mid \mathcal{F}_t^h] = h \cdot E[Z_{t+h}^h - Z_t^h \mid \mathcal{F}_t^h] = h\eta h Z_t^h = h\beta X_t^h,$$

and the corresponding condition variance is

$$\text{Var}[X_{t+h}^h - X_t^h \mid \mathcal{F}_t^h] = h^2 \cdot \text{Var}[Z_{t+h}^h - Z_t^h \mid \mathcal{F}_t^h] = h^2 \sigma^2 Z_t^h = h\sigma^2 X_t^h,$$

where $\mathcal{F}_t^h = \sigma(N_{i,l} \mid i \geq 1, 0 \leq l \leq k)$ for $t = k \cdot h$. Since both quantities are of order $O(h)$, we can expect a limit process (X_t) as $h \searrow 0$ with drift coefficient $\beta \cdot X_t$ and diffusion coefficient $\sqrt{\sigma^2 X_t}$, i.e., the scaling limit should be a diffusion process solving a s.d.e. of type (8.2.2). A rigorous derivation of this diffusion limit can be found e.g. in Section 8 of [Durrett: Stochastic Calculus].

We now analyze the asymptotics of solutions of (8.2.2). The ratio of drift and diffusion coefficient is $\beta x / (\sigma \sqrt{x})^2 = \beta / \sigma$, and hence the derivative of a scale function is

$$s'(x) = \text{const.} \cdot \exp(-2\beta x / \sigma).$$

Thus $s(0)$ is always finite, and $s(\infty) = \infty$ if and only if $\beta \leq 1$. Therefore, by Corollary 8.4, in the subcritical and critical case $\beta \leq 1$, we obtain

$$\lim_{t \nearrow T} X_t = 0 \quad \text{almost surely,}$$

whereas in the supercritical case $\beta > 1$,

$$P \left[\lim_{t \nearrow T} X_t = 0 \right] > 0 \quad \text{and} \quad P \left[\lim_{t \nearrow T} X_t = \infty \right] > 0.$$

This corresponds to the behaviour of Galton-Watson processes in discrete time. It can be shown by Feller's boundary classification for one-dimensional diffusion processes that if $X_t \rightarrow 0$ then the process actually dies out almost surely in finite time, cf. e.g. Section 6.5 in [Durrett: Stochastic Calculus]. On the other hand, for trajectories with $X_t \rightarrow \infty$, the explosion time T is almost surely infinite and X_t grows exponentially as $t \rightarrow \infty$.

Cox-Ingersoll-Ross model

The CIR model is a model for the stochastic evolution of interest rates or volatilities. The equation is

$$dR_t = (\alpha - \beta R_t) dt + \sigma \sqrt{R_t} dB_t \quad R_0 = x_0, \quad (8.2.3)$$

with a one-dimensional Brownian motion (B_t) and positive constants $\alpha, \beta, \sigma > 0$. Although the s.d.e. looks similar to the equation for Feller's branching diffusion, the behaviour of the drift coefficient near 0 is completely different. In fact, the idea is that the positive drift α pushes the process away from 0 so that a recurrent process on $(0, \infty)$ is obtained. We will see that this intuition is true for $\alpha \geq \sigma^2/2$ but not for $\alpha < \sigma^2/2$.

Again, there is no explicit solution for the s.d.e. (8.13), but existence of a strong solution holds. The ratio of the drift and diffusion coefficient is $(\alpha - \beta x)/\sigma^2 x$, which yields

$$s'(x) = \text{const.} \cdot x^{-2\alpha/\sigma^2} e^{2\beta x/\sigma^2}.$$

Hence $s(\infty) = \infty$ for any $\beta > 0$, and $s(0) = \infty$ if and only if $2\alpha \geq \sigma^2$. Therefore, the CIR process is recurrent if and only if $\alpha \geq \sigma^2/2$, whereas $X_t \rightarrow 0$ as $t \nearrow T$ almost surely otherwise.

By applying Itô's formula one can now prove that X_t has finite moments, and compute the expectation and variance explicitly. Indeed, taking expectation values in the s.d.e.

$$R_t = x_0 + \int_0^t (\alpha - \beta R_s) ds + \int_0^t \sigma \sqrt{R_s} dB_s,$$

we obtain informally

$$\frac{d}{dt}E[R_t] = \alpha - \beta E[R_t],$$

and hence by variation of constants,

$$E[R_t] = x_0 \cdot e^{-\beta t} + \frac{\alpha}{\beta}(1 - e^{-\beta t}).$$

To make this argument rigorous requires proving that the local martingale $t \mapsto \int_0^t \sigma \sqrt{R_s} dB_s$ is indeed a martingale:

Exercise. Consider a strong solution $(R_t)_{t \geq 0}$ of (8.13) for $\alpha \geq \sigma^2/2$.

- (1). Show by applying Itô's formula to $x \mapsto |x|^p$ that $E[|R_t|^p] < \infty$ for any $t \geq 0$ and $p \geq 1$.
- (2). Compute the expectation of R_t , e.g. by applying Itô's formula to $e^{\beta t}x$.
- (3). Proceed in a similar way to compute the variance of R_t . Find its asymptotic value $\lim_{t \rightarrow \infty} \text{Var}[R_t]$.

8.3 Linear SDE with additive noise

We now consider stochastic differential equations of the form

$$dX_t = \beta_t C_t dt + \sigma_t dB_t, \quad X_0 = x, \quad (8.3.1)$$

where (B_t) is a Brownian motion, and the coefficients are *deterministic* continuous functions $\beta, \sigma : [0, \infty) \rightarrow \mathbb{R}$. Hence the drift term $\beta_t X_t$ is linear in X_t , and the diffusion coefficient does not depend on X_t , i.e., the noise increment $\sigma_t dB_t$ is proportional to white noise dB_t with a proportionality factor that does not depend on X_t .

Variation of constants

An explicit strong solution of the SDE (8.3.1) can be computed by a “variation of constants” Ansatz. We first note that the general solution in the deterministic case $\sigma_t \equiv 0$ is given by

$$X_t = \text{const.} \cdot \exp \left(\int_0^t \beta_s ds \right).$$

To solve the SDE in general we try the ansatz

$$X_t = C_t \cdot \exp \left(\int_0^t \beta_s ds \right)$$

with a continuous Itô process (C_t) driven by the Brownian motion (B_t) . By the Itô product rule,

$$dX_t = \beta_t X_t dt + \exp \left(\int_0^t \beta_s ds \right) dC_t.$$

Hence (X_t) solves (8.3.1) if and only if

$$dC_t = \exp \left(- \int_0^t \beta_s ds \right) \sigma_t dB_t,$$

i.e.,

$$C_t = C_0 + \int_0^t \exp \left(- \int_0^r \beta_s ds \right) \sigma_r dB_r.$$

We thus obtain:

Theorem 8.5. *The almost surely unique strong solution of the SDE (8.3.1) with initial value x is given by*

$$X_t^x = x \cdot \exp \left(- \int_0^t \beta_s ds \right) + \int_0^t \exp \left(\int_r^t \beta_s ds \right) \sigma_r dB_r.$$

Note that the theorem not only yields an explicit solution but it also shows that the solution depends smoothly on the initial value x . The effect of the noise on the solution is additive and given by a Wiener-Itô integral, i.e., an Itô integral with deterministic integrand. The average value

$$E[X_t^x] = x \cdot \exp\left(\int_0^t B_s ds\right), \quad (8.3.2)$$

coincides with the solution in the absence of noise, and the mean-square deviation from this solution due to random perturbation of the equation is

$$\text{Var}[X_t^x] = \text{Var}\left[\int_0^t \exp\left(\int_r^t \beta_s ds\right) \sigma_r dB_r\right] = \int_0^t \exp\left(2 \int_r^t \beta_s ds\right) \sigma_r^2 dr$$

by the Itô isometry.

Solutions as Gaussian processes

We now prove that the solution (X_t) of a linear s.d.e. with additive noise is a Gaussian process. We first observe that X_t is normally distributed for any $t \geq 0$.

Lemma 8.6. *For any deterministic function $h \in L^2(0, t)$, the Wiener-Itô integral $I_t = \int_0^t h_s dB_s$ is normally distributed with mean 0 and variance $\int_0^t h_s^2 ds$.*

Proof. Suppose first that $h = \sum_{i=0}^{n-1} c_i \cdot I_{(t_i, t_{i+1}]}$ is a step function with $n \in \mathbb{N}$, $c_1, \dots, c_n \in \mathbb{R}$, and $0 \leq t_0 < t_1 < \dots < t_n$. Then $I_t = \sum_{i=0}^{n-1} c_i \cdot (B_{t_{i+1}} - B_{t_i})$ is normally distributed with mean zero and variance

$$\text{Var}[I_t] = \sum_{i=0}^{n-1} c_i^2 (t_{i+1} - t_i) = \int_0^t h_s^2 ds.$$

In general, there exists a sequence $(h^{(n)})_{n \in \mathbb{N}}$ of step functions such that $h^{(n)} \rightarrow h$ in $L^2(0, t)$, and

$$I_t = \int_0^t h dB = \lim_{n \rightarrow \infty} \int_0^t h^{(n)} dB \quad \text{in } L^2(\Omega, \mathcal{A}, P).$$

Hence I_t is again normally distributed with mean zero and

$$\text{Var}[I_t] = \lim_{n \rightarrow \infty} \text{Var} \left[\int_0^t h^{(n)} dB \right] = \int_0^t h^2 ds.$$

□

Theorem 8.7 (Wiener-Itô integrals are Gaussian processes). *Suppose that $h \in L^2_{loc}([0, \infty), \mathbb{R})$. Then $I_t = \int_0^t h_s dB_s$ is a continuous Gaussian process with*

$$E[I_t] = 0 \quad \text{and} \quad \text{Cov}[I_t, I_s] = \int_0^{t \wedge s} h_r^2 ds \quad \text{for any } t, s \geq 0.$$

Proof. Let $0 \leq t_1 < \dots < t_n$. To show that $(I_{t_1}, \dots, I_{t_n})$ has a normal distribution it suffices to prove that any linear combination of the random variables I_{t_1}, \dots, I_{t_n} is normally distributed. This holds true since any linear combination is again an Itô integral with deterministic integrand:

$$\sum_{i=1}^n \lambda_i I_{t_i} = \int_0^{t_n} \sum_{i=1}^n \lambda_i \cdot I_{(0, t_i)}(s) h_s dB_s$$

for any $n \in \mathbb{N}$ and $\lambda_1, \dots, \lambda_n \in \mathbb{R}$. Hence (I_t) is a Gaussian process with $E[I_t] = 0$ and

$$\begin{aligned} \text{Cov}[I_t, I_s] &= E[I_t I_s] \\ &= E \left[\int_0^\infty h_r \cdot I_{(0, t)}(r) dB_r \int_0^\infty h_r \cdot I_{(0, s)}(r) dB_r \right] \\ &= (h \cdot I_{(0, t)}, h \cdot I_{(0, s)})_{L^2(0, \infty)} \\ &= \int_0^{s \wedge t} h_r^2 dr. \end{aligned}$$

□

Example (Brownian motion). If $h \equiv 1$ then $I_t = B_t$. The Brownian motion (B_t) is a centered Gaussian process with $\text{Cov}[B_t, B_s] = t \wedge s$.

More generally, by Theorem 8.7 and Theorem 8.5, any solution (X_t) of a linear SDE with additive noise and deterministic (or Gaussian) initial value is a continuous Gaussian process. In fact by (8.3.1), the marginals of (X_t) are affine functions of the corresponding marginals of a Wiener-Itô integral:

$$X_t^x = \frac{1}{h_t} \cdot \left(x + \int_0^t h_r \sigma_r dB_r \right) \quad \text{with} \quad h_r = \exp \left(- \int_0^r \beta_u du \right).$$

Hence all finite dimensional marginals of (X_t^x) are normally distributed with

$$E[X_t^x] = x/H_t \quad \text{and} \quad \text{Cov}[X_t^x, X_s^x] = \frac{1}{h_t h_s} \cdot \int_0^{t \wedge s} h_r^2 \sigma_r^2 dr.$$

The Ornstein-Uhlenbeck process

In 1905, Einstein introduced a model for the movement of a “big” particle in a fluid. Suppose that V_t^{abs} is the absolute velocity of the particle, \bar{V}_t is the mean velocity of the fluid molecules and $V_t = V_t^{\text{abs}} - \bar{V}_t$ is the velocity of the particle relative to the fluid. Then the velocity approximatively can be described as a solution to an s.d.e.

$$dV_t = -\gamma V_t dt + \sigma dB_t. \tag{8.3.3}$$

Here (B_t) is a Brownian motion in \mathbb{R}^d , $d = 3$, and γ, σ are strictly positive constants that describe the damping by the viscosity of the fluid and the magnitude of the random collisions. A solution to the s.d.e. (8.3.3) is called an **Ornstein-Uhlenbeck process**. Although it has first been introduced as a model for the velocity of physical Brownian motion, the Ornstein-Uhlenbeck process is a fundamental stochastic process that is almost as important as Brownian motion for mathematical theory and stochastic modeling. In particular, it is a continuous-time analogue of an AR(1) autoregressive process. Note that (8.3.3) is a system of d decoupled one-dimensional stochastic differential equations $dV_t^{(i)} = -\gamma V_t^{(i)} dt + \sigma dB_t^{(i)}$. Therefore, we will assume w.l.o.g. $d = 1$. By the considerations above, the one-dimensional Ornstein-Uhlenbeck process is a continuous

Gaussian process. The unique strong solution of the s.d.e. (8.3.3) with initial condition x is given explicitly by

$$V_t^x = e^{-\gamma t} \left(x + \sigma \int_0^t e^{\gamma s} dB_s \right). \quad (8.3.4)$$

In particular,

$$E[V_t^x] = e^{-\gamma t} x,$$

and

$$\begin{aligned} \text{Cov}[V_t^x, V_s^x] &= e^{-\gamma(t+s)} \sigma^2 \int_0^{t \wedge s} e^{2\gamma r} dr \\ &= \frac{\sigma^2}{2\gamma} (e^{-\gamma|t-s|} - e^{-\gamma(t+s)}) \quad \text{for any } t, s \geq 0. \end{aligned}$$

Note that as $t \rightarrow \infty$, the effect of the initial condition decays exponentially fast with rate γ . Similarly, the correlations between V_t^x and V_s^x decay exponentially as $|t-s| \rightarrow \infty$. The distribution at time t is

$$V_t^x \sim N \left(e^{-\gamma t} x, \frac{\sigma^2}{2\gamma} (1 - e^{-2\gamma t}) \right). \quad (8.3.5)$$

In particular, as $t \rightarrow \infty$

$$V_t^x \xrightarrow{\mathcal{D}} N \left(0, \frac{\sigma^2}{2\gamma} \right).$$

One easily verifies that $N(0, \sigma^2/2\gamma)$ is an *equilibrium* for the process: If $V_0 \sim N(0, \sigma^2/2\gamma)$ and (B_t) is independent of V_0 then

$$\begin{aligned} V_t &= e^{-\gamma t} V_0 + \sigma \int_0^t e^{\gamma(s-t)} dB_s \\ &\sim N \left(0, \frac{\sigma^2}{2\gamma} e^{-2\gamma t} + \sigma^2 \int_0^t e^{2\gamma(s-t)} ds \right) = N(0, \sigma^2/2\gamma) \end{aligned}$$

for any $t \geq 0$.

Theorem 8.8. *The Ornstein-Uhlenbeck process (V_t^x) is a time-homogeneous Markov process w.r.t. the filtration $(\mathcal{F}_t^{B,P})$ with stationary distribution $N(0, \sigma^2/2\gamma)$ and transition probabilities*

$$p_t(x, A) = P \left[e^{-\gamma t} x + \frac{\sigma}{\sqrt{2\gamma}} \sqrt{1 - e^{-2\gamma t}} Z \in A \right], \quad Z \sim N(0, 1).$$

Proof. We first note that by (8.3.5),

$$V_t^x \sim e^{-\gamma t} x + \frac{\sigma}{\sqrt{2\gamma}} \sqrt{1 - e^{-2\gamma t}} Z \quad \text{for any } t \geq 0$$

with $Z \sim N(0, 1)$. Hence,

$$E[f(V_t^x)] = (p_t f)(x)$$

for any non-negative measurable function $f : \mathbb{R} \rightarrow \mathbb{R}$. We now prove a *pathwise counterpart to the Markov property*: For $t, r \geq 0$, by (8.3.4)

$$\begin{aligned} V_{t+r}^x &= e^{-\gamma(t+r)} \left(x + \sigma \int_0^t e^{\gamma s} dB_s \right) + \sigma \int_0^{t+r} e^{\gamma(s-t-r)} dB_s \\ &= e^{-\gamma r} V_t^x + \sigma \int_0^r e^{\gamma(u-r)} d\bar{B}_u, \end{aligned} \quad (8.3.6)$$

where $\bar{B}_u := B_{t+u} - B_t$ is a Brownian motion that is independent of $\mathcal{F}_t^{B,P}$. Hence, the random variable $\sigma \cdot \int_0^r e^{\gamma(u-r)} d\bar{B}_u$ is also independent of $\mathcal{F}_t^{B,P}$ and, by (8.3.4), it has the same distribution as the Ornstein-Uhlenbeck process with initial condition 0:

$$\sigma \cdot \int_0^r e^{\gamma(u-r)} d\bar{B}_u \sim V_r^0.$$

Therefore, by (8.3.6), the conditional distribution of V_{t+r}^x given $\mathcal{F}_t^{B,P}$ coincides with the distribution of the process with initial V_t^x at time r :

$$\begin{aligned} E[f(V_{t+r}^x) | \mathcal{F}_t^{B,P}] &= E[f(e^{-\gamma r} V_t^x(\omega) + V_r^0)] \\ &= E[f(V_r^{V_t^x(\omega)})] = (p_r f)(V_t^x(\omega)) \quad \text{for } P\text{-a.e. } \omega. \end{aligned}$$

This proves that (V_t^x) is a Markov process with transition kernels $p_r, r \geq 0$. \square

Remark. The pathwise counterpart of the Markov property used in the proof above is called **cocycle property** of the stochastic flow $x \mapsto V_t^x$.

The Itô-Doebelin formula can now be used to identify the generator of the Ornstein-Uhlenbeck process: Taking expectation values, we obtain the forward equation

$$E[F(V_t^x)] = F(x) + \int_0^t E[(\mathcal{L}F)(V_s^x)] ds$$

for any function $F \in C_0^2(\mathbb{R})$ and $t \geq 0$, where

$$(\mathcal{L}F)(x) = \frac{1}{2}\sigma^2 f''(x) - \gamma x f'(x).$$

For the transition function this yields

$$(p_t F)(x) = F(x) + \int_0^t (p_s \mathcal{L}F)(x) \quad \text{for any } x \in \mathbb{R},$$

whence

$$\lim_{t \searrow 0} \frac{(p_t f)(x) - f(x)}{t} = \lim_{t \searrow 0} \frac{1}{t} \int_0^t E[(\mathcal{L}f)(V_s^x)] ds = (\mathcal{L}f)(x)$$

by continuity and dominated convergence. This shows that the infinitesimal generator of the Ornstein-Uhlenbeck process is an extension of the operator $(\mathcal{L}, C_0^2(\mathbb{R}))$.

Change of time-scale

We will now prove that Wiener-Itô integrals can also be represented as Brownian motion with a coordinate transformation on the time axis. Hence solutions of one-dimensional linear SDE with additive noise are affine functions of time changed Brownian motions. We first note that a Wiener-Itô integral $I_t = \int_0^t h_r dB_r$ with $h \in L_{\text{loc}}^2(0, \infty)$ is a continuous centered Gaussian process with covariance

$$\text{Cov}[I_t, I_s] = \int_0^{t \wedge s} h_r^2 dr = \tau(t) \wedge \tau(s)$$

where

$$\tau(t) := \int_0^t h_r^2 dr = \text{Var}[I_t]$$

is the corresponding variance process. The variance process should be thought of as an “internal clock” for the process (I_t) . Indeed, suppose $h > 0$ almost everywhere. Then τ is strictly increasing and continuous, and

$$\tau : [0, \infty) \rightarrow [0, \tau(\infty)) \quad \text{is a homeomorphism.}$$

Transforming the time-coordinate by τ , we have

$$\text{Cov}[I_{\tau^{-1}(t)}, I_{\tau^{-1}(s)}] = t \wedge s \quad \text{for any } t, s \in [0, \tau(\infty)].$$

These are exactly the covariance of a Brownian motion. Since a continuous Gaussian process is uniquely determined by its expectations and covariances, we can conclude:

Theorem 8.9 (Wiener-Itô integrals as time changed Brownian motions). *The process $\tilde{B}_s := I_{\tau^{-1}(s)}$, $0 \leq s < \tau(\infty)$, is a Brownian motion, and*

$$I_t = \tilde{B}_{\tau(t)} \quad \text{for any } t \geq 0, P\text{-almost surely.}$$

Proof. Since $(\tilde{B}_s)_{0 \leq s < \tau(\infty)}$ has the same marginal distributions as the Wiener-Itô integral $(I_t)_{t \geq 0}$ (but at different times), (\tilde{B}_s) is again a continuous centered Gaussian process. Moreover, $\text{Cov}[B_t, B_s] = t \wedge s$, so that (B_s) is indeed a Brownian motion. \square

Note that the argument above is different from previous considerations in the sense that the Brownian motion (\tilde{B}_s) is constructed from the process (I_t) and not vice versa.

This means that we can not represent (I_t) as a time-change of a given Brownian motion (e.g. (B_t)) but we can only show that there exists a Brownian motion (\tilde{B}_s) such that I is a time-change of \tilde{B} . This way of representing stochastic processes w.r.t. Brownian motions that are constructed from the process corresponds to the concept of weak solutions of stochastic differential equations, where driving Brownian motion is not given a

priori. We return to these ideas in Section 9, where we will also prove that continuous local martingales can be represented as time-changed Brownian motions.

Theorem 8.9 enables us to represent solution of linear SDE with additive noise by time-changed Brownian motions. We demonstrate this with an example: By the explicit formula (8.3.4) for the solution of the Ornstein-Uhlenbeck SDE, we obtain:

Corollary 8.10 (Mehler formula). *A one-dimensional Ornstein-Uhlenbeck process V_t^x with initial condition x can be represented as*

$$V_t^x = e^{-\gamma t} \left(x + \sigma \tilde{B}_{\frac{1}{2\gamma}(e^{2\gamma t} - 1)} \right)$$

with a Brownian motion $(\tilde{B}_t)_{t \geq 0}$ such that $\tilde{B}_0 = 0$.

Proof. The corresponding time change for the Wiener-Itô integral is given by

$$\tau(t) = \int_0^t \exp(2\gamma s) ds = (\exp(2\gamma t) - 1)/2\gamma.$$

□

8.4 Brownian bridge

In many circumstances one is interested in conditioning diffusion process on taking a given value at specified times. A basic example is the Brownian bridge which is Brownian motion conditioned to end at a given point x after time t_0 . We now present several ways to describe and characterize Brownian bridges. The first is based on the Wiener-Lévy construction and specific to Brownian motion, the second extends to Gaussian processes, whereas the final characterization of the bridge process as the solution of a time-homogeneous SDE can be generalized to other diffusion processes. From now on, we consider a one-dimensional Brownian motion $(B_t)_{0 \leq t \leq 1}$ with $B_0 = 0$ that we would like to condition on taking a given value y at time 1

Wiener-Lévy construction

Recall that the Brownian motion (B_t) has the Wiener-Lévy representation

$$B_t(\omega) = Y(\omega)t + \sum_{n=0}^{\infty} \sum_{k=0}^{2^n-1} 2^{n-1} Y_{n,k}(\omega) e_{n,k}(t) \quad \text{for } t \in [0, 1] \quad (8.4.1)$$

where $e_{n,k}$ are the Schauder functions, and Y and $Y_{n,k}$ ($n \geq 0, k = 0, 1, 2, \dots, 2^n - 1$) are independent and standard normally distributed. The series in (8.4.1) converges almost surely uniformly on $[0, 1]$, and the approximating partial sums are piecewise linear approximations of B_t . The random variables $Y = B_1$ and

$$X_t := \sum_{n=0}^{\infty} \sum_{k=0}^{2^n-1} Y_{n,k} e_{n,k}(t) = B_t - tB_1$$

are independent. This suggests that we can construct the bridge by replacing $Y(\omega)$ by the constant value y . Let

$$X_t^y := yt + X_t = B_t + (y - B_1) \cdot t,$$

and let μ_y denote the distribution of the process $(X_t^y)_{0 \leq t \leq 1}$ on $C([0, 1])$. The next theorem shows that X_t^y is indeed a Brownian motion conditioned to end at y at time 1:

Theorem 8.11. *The map $y \mapsto \mu_y$ is a regular version of the conditional distribution of $(B_t)_{0 \leq t \leq 1}$ given B_1 , i.e.,*

- (1). μ_y is a probability measure on $C([0, 1])$ for any $y \in \mathbb{R}$,
- (2). $P[(B_t)_{0 \leq t \leq 1} \in A \mid B_1] = \mu_{B_1}[A]$ holds P -almost surely for any given Borel subset $A \subseteq C([0, 1])$.
- (3). If $F : C([0, 1]) \rightarrow \mathbb{R}$ is a bounded and continuous function (w.r.t. the supremum norm on $C([0, 1])$) then the map $y \mapsto \int F d\mu_y$ is continuous.

The last statement says that $y \mapsto \mu_y$ is a continuous function w.r.t. the topology of weak convergence.

Proof. By definition, μ_y is a probability measure for any $y \in \mathbb{R}$. Moreover, for any Borel set $A \subseteq C([0, 1])$,

$$\begin{aligned} P[(B_t)_{0 \leq t \leq 1} \in A \mid B_1](\omega) &= P[(X_t + tB_1) \in A \mid B_1](\omega) \\ &= P[(X_t + tB_1(\omega)) \in A] = P[(X_t^{B_1(\omega)}) \in A] = \mu_{B_1(\omega)}[A] \end{aligned}$$

for P -almost every ω by independence of (X_T) and B_1 . Finally, if $F : C([0, 1]) \rightarrow \mathbb{R}$ is continuous and bounded then

$$\int F d\mu_y = E[F((y_t + X_t)_{0 \leq t \leq 1})]$$

is continuous in y by dominated convergence. \square

Finite-dimensional distributions

We now compute the marginals of the Brownian bridge X_t^y :

Corollary 8.12. *For any $n \in \mathbb{N}$ and $0 < t_1 < \dots < t_n < 1$, the distribution of $(X_{t_1}^y, \dots, X_{t_n}^y)$ on \mathbb{R}^n is absolutely continuous with density*

$$f_y(x_1, \dots, x_n) = \frac{p_{t_1}(0, x_1)p_{t_2-t_1}(x_1, x_2) \cdots p_{t_n-t_{n-1}}(x_{n-1}, x_n)p_{1-t_n}(x_n, y)}{p_1(0, y)}. \quad (8.4.2)$$

Proof. The distribution of $(B_{t_1}, \dots, B_{t_n}, B_1)$ is absolutely continuous with density

$$f_{B_{t_1}, \dots, B_{t_n}, B_1}(x_1, \dots, x_n, y) = p_{t_1}(0, x_0)p_{t_2-t_1}(x_1, x_2) \cdots p_{t_n-t_{n-1}}(x_{n-1}, x_n)p_{1-t_n}(x_n, y).$$

Since the distribution of $(X_{t_1}^y, \dots, X_{t_n}^y)$ is a regular version of the conditional distribution of $(B_{t_1}, \dots, B_{t_n})$ given B_1 , it is absolutely continuous with the conditional density

$$\begin{aligned} f_{B_{t_1}, \dots, B_{t_n} \mid B_1}(x_1, \dots, x_n \mid y) &= \frac{f_{B_{t_1}, \dots, B_{t_n}, B_1}(x_1, \dots, x_n, y)}{\int \cdots \int f_{B_{t_1}, \dots, B_{t_n}, B_1}(x_1, \dots, x_n, y) dx_1 \cdots dx_n} \\ &= f_y(x_1, \dots, x_n). \end{aligned}$$

\square

In general, any almost surely continuous process on $[0, 1]$ with marginals given by (8.4.2) is called a **Brownian bridge from 0 to y in time 1**. A Brownian bridge from x to y in time t is defined correspondingly for any $x, y \in \mathbb{R}$ and any $t > 0$. In fact, this definition of the bridge process in terms of the marginal distributions carries over from Brownian motion to arbitrary Markov processes with strictly positive transition densities. In the case of the Brownian bridge, the marginals are again normally distributed:

Theorem 8.13 (Brownian bridge as a Gaussian process). *The Brownian bridge from 0 to y in time 1 is the (in distribution unique) continuous Gaussian process $(X_t^y)_{t \in [0,1]}$ with*

$$E[X_t^y] = ty \quad \text{and} \quad \text{Cov}[X_t^y, X_s^y] = t \wedge s - ts \quad \text{for any } s, t \in [0, 1]. \quad (8.4.3)$$

Proof. A continuous Gaussian process is determined uniquely in distribution by its means and covariances. Therefore, it suffices to show that the bridge $X_t^y = B_t + (y - B_1)t$ defined above is a continuous Gaussian process such that (8.4.3) holds. This holds true: By (8.4.2), the marginals are normally distributed, and by definition, $t \mapsto X_t^y$ is almost surely continuous. Moreover,

$$\begin{aligned} E[X_t^y] &= E[B_t] + E[y - B_1] \cdot t = yt, & \text{and} \\ \text{Cov}[X_t^y, X_s^y] &= \text{Cov}[B_t, B_s] - t \cdot \text{Cov}[B_1, B_s] - s \cdot \text{Cov}[B_t, B_1] + ts \text{Var}[B_1] \\ &= t \wedge s - ts - st + ts = t \wedge s - ts. \end{aligned}$$

□

Remark (Covariance as Green function, Cameron-Martin space). The covariances of the Brownian bridge are given by

$$c(t, s) = \text{Cov}[X_t^y, X_s^y] = \begin{cases} t \cdot (1 - s) & \text{for } t \leq s, \\ (1 - t) \cdot s & \text{for } t \geq s. \end{cases}$$

The function $c(t, s)$ is the Green function of the operator d^2/dt^2 with Dirichlet boundary conditions on the interval $[0, 1]$. This is related to the fact that the distribution of the

Brownian bridge from 0 to 0 can be viewed as a standard normal distribution on the space of continuous paths $\omega : [0, 1] \rightarrow \mathbb{R}$ with $\omega(0) = \omega(1) = 0$ w.r.t. the Cameron-Martin inner product

$$(g, h)_H = \int_0^1 g'(s)h'(s) ds.$$

The second derivative d^2/dt^2 is the linear operator associated with this quadratic form.

SDE for the Brownian bridge

Our construction of the Brownian bridge by an affine transformation of Brownian motion has two disadvantages:

- It can not be carried over to more general diffusion processes with possibly non-linear drift and diffusion coefficients.
- The bridge $X_t^y = B_t + t(y - B_1)$ does not depend on (B_t) in an adapted way, because the terminal value B_1 is required to define X_t^y for any $t > 0$.

We will now show how to construct a Brownian bridge from a Brownian motion in an adapted way. The idea is to consider an SDE w.r.t. the given Brownian motion with a drift term that forces the solution to end at a given point at time 1. The size of the drift term will be large if the process is still far away from the given terminal point at a time close to 1. For simplicity we consider a bridge (X_t) from 0 to 0 in time 1. Brownian bridges with other end points can be constructed similarly. Since the Brownian bridge is a Gaussian process, we may hope that there is a linear stochastic differential equation with additive noise that has a Brownian bridge as a solution. We therefore try the Ansatz

$$dX_t = -\beta_t X_t dt + dB_t, \quad X_0 = 0 \quad (8.4.4)$$

with a given continuous deterministic function $\beta_t, 0 \leq t < 1$. By variation of constants, the solution of (8.4.4) is the Gaussian process $X_t, 0 \leq t < 1$, given by

$$X_t = \frac{1}{h_t} \int_0^t h_r dB_r \quad \text{where} \quad h_t = \exp\left(\int_0^t \beta_s ds\right).$$

The process (X_t) is centered and has covariances

$$\text{Cov}[X_t, X_s] = \frac{1}{h_t h_s} \int_0^{t \wedge s} h_r^2 dr.$$

Therefore, (X_t) is a Brownian bridge if and only if

$$\text{Cov}[X_t, X_s] = t \cdot (1 - s) \quad \text{for any } t \leq s,$$

i.e., if and only if

$$\frac{1}{t h_t} \int_0^t h_r^2 dr = h_s \cdot (1 - s) \quad \text{for any } 0 < t \leq s. \quad (8.4.5)$$

The equation (8.4.5) holds if and only if h_t is a constant multiple of $1/1 - t$, and in this case

$$\beta_t = \frac{d}{dt} \log h_t = \frac{h'_t}{h_t} = \frac{1}{1 - t} \quad \text{for } t \in [0, 1].$$

Summarizing, we have shown:

Theorem 8.14. *If (B_t) is a Brownian motion then the process (X_t) defined by*

$$X_t = \int_0^t \frac{1-t}{1-r} dB_r \quad \text{for } t \in [0, 1], \quad X_1 = 0,$$

is a Brownian bridge from 0 to 0 in time 1. It is the unique continuous process solving the SDE

$$dX_t = -\frac{X_t}{1-t} dt + dB_t \quad \text{for } t \in [0, 1). \quad (8.4.6)$$

Proof. As shown above, $(X_t)_{t \in [0,1]}$ is a continuous centered Gaussian process with the covariances of the Brownian bridge. Hence its distribution on $C([0, 1])$ coincides with that of the Brownian bridge from 0 to 0. In particular, this implies $\lim_{t \nearrow 1} X_t = 0$ almost surely, so the trivial extension from $[0, 1)$ to $[0, 1]$ defined by $X_1 = 0$ is a Brownian bridge. \square

If the Brownian bridge is replaced by a more general conditioned diffusion process, the Gaussian characterization does not apply. Nevertheless, it can still be shown by different means (the keyword is “ h -transform”) that the bridge process solves an SDE generalizing (8.4.6), cf. ?? below.

8.5 Stochastic differential equations in \mathbb{R}^n

We now explain how to generalize our considerations to systems of stochastic differential equations, or, equivalently, SDE in several dimensions. For the moment, we will not initiate a systematic study but rather consider some examples. The setup is the following: We are given a d -dimensional Brownian motion $B_t = (B_t^1, \dots, B_t^d)$. The component processes B_t^k , $1 \leq k \leq d$, are independent one-dimensional Brownian motions that drive the stochastic dynamics. We are looking for a stochastic process $X_t : \Omega \rightarrow \mathbb{R}^n$ solving an SDE of the form

$$dX_t = b(t, X_t) dt + \sum_{k=1}^d \sigma_k(t, X_t) dB_t^k. \quad (8.5.1)$$

Here n and d may be different, and $b, \sigma_1, \dots, \sigma_d : \mathbb{R}_+ \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ are time-dependent continuous vector fields on \mathbb{R}^n . In matrix notation,

$$dX_t = b(t, X_t) dt + \sigma(t, X_t) dB_t \quad (8.5.2)$$

where $\sigma(t, x) = (\sigma_1(t, x)\sigma_2(t, x) \cdots \sigma_d(t, x))$ is an $n \times d$ -matrix.

Existence, uniqueness and stability

Assuming Lipschitz continuity of the coefficients, existence, uniqueness and stability of strong solutions of the SDE (8.5.2) can be shown by similar arguments as for ordinary differential equations.

Theorem 8.15 (Existence, uniqueness and stability under global Lipschitz conditions). *Suppose that b and σ satisfy a global Lipschitz condition of the following form: For any $t_0 \in \mathbb{R}$, there exists a constant $L \in \mathbb{R}_+$ such that*

$$|b(t, x) - b(t, \tilde{x})| + \|\sigma(t, x) - \sigma(t, \tilde{x})\| \leq L \cdot |x - \tilde{x}| \quad \forall t \in [0, t_0], x, \tilde{x} \in \mathbb{R}^n. \quad (8.5.3)$$

Then for any initial value $x \in \mathbb{R}^n$, the SDE (8.5.2) has a unique (up to equivalence) strong solution $(X_t)_{t \in [0, \infty)}$ such that $X_0 = x$ P -almost surely.

Furthermore, if (X_t) and (\tilde{X}_t) are two strong solutions with arbitrary initial conditions, then for any $t \in \mathbb{R}_+$, there exists a finite constant $C(t)$ such that

$$E \left[\sup_{s \in [0, t]} |X_s - \tilde{X}_s| \right] \leq C(t) \cdot E \left[|X_0 - \tilde{X}_0|^2 \right].$$

The proof of Theorem 8.15 is outlined in the exercises below. In Section 12.1, we will prove more general results that contain the assertion of the theorem as a special case. In particular, we will see that existence up to an explosion time and uniqueness of strong solutions still hold true if one assumes only a local Lipschitz condition.

The key step for proving stability and uniqueness is to control the deviation

$$\varepsilon_t := E \left[\sup_{s \leq t} |X_s - \tilde{X}_s|^2 \right]$$

between two solutions up to time t . Existence of strong solutions can then be shown by a Picard-Lindelöf approximation based on a corresponding norm:

Exercise (Proof of stability and uniqueness). Suppose that (X_t) and (\tilde{X}_t) are strong solutions of (8.5.2), and let $t_0 \in \mathbb{R}_+$. Apply Itô's isometry and Gronwall's inequality to show that if (8.5.3) holds, then there exists a finite constant $C \in \mathbb{R}_+$ such that for any $t \leq t_0$,

$$\varepsilon_t \leq C \cdot \left(\varepsilon_0 + \int_0^t \varepsilon_s ds \right), \quad \text{and} \quad (8.5.4)$$

$$\varepsilon_t \leq C \cdot e^{Ct} \varepsilon_0. \quad (8.5.5)$$

Hence conclude that two strong solutions with the same initial value coincide almost surely.

Exercise (Existence of strong solutions). Define approximate solutions of (8.5.2) with initial value $x \in \mathbb{R}^n$ inductively by setting $X_t^0 := x$ for all t , and

$$X_t^{n+1} := x + \int_0^t b(s, X_s^n) ds + \int_0^t \sigma(s, X_s^n) dB_s.$$

Let $\Delta_t^n := E[\sup_{s \leq t} |X_s^{n+1} - X_s^n|^2]$. Show that if (8.5.3) holds, then for any $t_0 \in \mathbb{R}_+$, there exists a finite constant $C(t_0)$ such that

$$\begin{aligned} \Delta_t^{n+1} &\leq C(t_0) \int_0^t \Delta_s^n ds && \text{for any } n \geq 0 \text{ and } t \leq t_0, && \text{and} \\ \Delta_t^n &\leq C(t_0)^n \frac{t^n}{n!} \Delta_t^0 && \text{for any } n \in \mathbb{N} \text{ and } t \leq t_0. \end{aligned}$$

Hence conclude that the limit $X_s = \lim_{n \rightarrow \infty} X_s^n$ exists uniformly for $s \in [0, t_0]$ with probability one, and X is a strong solution of (8.5.2) with $X_0 = x$.

Itô processes driven by several Brownian motions

Any solution to the SDE (8.5.1) is an Itô process pf type

$$X_t = \int_0^t G_s ds + \sum_{k=1}^d \int_0^t H_s^k dB_s^k \quad (8.5.6)$$

with continuous $(\mathcal{F}_t^{B,P})$ adapted stochastic processes $G_s, H_s^1, H_s^2, \dots, H_s^d$. We now extend the stochastic calculus rules to such Itô processes that are driven by several independent Brownian motions. Let H_s and \tilde{H}_s be continuous $(\mathcal{F}_t^{B,P})$ adapted processes.

Lemma 8.16. *If (π_n) is a sequence of partitions of \mathbb{R}_+ with $\text{mesh}(\pi_n) \rightarrow 0$ then for any $1 \leq k, l \leq d$ and $a \in \mathbb{R}_+$, the covariation of the Itô integrals $t \mapsto \int_0^t H_s dB_s^k$ and $t \mapsto \int_0^t \tilde{H}_s dB_s^l$ exists almost surely uniformly for $t \in [0, a]$ along a subsequence of (π_n) , and*

$$\left[\int_0^\bullet H dB^k, \int_0^\bullet \tilde{H} dB^l \right]_t = \int_0^t H \tilde{H} d[B^k, B^l] = \delta_{kl} \int_0^t H_s \tilde{H}_s ds.$$

The proof is an extension of the proof of Theorem 8.1(ii), where the assertion has been derived for $k = l$ and $H = \tilde{H}$. The details are left as an exercise.

Similarly to the one-dimensional case, the lemma can be used to compute the covariation of Itô integrals w.r.t. arbitrary Itô processes. If X_s and Y_s are Itô processes as in (8.5.1), and K_s and L_s are adapted and continuous then we obtain

$$\left[\int_0^\bullet K dX, \int_0^\bullet L dY \right]_t = \int_0^t K_s L_s d[X, Y]_s$$

almost surely uniformly for $t \in [0, u]$, along an appropriate subsequence of (π_n) .

Multivariate Itô-Doeblin formula

We now assume again that $(X_t)_{t \geq 0}$ is a solution of a stochastic differential equation of the form (8.5.1). By Lemma 8.16, we can apply Itô's formula to almost every sample path $t \mapsto X_t(\omega)$:

Theorem 8.17 (Itô-Doeblin). *Let $F \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^n)$. Then almost surely,*

$$\begin{aligned} F(t, X_t) &= F(0, X_0) + \int_0^t (\sigma^\top \nabla_x F)(s, X_s) \cdot dB_s \\ &\quad + \int_0^t \left(\frac{\partial F}{\partial t} + \mathcal{L}F \right) (s, X_s) ds \quad \text{for all } t \geq 0, \end{aligned}$$

where ∇_x denotes the gradient in the space variable, and

$$(\mathcal{L}F)(t, x) := \frac{1}{2} \sum_{i,j=1}^n a_{i,j}(t, x) \frac{\partial^2 F}{\partial x_i \partial x_j}(t, x) + \sum_{i=1}^n b_i(t, x) \frac{\partial F}{\partial x_i}(t, x)$$

with $a(t, x) := \sigma(t, x)\sigma(t, x)^\top \in \mathbb{R}^{n \times n}$.

Proof. If X is a solution to the SDE then

$$\begin{aligned} [X^i, X^j]_t &= \sum_{k,l} \left[\int \sigma_k^i(s, X) dB^k, \int \sigma_l^j(s, X) dB^l \right]_t \\ &= \sum_{k,l} \int_0^t (\sigma_k^i \sigma_l^j)(s, X) d[B^k, B^l] = \int_0^t a^{ij}(s, X_s) ds \end{aligned}$$

where $a^{ij} = \sum_k \sigma_k^i \sigma_k^j$, i.e.,

$$a(s, x) = \sigma(s, x) \sigma(s, x)^T \in \mathbb{R}^{n \times n}.$$

Therefore, Itô's formula applied to the process (t, X_t) yields

$$\begin{aligned} dF(t, X) &= \frac{\partial F}{\partial t}(t, X) dt + \nabla_x F(t, X) \cdot dX + \frac{1}{2} \sum_{i,j=1}^d \frac{\partial^2 F}{\partial x^i \partial x^j}(t, X) d[X^i, X^j] \\ &= (\sigma^T \nabla_x F)(t, X) \cdot dB + \left(\frac{\partial F}{\partial t} + \mathcal{L}F \right)(t, X) dt, \end{aligned}$$

for any $F \in C^{1,2}(\mathbb{R}_+ \times \mathbb{R}^n)$. □

The Itô-Doeblin formula shows that for any $F \in C^2(\mathbb{R}_+ \times \mathbb{R}^n)$, the process

$$M_s^F = F(s, X_s) - F(0, X_0) - \int_0^s \left(\frac{\partial F}{\partial t} + \mathcal{L}F \right)(t, X_t) dt$$

is a local martingale. If $\sigma^T \nabla_x F$ is bounded then M^F is a global martingale.

Exercise (Drift and diffusion coefficients). Show that the processes

$$M_s^i = X_s^i - X_0^i - \int_0^s b^i(s, X_s) ds, \quad 1 \leq i \leq n,$$

are local martingales with covariations

$$[M^i, M^j]_s = a_{i,j}(s, X_s) \quad \text{for any } s \geq 0, P\text{-almost surely.}$$

The vector field $b(s, x)$ is called the *drift vector field* of the SDE, and the coefficients $a_{i,j}(s, x)$ are called *diffusion coefficients*.

General Ornstein-Uhlenbeck processes

XXX to be included

Example (Stochastic oscillator).

Examples

Example (Physical Brownian motion with external force).

Example (Kalman-Bucy filter).

Example (Heston model for stochastic volatility).

Chapter 9

Change of measure

9.1 Local and global densities of probability measures

A thorough understanding of absolute continuity and relative densities of probability measures is crucial at many places in stochastic analysis. Martingale convergence yields an elegant approach to these issues including a proof of the Radon-Nikodym and the Lebesgue Decomposition Theorem. We first recall the definition of absolute continuity.

Absolute Continuity

Suppose that P and Q are probability measures on a measurable space (Ω, \mathcal{A}) , and \mathcal{F} is a sub- σ -algebra of \mathcal{A} .

Definition. (1). The measure Q is called **absolutely continuous w.r.t. P on the σ -algebra \mathcal{F}** if and only if $Q[A] = 0$ for any $A \in \mathcal{F}$ with $P[A] = 0$.

(2). The measures Q and P are called **singular on \mathcal{F}** if and only if there exists $A \in \mathcal{F}$ such that $P[A] = 0$ and $Q[A^C] = 0$.

We use the notations $Q \ll P$ for absolute continuity of Q w.r.t. P , $Q \approx P$ for mutual absolute continuity, and $Q \perp P$ for singularity of Q and P . The definitions above extend to signed measures.

Example. The Dirac measure $\delta_{1/2}$ is obviously singular w.r.t. Lebesgue measure $\lambda_{(0,1]}$ on the Borel σ -algebra $\mathcal{B}((0, 1])$. However, $\delta_{1/2}$ is absolutely continuous w.r.t. $\lambda_{(0,1]}$ on each of the σ -algebras $\mathcal{F}_n = \sigma(\mathcal{D}_n)$ generated by the dyadic partitions $\mathcal{D}_n = \{(k \cdot 2^{-n}, (k + 1)2^{-n}] : 0 \leq k < 2^n\}$, and $\mathcal{B}([0, 1)) = \sigma(\bigcup \mathcal{D}_n)$.

The next lemma clarifies the term “absolute continuity.”

Lemma 9.1. *The probability measure Q is absolutely continuous w.r.t. P on the σ -algebra \mathcal{F} if and only if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for $A \in \mathcal{F}$,*

$$P[A] < \delta \quad \Rightarrow \quad Q[A] < \varepsilon. \tag{9.1.1}$$

Proof. The “if” part is obvious. If $P[A] = 0$ and (9.1.1) holds for each $\varepsilon > 0$ with δ depending on ε then $Q[A] < \varepsilon$ for any $\varepsilon > 0$, and hence $Q[A] = 0$.

To prove the “only if” part, we suppose that there exists $\varepsilon > 0$ such that (9.1.1) does not hold for any $\delta > 0$. Then there exists a sequence (A_n) of events in \mathcal{F} such that

$$Q[A_n] \geq \varepsilon \quad \text{and} \quad P[A_n] \leq 2^{-n}.$$

Hence, by the Borel-Cantelli-Lemma,

$$P[A_n \text{ infinitely often}] = 0,$$

whereas

$$Q[A_n \text{ infinitely often}] = Q \left[\bigcap_{n} \bigcup_{m \geq n} A_m \right] = \lim_{n \rightarrow \infty} Q \left[\bigcup_{m \geq n} A_m \right] \geq \varepsilon.$$

Therefore Q is not absolutely continuous w.r.t. P . □

Example (Absolute continuity on \mathbb{R}). A probability measure μ on a real interval is absolutely continuous w.r.t. Lebesgue measure if and only if the distribution function $F(t) = \mu[(-\infty, t]]$ satisfies:

For any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $n \in \mathbb{N}$ and $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}$,

$$\sum_{i=1}^n |b_i - a_i| < \varepsilon \quad \Rightarrow \quad \sum_{i=1}^n |F(b_i) - F(a_i)| < \delta, \quad (9.1.2)$$

cf. e.g. [Billingsley: Probability and Measures].

Definition (Absolutely continuous functions). A function $F : (a, b) \subset \mathbb{R} \rightarrow \mathbb{R}$ is called *absolutely continuous* iff (9.1.2) holds.

The Radon-Nikodym Theorem states that absolute continuity is equivalent to the existence of a relative density.

Theorem 9.2 (Radon-Nikodym). The probability measure Q is absolutely continuous w.r.t. P on the σ -algebra \mathcal{F} if and only if there exists a non-negative random variable $Z \in \mathcal{L}^1(\Omega, \mathcal{F}, P)$ such that

$$Q[A] = \int_A Z dP \quad \text{for any } A \in \mathcal{F}. \quad (9.1.3)$$

The relative density Z of Q w.r.t. P on \mathcal{F} is determined by (9.1.3) uniquely up to modification on P -measure zero sets. It is also called the **Radon-Nikodym derivative** or the **likelihood ratio** of Q w.r.t. P on \mathcal{F} . We use the notation

$$Z = \left. \frac{dQ}{dP} \right|_{\mathcal{F}},$$

and omit the \mathcal{F} when the choice of the σ -algebra is clear.

Example (Finitely generated σ -algebra). Suppose that the σ -algebra \mathcal{F} is generated by finitely many disjoint atoms B_1, \dots, B_k with $\Omega = \bigcup B_i$. Then Q is absolutely continuous w.r.t. P if and only if for any i ,

$$P[B_i] = 0 \quad \Rightarrow \quad Q[B_i] = 0.$$

In this case, the relative density is given by

$$\frac{dQ}{dP}\Big|_{\mathcal{F}} = \sum_{i: P[B_i] > 0} \frac{Q[B_i]}{P[B_i]} \cdot I_{B_i}.$$

From local to global densities

Let (\mathcal{F}_n) be a given filtration on (Ω, \mathcal{A}) .

Definition (Local absolutely continuity). *The measure Q is called **locally absolutely continuous** w.r.t. P and the filtration (\mathcal{F}_n) if and only if Q is absolutely continuous w.r.t. P on the σ -algebra \mathcal{F}_n for each n .*

Example (Dyadic partitions). Any probability measure on the unit interval $[0, 1]$ is locally absolutely continuous w.r.t. Lebesgue measure on the filtration $\mathcal{F}_n = \sigma(\mathcal{D}_n)$ generated by the dyadic partitions of the unit interval. The Radon-Nikodym derivative on \mathcal{F}_n is the dyadic difference quotient defined by

$$\frac{d\mu}{d\lambda}\Big|_{\mathcal{F}_n}(x) = \frac{\mu[(k-1) \cdot 2^{-n}, k \cdot 2^{-n}]}{\lambda[(k-1) \cdot 2^{-n}, k \cdot 2^{-n}]} = \frac{F(k \cdot 2^{-n}) - F((k-1) \cdot 2^{-n})}{2^{-n}} \quad (9.1.4)$$

for $x \in ((k-1)2^{-n}, k2^{-n}]$.

Example (Product measures). If $Q = \bigotimes_{i=1}^{\infty} \nu$ and $P = \bigotimes_{i=1}^{\infty} \mu$ are infinite products of probability measures ν and μ , and ν is absolutely continuous w.r.t. μ with density ϱ , then Q is locally absolutely continuous w.r.t. P on the filtration

$$\mathcal{F}_n = \sigma(X_1, \dots, X_n)$$

generated by the coordinate maps $X_i(\omega) = \omega_i$. The local relative density is

$$\frac{dQ}{dP}\Big|_{\mathcal{F}_n} = \prod_{i=1}^n \varrho(X_i)$$

However, if $\nu \neq \mu$, then Q is not absolutely continuous w.r.t. P on $\mathcal{F}_{\infty} = \sigma(X_1, X_2, \dots)$, since by the LLN, $n^{-1} \sum_{i=1}^n I_A(X_i)$ converges Q almost surely to $\nu[A]$ and P -almost surely to $\mu[A]$.

Now suppose that Q is locally absolutely continuous w.r.t. P on a filtration (\mathcal{F}_n) with relative densities

$$Z_n = \frac{dQ}{dP} \Big|_{\mathcal{F}_n}.$$

The L^1 martingale convergence theorem can be applied to study the existence of a global density on the σ -algebra

$$\mathcal{F}_\infty = \sigma\left(\bigcup \mathcal{F}_n\right).$$

Let $Z_\infty := \limsup Z_n$.

Theorem 9.3 (Convergence of local densities, Lebesgue decomposition).

(1). *The sequence (Z_n) of successive relative densities is an (\mathcal{F}_n) -martingale w.r.t. P . In particular, (Z_n) converges P -almost surely to Z_∞ , and Z_∞ is integrable w.r.t. P .*

(2). *The following statements are equivalent:*

(a) *(Z_n) is uniformly integrable w.r.t. P .*

(b) *Q is absolutely continuous w.r.t. P on \mathcal{F}_∞ .*

(c) *$Q[A] = \int_A Z_\infty dP$ for any P on \mathcal{F}_∞ .*

(3). *In general, the decomposition $Q = Q_a + Q_s$ holds with*

$$Q_a[A] = \int_A Z_\infty dP, \quad Q_s[A] = Q[A \cap \{Z_\infty = \infty\}]. \quad (9.1.5)$$

Q_a and Q_s are positive measure with $Q_a \ll P$ and $Q_s \perp P$.

The decomposition $Q = Q_a + Q_s$ into an absolutely continuous and a singular part is called the **Lebesgue decomposition** of the measure Q w.r.t. P on the σ -algebra \mathcal{F}_∞ .

Proof. (1). For $n \geq 0$, the density Z_n is in $\mathcal{L}^1(\Omega, \mathcal{F}_n, P)$, and

$$E_P[Z_n; A] = Q[A] = E_P[Z_{n+1}; A] \quad \text{for any } A \in \mathcal{F}_n.$$

Hence $Z_n = E_P[Z_{n+1} | \mathcal{F}_n]$, i.e., (Z_n) is a martingale w.r.t. P . Since $Z_n \geq 0$, the martingale converges P -almost surely, and the limit is integrable.

(2). (a) \Rightarrow (c): If (Z_n) is uniformly integrable w.r.t. P , then

$$Z_n = E_P[Z_\infty | \mathcal{F}_n] \quad P\text{-almost surely for any } n,$$

by the L^1 convergence theorem. Hence for $A \in \mathcal{F}_n$,

$$Q[A] = E_P[Z_n; A] = E_P[Z_\infty; A].$$

This shows that $Q[A] = E_P[Z_\infty; A]$ holds for any $A \in \bigcup \mathcal{F}_n$, and thus for any $A \in \mathcal{F}_\infty = \sigma(\bigcup \mathcal{F}_n)$.

(c) \Rightarrow (b) is evident.

(b) \Rightarrow (a): If $Q \ll P$ on \mathcal{F}_∞ then Z_n converges also Q -almost surely to a finite limit Z_∞ . Hence for $n_0 \in \mathbb{N}$ and $c > 1$,

$$\begin{aligned} \sup_n E_P[|Z_n|; |Z_n| \geq c] &= \sup_n E_P[Z_n; Z_n \geq c] = \sup_n Q[Z_n \geq c] \\ &\leq \max_{n < n_0} Q[Z_n \geq c] + \sup_{n \geq n_0} Q[Z_n \geq c] \\ &\leq \max_{n < n_0} Q[Z_n \geq c] + Q[Z_\infty \geq c - 1] + \sup_{n \geq n_0} Q[|Z_n - Z_\infty| \geq 1]. \end{aligned}$$

Given $\varepsilon > 0$, the last summand is smaller than $\varepsilon/3$ for n_0 sufficiently large, and the other two summands on the right hand side are smaller than $\varepsilon/3$ if c is chosen sufficiently large depending on n_0 . Hence (Z_n) is uniformly integrable w.r.t. P .

(3). In general, $Q_a[A] = E_P[Z_\infty; A]$ is a positive measure on \mathcal{F}_∞ with $Q_a \leq Q$, since for $n \geq 0$ and $A \in \mathcal{F}_n$,

$$Q_a[A] = E_P[\liminf_{k \rightarrow \infty} Z_k; A] \leq \liminf_{k \rightarrow \infty} E_P[Z_k; A] = E_P[Z_n; A] = Q[A]$$

by Fatou's Lemma and the martingale property. It remains to show that

$$Q_a[A] = Q[A \cap \{Z_\infty < \infty\}] \quad \text{for any } A \in \mathcal{F}_\infty. \quad (9.1.6)$$

If (9.1.6) holds, then $Q = Q_a + Q_s$ with Q_s defined by (9.1.5). In particular, Q_s is then singular w.r.t. P , since $P[Z_\infty = \infty] = 0$ and $Q_s[Z_\infty = \infty] = 0$, whereas

Q_a is absolutely continuous w.r.t. P by definition.

Since $Q_a \leq Q$, it suffices to verify (9.1.6) for $A = \Omega$. Then

$$(Q - Q_a)[A \cap \{Z_\infty < \infty\}] = (Q - Q_a)[Z_\infty < \infty] = 0,$$

and therefore

$$Q[A \cap \{Z_\infty < \infty\}] = Q_a[A \cap \{Z_\infty < \infty\}] = Q_a[A]$$

for any $A \in \mathcal{F}_\infty$.

To prove (9.1.6) for $A = \Omega$ we observe that for $c \in (0, \infty)$,

$$\begin{aligned} Q \left[\limsup_{n \rightarrow \infty} Z_n < c \right] &\leq \limsup_{n \rightarrow \infty} Q[Z_n < c] = \limsup_{n \rightarrow \infty} E_P[Z_n; Z_n < c] \\ &\leq E_P \left[\limsup_{n \rightarrow \infty} Z_n \cdot I_{\{Z_n < c\}} \right] \leq E_P[Z_\infty] = Q_a[\Omega] \end{aligned}$$

by Fatou's Lemma. As $c \rightarrow \infty$, we obtain

$$Q[Z_\infty < \infty] \leq Q_a[\Omega] = Q_a[Z_\infty < \infty] \leq Q[Z_\infty < \infty]$$

and hence (9.1.6) with $A = \Omega$. This completes the proof □

As a first consequence of Theorem 9.3, we prove the Radon-Nikodym Theorem on a separable σ -algebra \mathcal{A} . Let P and Q be probability measures on (Ω, \mathcal{A}) with $Q \ll P$.

Proof of the Radon-Nikodym Theorem for separable σ -algebras. We fix a filtration (\mathcal{F}_n) consisting of finitely generated σ -algebras $\mathcal{F}_n \subseteq \mathcal{A}$ with $\mathcal{A} = \sigma(\bigcup \mathcal{F}_n)$. Since Q is absolutely continuous w.r.t. P , the local densities $Z_n = dQ/dP|_{\mathcal{F}_n}$ on the finitely generated σ -algebras \mathcal{F}_n exist, cf. the example above. Hence by Theorem 9.3,

$$Q[A] = \int_A Z_\infty dP \quad \text{for any } A \in \mathcal{A}.$$

□

The approach above can be generalized to probability measures that are not absolutely continuous:

Exercise (Lebesgue decomposition, Lebesgue densities). Let P and Q be arbitrary (not necessarily absolutely continuous) probability measures on (Ω, \mathcal{A}) . A **Lebesgue density** of Q w.r.t. P is a random variable $Z : \Omega \rightarrow [0, \infty]$ such that $Q = Q_a + Q_s$ with

$$Q_a[A] = \int_A Z dP, \quad Q_s[A] = Q[A \cap \{Z = \infty\}] \quad \text{for any } A \in \mathcal{A}.$$

The goal of the exercise is to prove that a Lebesgue density exists if the σ -algebra \mathcal{A} is separable.

- (1). Show that if Z is a Lebesgue density of Q w.r.t. P then $1/Z$ is a Lebesgue density of P w.r.t. Q . Here $1/\infty := 0$ and $1/0 := \infty$.

From now on suppose that the σ -algebra is separable with $\mathcal{A} = \sigma(\bigcup \mathcal{F}_n)$ where (\mathcal{F}_n) is a filtration consisting of σ -algebras generated by finitely many atoms.

- (1). Write down Lebesgue densities Z_n of Q w.r.t. P on each \mathcal{F}_n . Show that

$$Q[Z_n = \infty \text{ and } Z_{n+1} < \infty] = 0 \quad \text{for any } n,$$

and conclude that (Z_n) is a non-negative supermartingale under P , and $(1/Z_n)$ is a non-negative supermartingale under Q .

- (2). Prove that the limit $Z_\infty = \lim Z_n$ exists both P -almost surely and Q -almost surely, and $P[Z_\infty < \infty] = 1$ and $Q[Z_\infty > 0] = 1$.
- (3). Conclude that Z_∞ is a Lebesgue density of P w.r.t. Q on \mathcal{A} , and $1/Z_\infty$ is a Lebesgue density of Q w.r.t. P on \mathcal{A} .

Derivatives of monotone functions

Suppose that $F : [0, 1] \rightarrow \mathbb{R}$ is a monotone and right-continuous function. After an appropriate linear transformation we may assume that F is non decreasing with $F(0) = 0$ and $F(1) = 1$. Let μ denote the probability measure with distribution function F . By the example above, the Radon-Nikodym derivative of μ w.r.t. Lebesgue measure on the σ -algebra $\mathcal{F}_n = \sigma(\mathcal{D}_n)$ generated by the n -th dyadic partition of the unit interval is given by the dyadic difference quotients (9.1.4) of F . By Theorem 9.3, we obtain a version of Lebesgue's Theorem on derivatives of monotone functions:

Corollary 9.4 (Lebesgue's Theorem). *Suppose that $F : [0, 1] \rightarrow \mathbb{R}$ is monotone (and right continuous). Then the dyadic derivative*

$$F'(t) = \lim_{n \rightarrow \infty} \left. \frac{d\mu}{d\lambda} \right|_{\mathcal{F}_n} (t)$$

exists for almost every t and F' is an integrable function on $(0, 1)$. Furthermore, if F is absolutely continuous then

$$F(s) = \int_0^s F'(t) dt \quad \text{for all } s \in [0, 1]. \quad (9.1.7)$$

Remark. Right continuity is only a normalization and can be dropped from the assumptions. Moreover, the assertion extends to function of finite variation since these can be represented as the difference of two monotone functions, cf. ?? below. Similarly, (9.1.7) also holds for absolutely continuous functions that are not monotone. See e.g. [Elstrodt: Maß- und Integrationstheorie] for details.

Absolute continuity of infinite product measures

Suppose that $\Omega = \prod_{i=1}^{\infty} S_i$, and

$$Q = \bigotimes_{i=1}^{\infty} \nu_i \quad \text{and} \quad P = \bigotimes_{i=1}^{\infty} \mu_i$$

are products of probability measures ν_i and μ_i defined on measurable spaces (S_i, \mathcal{S}_i) . We assume that ν_i and μ_i are mutually absolutely continuous for every $i \in \mathbb{N}$. Denoting by $X_k : \Omega \rightarrow S_k$ the evaluation of the k -th coordinate, the product measures are mutually absolutely continuous on each of the σ -algebras

$$\mathcal{F}_n = \sigma(X_1, \dots, X_n), \quad n \in \mathbb{N},$$

with relative densities

$$\left. \frac{dQ}{dP} \right|_{\mathcal{F}_n} = Z_n \quad \text{and} \quad \left. \frac{dP}{dQ} \right|_{\mathcal{F}_n} = 1/Z_n,$$

where

$$Z_n = \prod_{i=1}^n \frac{d\nu_i}{d\mu_i}(X_i) \in (0, \infty) \quad P\text{-almost surely.}$$

In particular, (Z_n) is a martingale under P , and $(1/Z_n)$ is a martingale under Q . Let $\mathcal{F}_\infty = \sigma(X_1, X_2, \dots)$ denote the product σ -algebra.

Theorem 9.5 (Kakutani’s dichotomy). *The infinite product measures Q and P are either singular or mutually absolutely continuous with relative density Z_∞ . More precisely, the following statements are equivalent:*

- (1). $Q \ll P$ on \mathcal{F}_∞ .
- (2). $Q \approx P$ on \mathcal{F}_∞ .
- (3). $\prod_{i=1}^\infty \int \sqrt{\frac{d\nu_i}{d\mu_i}} d\mu_i > 0$.
- (4). $\sum_{i=1}^\infty d_H^2(\nu_i, \mu_i) < \infty$.

Here the squared Hellinger distance $d_H^2(\nu_i, \mu_i)$ of mutually absolutely continuous probability measures ν and μ is defined by

$$\begin{aligned} d_H^2 &= \frac{1}{2} \int \left(\sqrt{\frac{d\nu}{d\mu}} - 1 \right)^2 d\mu = \frac{1}{2} \int \left(\sqrt{\frac{d\mu}{d\nu}} - 1 \right)^2 d\nu \\ &= 1 - \int \sqrt{\frac{d\nu}{d\mu}} d\mu = 1 - \int \sqrt{\frac{d\mu}{d\nu}} d\nu. \end{aligned}$$

Remark. (1). If mutual absolutely continuity holds then the relative densities on \mathcal{F}_∞ are

$$\frac{dQ}{dP} = \lim_{n \rightarrow \infty} Z_n \quad P\text{-almost surely, and} \quad \frac{dP}{dQ} = \lim_{n \rightarrow \infty} \frac{1}{Z_n} \quad Q\text{-almost surely.}$$

(2). If ν and μ are absolutely continuous w.r.t. a measure dx then

$$d_H^2(\nu, \mu) = \frac{1}{2} \int \left(\sqrt{f(x)} - \sqrt{g(x)} \right)^2 dx = 1 - \int \sqrt{f(x)g(x)} dx.$$

Proof. (1) \iff (3): For $i \in \mathbb{N}$ let $Y_i := \frac{d\nu_i}{d\mu_i}(X_i)$. Then the random variables Y_i are independent under both P and Q with $E_P[Y_i] = 1$, and

$$Z_n = Y_1 \cdot Y_2 \cdots Y_n.$$

By Theorem 9.3, the measure Q is absolutely continuous w.r.t. P if and only if the martingale (Z_n) is uniformly integrable. To obtain a sharp criterion for uniform integrability we switch from L^1 to L^2 , and consider the non-negative martingale

$$M_n = \frac{\sqrt{Y_1}}{\beta_1} \cdot \frac{\sqrt{Y_2}}{\beta_2} \cdots \frac{\sqrt{Y_n}}{\beta_n} \quad \text{with } \beta_i = E_P[\sqrt{Y_i}] = \int \sqrt{\frac{d\nu_i}{d\mu_i}} d\mu_i$$

under the probability measure P . Note that for $n \in \mathbb{N}$,

$$E[M_n^2] = \prod_{i=1}^n E[Y_i]/\beta_i^2 = 1 / \left(\prod_{i=1}^n \beta_i \right)^2.$$

If (3) holds then (M_n) is bounded in $L^2(\Omega, \mathcal{A}, P)$. Therefore, by Doob's L^2 inequality, the supremum of M_n is in $\mathcal{L}^2(\Omega, \mathcal{A}, P)$, i.e.,

$$E[\sup |Z_n|] = E[\sup M_n^2] < \infty.$$

Thus (Z_n) is uniformly integrable and $Q \ll P$ on \mathcal{F}_∞ .

Conversely, if (3) does not hold then

$$Z_n = M_n^2 \cdot \prod_{i=1}^n \beta_i \longrightarrow 0 \quad P\text{-almost surely,}$$

since M_n converges to a finite limit by the martingale convergence theorem. Therefore, the absolute continuous part Q_a vanishes by Theorem 9.3 (3), i.e., Q is singular w.r.t. P .

(3) \iff (4): For reals $\beta_i \in (0, 1)$, the condition $\prod_{i=1}^{\infty} \beta_i > 0$ is equivalent to $\sum_{i=1}^{\infty} (1 - \beta_i) < \infty$. For β_i as above, we have

$$1 - \beta_i = 1 - \int \sqrt{\frac{d\nu_i}{d\mu_i}} d\mu_i = d_H^2(\nu_i, \mu_i).$$

(2) \Rightarrow (1) is obvious.

(4) \Rightarrow (2): Condition (4) is symmetric in ν_i and μ_i . Hence, if (4) holds then both $Q \ll P$ and $P \ll Q$. \square

Example (Gaussian products). Let $P = \bigotimes_{i=1}^{\infty} N(0, 1)$ and $Q = \bigotimes_{i=1}^{\infty} N(a_i, 1)$ where $(a_i)_{i \in \mathbb{N}}$ is a sequence of reals. The relative density of the normal distributions $\nu_i := N(a_i, 1)$ and $\mu := N(0, 1)$ is

$$\frac{d\nu_i}{d\mu}(x) = \frac{\exp(-(x - a_i)^2/2)}{\exp(-x^2/2)} = \exp(a_i x - a_i^2/2),$$

and

$$\int \sqrt{\frac{d\nu_i}{d\mu}} d\mu = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left(-\frac{1}{2}(x^2 - a_i x + a_i^2/2)\right) dx = \exp(-a_i^2/8).$$

Therefore, by condition (3) in Theorem 9.5,

$$Q \ll P \iff Q \approx P \iff \sum_{i=1}^{\infty} a_i^2 < \infty.$$

Hence mutual absolute continuity holds for the infinite products if and only if the sequence (a_i) is contained in ℓ^2 , and otherwise Q and P are singular.

Remark (Relative entropy). (1). In the singular case, the exponential rate of degeneration of the relative densities on the σ -algebras \mathcal{F}_n is related to the relative entropies

$$H(\nu_i | \mu_i) = \int \frac{d\nu_i}{d\mu_i} \log \frac{d\nu_i}{d\mu_i} d\mu_i = \int \log \frac{d\nu_i}{d\mu_i} d\nu_i.$$

For example in the i.i.d. case $\mu_i \equiv \mu$ and $\nu_i \equiv \nu$, we have

$$\frac{1}{n} \log Z_n = \frac{1}{n} \sum_{i=1}^n \log \frac{d\nu}{d\mu}(X_i) \longrightarrow H(\nu | \mu) \quad Q\text{-a.s., and}$$

$$-\frac{1}{n} \log Z_n = \frac{1}{n} \log Z^{-1} \longrightarrow H(\mu | \nu) \quad P\text{-a.s.}$$

as $n \rightarrow \infty$ by the Law of Large Numbers.

In general, $\log Z_n - \sum_{i=1}^n H(\nu_i | \mu_i)$ is a martingale w.r.t. Q , and $\log Z_n + \sum_{i=1}^n H(\nu_i | \mu_i)$ is a martingale w.r.t. P .

(2). The relative entropy is related to the squared Hellinger distance by the inequality

$$\frac{1}{2} H(\nu | \mu) \geq d_H^2(\nu | \mu),$$

which follows from the elementary inequality

$$\frac{1}{2} \log x^{-1} = -\log \sqrt{x} \geq 1 - \sqrt{x} \quad \text{for } x > 0.$$

9.2 Translations of Wiener measure

We now return to stochastic processes in continuous time. We endow the continuous path space $C([0, \infty), \mathbb{R}^d)$ with the σ -algebra generated by the evolution maps $X_t(\omega) = \omega(t)$, and with the filtration


$$\mathcal{F}_t^X = \sigma(X_s \mid s \in [0, t]), \quad t \geq 0.$$

Note that \mathcal{F}_t^X consists of all sets of type

$$\{\omega \in C([0, \infty), \mathbb{R}^d) : \omega|_{[0, t]} \in \Gamma\} \quad \text{with } \Gamma \in \mathcal{B}(C([0, t], \mathbb{R}^d)).$$

In many situations one is interested in the distribution on path space of a process

$$B_t^h = B_t + h(t)$$



obtained by translating a Brownian motion (B_t) by a deterministic function $h : [0, \infty) \rightarrow \mathbb{R}^d$. In particular, it is important to know if the distribution of (B_t^h) has a density w.r.t. the Wiener measure on the σ -algebras \mathcal{F}_t^X , and how to compute the densities if they exist.

Example. (1). Suppose we would like to evaluate the probability that $\sup_{s \in [0, t]} |B_s - g(s)| < \varepsilon$ for a given $t > 0$ and a given function $g \in C([0, \infty), \mathbb{R}^d)$ asymptotically as $\varepsilon \searrow 0$. One approach is to study the distribution of the translated process $B_t - g(t)$ near 0.

- (2). Similarly, computing the passage probability $P[B_s \geq a+bs \text{ for some } s \in [0, t]]$ to a line $s \mapsto a + bs$ for a one-dimensional Brownian motion is equivalent to computing the passage probability to the point a for the translated process $B_t - bt$.
- (3). A solution to a stochastic differential equation

$$dY_t = dB_t + b(t, Y_t)dt$$

is a translation of the Brownian motion $B_t - B_0$ by the stochastic process $H_t = Y_0 + \int_0^t b(s, Y_s) ds$. Again, in the simplest case (when $b(t, y)$ only depends on t), H_t is a deterministic function.

The Cameron-Martin Theorem

Let (B_t) denote a continuous Brownian motion with $B_0 = 0$, and let $h \in C([0, \infty), \mathbb{R}^d)$. The distribution

$$\mu_h := P \circ (B + h)^{-1}$$

of the translated process $B_t^h = B_t + h(t)$ is the image of Wiener measure μ_0 under the translation map

$$\tau_h : C([0, \infty), \mathbb{R}^d) \longrightarrow C([0, \infty), \mathbb{R}^d), \quad \tau_h(x) = x + h.$$

Recall that Wiener measure is a Gaussian measure on the infinite dimensional space $C([0, \infty), \mathbb{R}^d)$. The next exercise discusses translations of Gaussian measures in \mathbb{R}^n :

Exercise (Translations of normal distributions). Let $C \in \mathbb{R}^{n \times n}$ be a symmetric non-negative definite matrix, and let $h \in \mathbb{R}^n$. the image of the normal distribution $N(0, C)$ under the translation map $x \mapsto x + h$ on \mathbb{R}^n is the normal distribution $N(h, C)$.

- (1). Show that if C is non-degenerate then $N(h, C) \approx N(0, C)$ with relative density

$$\frac{dN(h, C)}{dN(0, C)}(x) = e^{(h,x) - \frac{1}{2}(h,h)} \quad \text{for } x \in \mathbb{R}^n, \quad (9.2.1)$$

where $(g, h) = (g, C^{-1}h)$ for $g, h \in \mathbb{R}^n$.

- (2). Prove that in general, $N(h, C)$ is absolutely continuous w.r.t. $N(0, C)$ if and only if h is orthogonal to the kernel of C w.r.t. the Euclidean inner product.

On $C([0, \infty), \mathbb{R}^d)$, we can usually not expect the existence of a global density of the translated measures μ_h w.r.t. μ_0 . The Cameron-Martin Theorem states that for $t \geq 0$, a relative density on \mathcal{F}_t^X exists if and only if h is contained in the corresponding Cameron-Martin space:

Theorem 9.6 (Cameron, Martin). For $h \in C([0, \infty), \mathbb{R}^d)$ and $t \in \mathbb{R}_+$ the translated measure $\mu_h = \mu \circ \tau_h^{-1}$ is absolutely continuous w.r.t. Wiener measure μ_0 on \mathcal{F}_t^X if and only if h is an absolutely continuous function on $[0, t]$ with $h(0) = 0$ and $\int_0^t |h'(s)|^2 ds < \infty$. In this case, the relative density is given by

$$\left. \frac{d\mu_h}{d\mu_0} \right|_{\mathcal{F}_t^X} = \exp \left(\int_0^t h'(s) dX_s - \frac{1}{2} \int_0^t |h'(s)|^2 ds \right). \quad (9.2.2)$$

where $\int_0^t h'(s) dX_s$ is the Itô integral w.r.t. the canonical Brownian motion (X, μ_0) .

Before giving a rigorous proof let us explain heuristically why the result should be true. Clearly, absolute continuity does not hold if $h(0) \neq 0$, since then the translated paths do not start at 0. Now suppose $h(0) = 0$, and fix $t \in (0, \infty)$. Absolute continuity on \mathcal{F}_t^X means that the distribution μ_h^t of $(B_s^h)_{0 \leq s \leq t}$ on $C([0, t], \mathbb{R}^d)$ is absolutely continuous w.r.t. Wiener measure μ_0^t on this space. The measure μ_0^t , however, is a kind of infinite dimensional standard normal distribution w.r.t. the inner product

$$(x, y)_H = \int_0^t x'(s) \cdot y'(s) ds$$

on functions $x, y : [0, t] \rightarrow \mathbb{R}^d$ vanishing at 0, and the translated measure μ_h^t is a Gaussian measure with mean h and the same covariances.

Choosing an orthonormal basis $(e_i)_{i \in \mathbb{N}}$ w.r.t. the H -inner product (e.g. Schauder functions), we can identify μ_0^t and μ_h^t with the product measures $\bigotimes_{i=1}^{\infty} N(0, 1)$ and $\bigotimes_{i=1}^{\infty} N(\alpha_i, 1)$

respectively where $\alpha_i = (h, e_i)_H$ is the i -th coefficient of h in the basis expansion. Therefore, μ_h^t should be absolutely continuous w.r.t. μ_0^t if and only if

$$(h, h)_H = \sum_{i=1}^{\infty} \alpha_i^2 < \infty,$$

i.e., if and only if h is absolutely continuous with $h' \in \mathcal{L}^2(0, t)$.

Moreover, in analogy to the finite-dimensional case (9.2.1), we would expect informally a relative density of the form

$$\text{“} \frac{d\mu_h^t}{d\mu_0^t}(x) = e^{(h,x)_H - \frac{1}{2}(h,h)_H} = \exp\left(\int_0^t h'(s) \cdot x'(s) ds - \frac{1}{2} \int_0^t |h'(s)|^2 ds\right) \text{”}$$

Since μ_0^t -almost every path $x \in C([0, \infty), \mathbb{R}^d)$ is not absolutely continuous, this expression does not make sense. Nevertheless, using finite dimensional approximations, we can derive the rigorous expression (9.2.2) for the relative density where the integral $\int_0^t h'x' ds$ is replaced by the almost surely well-defined stochastic integral $\int_0^t h' dx$:

Proof of Theorem 9.6. We assume $t = 1$. The proof for other values of t is similar. Moreover, as explained above, it is enough to consider the case $h(0) = 0$.

- (1). *Local densities:* We first compute the relative densities when the paths are only evaluated at dyadic time points. Fix $n \in \mathbb{N}$, let $t_i = i \cdot 2^{-n}$, and let

$$\delta_i x = x_{t_{i+1}} - x_{t_i}$$

denote the i -th dyadic increment. Then the increments $\delta_i B^h$ ($i = 0, 1, \dots, 2^n - 1$) of the translated Brownian motion are independent random variables with distributions

$$\delta_i B^h = \delta_i B + \delta_i h \sim N(\delta_i h, (\delta t) \cdot I_d), \quad \delta t = 2^{-n}.$$

Consequently, the marginal distribution of $(B_{t_1}^h, B_{t_2}^h, \dots, B_{t_{2^n}}^h)$ is a normal distribution with density w.r.t. Lebesgue measure proportional to

$$\exp\left(-\sum_{i=0}^{2^n-1} \frac{|\delta_i x - \delta_i h|^2}{2\delta t}\right), \quad x = (x_{t_1}, x_{t_2}, \dots, x_{t_{2^n}}) \in \mathbb{R}^{2^n d}.$$

Since the normalization constant does not depend on h , the joint distribution of $(B_{t_1}^h, B_{t_2}^h, \dots, B_{t_{2^n}}^h)$ is absolutely continuous w.r.t. that of $(B_{t_1}, B_{t_2}, \dots, B_{t_{2^n}})$ with relative density

$$\exp\left(\sum \frac{\delta_i h}{\delta t} \cdot \delta_i x - \frac{1}{2} \sum \left|\frac{\delta_i h}{\delta t}\right|^2 \delta t\right). \quad (9.2.3)$$

Consequently, μ_h is always absolutely continuous w.r.t. μ_0 on each of the σ -algebras

$$\mathcal{F}_n = \sigma(X_{i \cdot 2^{-n}} : i = 0, 1, \dots, 2^n - 1), \quad n \in \mathbb{N},$$

with relative densities

$$Z_n = \exp\left(\sum_{i=0}^{2^n-1} \frac{\delta_i h}{\delta t} \cdot \delta_i X - \frac{1}{2} \sum_{i=0}^{2^n-1} \left|\frac{\delta_i h}{\delta t}\right|^2 \delta t\right). \quad (9.2.4)$$

(2). *Limit of local densities:* Suppose that h is absolutely continuous with

$$\int_0^1 |h'(t)|^2 dt < \infty.$$

We now identify the limit of the relative densities Z_n as $n \rightarrow \infty$.

First, we note that

$$\sum_{i=0}^{2^n-1} \left|\frac{\delta_i h}{\delta t}\right|^2 \delta t \longrightarrow \int_0^1 |h'(t)|^2 dt \quad \text{as } n \rightarrow \infty.$$

In fact, the sum on the right hand side coincides with the squared L^2 norm

$$\int_0^1 \left|dh/dt\Big|_{\sigma(\mathcal{D}_n)}\right|^2 dt$$

of the dyadic derivative

$$\frac{dh}{dt}\Big|_{\sigma(\mathcal{D}_n)} = \sum_{i=0}^{2^n-1} \frac{\delta_i h}{\delta t} \cdot I_{((i-1) \cdot 2^{-n}, i \cdot 2^{-n}]}$$

on the σ -algebra generated by the intervals $((i-1) \cdot 2^{-n}, i \cdot 2^{-n}]$. If h is absolutely continuous with $h' \in L^2(0, 1)$ then $\frac{dh}{dt}\Big|_{\sigma(\mathcal{D}_n)} \rightarrow h'(t)$ in $L^2(0, 1)$ by the L^2 martingale convergence theorem.

Furthermore, by Itô's isometry,

$$\sum_{i=0}^{2^n-1} \frac{\delta_i h}{\delta t} \cdot \delta_i X \rightarrow \int_0^1 h'(s) dX_s \quad \text{in } L^2(\mu_0) \text{ as } n \rightarrow \infty. \quad (9.2.5)$$

Indeed, the sum on the right-hand side is the Itô integral of the step function $\left. \frac{dh}{dt} \right|_{\sigma(\mathcal{D}_n)}$ w.r.t. X , and as remarked above, these step functions converge to h' in $L^2(0, 1)$. Along a subsequence, the convergence in (9.2.5) holds μ_0 -almost surely, and hence by (9.2.4),

$$\lim_{n \rightarrow \infty} Z_n = \exp \left(\int_0^1 h'(s) dX_s - \frac{1}{2} \int_0^1 |h'(s)|^2 ds \right) \quad \mu_0\text{-a.s.} \quad (9.2.6)$$

- (3). *Absolute continuity on \mathcal{F}_1^X* : We still assume $h' \in L^2(0, 1)$. Note that $\mathcal{F}_1^X = \sigma(\bigcup \mathcal{F}_n)$. Hence for proving that μ_h is absolutely continuous w.r.t. μ_0 on \mathcal{F}_1^X with density given by (9.2.6), it suffices to show that $\limsup Z_n < \infty$ μ_h -almost surely (i.e., the singular part in the Lebesgue decomposition of μ_h w.r.t. μ_0 vanishes). Since $\mu_h = \mu_0 \circ \tau_h^{-1}$, the process

$$W_t = X_t - h(t) \quad \text{is a Brownian motion w.r.t. } \mu_h,$$

and by (9.2.3) and (9.2.4),

$$Z_n = \exp \left(\sum_{i=0}^{2^n-1} \frac{\delta_i h}{\delta t} \cdot \delta_i W + \frac{1}{2} \sum_{i=0}^{2^n-1} \left| \frac{\delta_i h}{\delta t} \right|^2 \delta t \right).$$

Note that the minus sign in front of the second sum has turned into a plus by the translation! Arguing similarly as above, we see that along a subsequence, (Z_n) converges μ_h -almost surely to a finite limit:

$$\lim Z_n = \exp \left(\int_0^1 h'(s) dW_s + \frac{1}{2} \int_0^1 |h'(s)|^2 ds \right) \quad \mu_h\text{-a.s.}$$

Hence $\mu_h \ll \mu_0$ with density $\lim Z_n$.

(4). *Singularity on \mathcal{F}_1^X* : Conversely, let us suppose now that h is not absolutely continuous or h' is not in $L^2(0, 1)$. Then

$$\sum_{i=0}^{2^n-1} \left| \frac{\delta_i h}{\delta_i t} \right|^2 \delta t = \int_0^1 \left| \frac{dh}{dt} \right|_{\sigma(\mathcal{D}_n)}^2 dt \longrightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Since

$$\left\| \sum_{i=0}^{2^n-1} \frac{\delta_i h}{\delta t} \cdot \delta_i X \right\|_{L^2(\mu_0)} = \left(\sum_{i=0}^{2^n-1} \left(\frac{\delta_i h}{\delta t} \right)^2 \delta t \right)^{1/2},$$

we can conclude by (9.2.4) that

$$\lim Z_n = 0 \quad \mu_0\text{-almost surely,}$$

i.e., μ_h is singular w.r.t. μ_0 .

□

In Section 11.5, we will give an alternative proof of the Cameron-Martin Theorem.

Passage times for Brownian motion with constant drift

We now consider a one-dimensional Brownian motion with constant drift β , i.e., a process

$$Y_t = B_t + \beta t, \quad t \geq 0,$$

where B_t is a Brownian motion starting at 0 and $\beta \in \mathbb{R}$. We will apply the Cameron-Martin Theorem to compute the distributions of the first passage times

$$T_a^Y = \min\{t \geq 0 : Y_t = a\}, \quad a > 0.$$

Note that T_a^Y is also the first passage time to the line $t \mapsto a - \beta t$ for the Brownian motion (B_t) .

Theorem 9.7. For $a > 0$ and $\beta \in \mathbb{R}$, the restriction of the distribution of T_a^Y to $(0, \infty)$ is absolutely continuous with density

$$f_{a,\beta}(t) = \frac{a}{\sqrt{2\pi t^3}} \exp\left(-\frac{(a - \beta t)^2}{2t}\right).$$

In particular,

$$P[T_a^Y < \infty] = \int_0^\infty f_{a,\beta}(s) ds.$$

Proof. Let $h(t) = \beta t$. By the Cameron-Martin Theorem, the distribution μ_h of (Y_t) is absolutely continuous w.r.t. Wiener measure on \mathcal{F}_t^X with density

$$Z_t = \exp(\beta \cdot X_t - \beta^2 t/2).$$

Therefore, denoting by $T_a = \inf\{t \geq 0 : X_t = a\}$ the passage time of the canonical process, we obtain

$$\begin{aligned} P[T_a^Y \leq t] &= \mu_h[T_a \leq t] = E_{\mu_0}[Z_t ; T_a \leq t] \\ &= E_{\mu_0}[Z_{T_a} ; T_a \leq t] = E_{\mu_0}[\exp(\beta a - \frac{1}{2}\beta^2 T_a) ; T_a \leq t] \\ &= \int_0^t \exp(\beta a - \beta^2 s/2) f_{T_a}(s) ds \end{aligned}$$

by the optional sampling theorem. The claim follows by inserting the explicit expression for f_{T_a} derived in Corollary 1.25. \square

9.3 Girsanov transform

We will now extend the results in the previous section 9.2 considerably. To this end, we will consider locally absolutely continuous changes of measure with local densities of type

$$Z_t = \exp\left(\int_0^t G_s \cdot dX_s - \frac{1}{2} \int_0^t |G_s|^2 ds\right),$$

where (G_s) is an adapted process. Recall that the densities in the Cameron-Martin-Theorem took this form with the *deterministic* function $G_s = h'(s)$. We start with a general discussion about changing measure on filtered probability spaces that will be useful in other contexts as well.

Change of measure on filtered probability spaces

Let (\mathcal{F}_t) be a filtration on a measurable space (Ω, \mathcal{A}) , and fix $t_0 \in (0, \infty)$. We consider two probability measures P and Q on (Ω, \mathcal{A}) that are mutually absolutely continuous on the σ -algebra \mathcal{F}_{t_0} with relative density

$$Z_{t_0} = \left. \frac{dP}{dQ} \right|_{\mathcal{F}_{t_0}} > 0 \quad Q\text{-almost surely.}$$

Then P and Q are also mutually absolutely continuous on each of the σ -algebras \mathcal{F}_t , $t \leq t_0$, with Q - and P -almost surely strictly positive relative densities

$$Z_t = \left. \frac{dP}{dQ} \right|_{\mathcal{F}_t} = E_Q[Z_{t_0} | \mathcal{F}_t] \quad \text{and} \quad \left. \frac{dQ}{dP} \right|_{\mathcal{F}_t} = \frac{1}{Z_t}.$$

The process $(Z_t)_{t \leq t_0}$ is a martingale w.r.t. Q , and, correspondingly, $(1/Z_t)_{t \leq t_0}$ is a martingale w.r.t. P . From now on, we always choose a right continuous version of these martingales.

Lemma 9.8. 1) For any $0 \leq s \leq t \leq t_0$, and for any \mathcal{F}_t -measurable random variable $X : \Omega \rightarrow [0, \infty]$,

$$E_P[X | \mathcal{F}_s] = \frac{E_Q[X Z_t | \mathcal{F}_s]}{E_Q[Z_t | \mathcal{F}_s]} = \frac{E_Q[X Z_t | \mathcal{F}_s]}{Z_s} \quad P\text{-a.s. and } Q\text{-a.s.} \quad (9.3.1)$$

2) Suppose that $(M_t)_{t \leq t_0}$ is an (\mathcal{F}_t) adapted right continuous stochastic process. Then

- (i) M is a martingale w.r.t. P \Leftrightarrow $M \cdot Z$ is a martingale w.r.t. Q ,
- (ii) M is a local martingale w.r.t. P \Leftrightarrow $M \cdot Z$ is a local martingale w.r.t. Q .

Proof. 1) The right hand side of (9.3.1) is \mathcal{F}_s -measurable. Moreover, for any $A \in \mathcal{F}_s$,

$$\begin{aligned} E_P[E_Q[XZ_t|\mathcal{F}_s]/Z_s; A] &= E_Q[E_Q[XZ_t|\mathcal{F}_s]; A] \\ &= E_Q[XZ_t; A] = E_Q[X; A]. \end{aligned}$$

2) (i) is a direct consequence of 1). Moreover, by symmetry, it is enough to prove the implication " \Leftarrow " in (ii). Hence suppose that $M \cdot Z$ is a local Q -martingale with localizing sequence (T_n) . We show that M^{T_n} is a P -martingale, i.e.,

$$E_P[M_{t \wedge T_n}; A] = E_P[M_{s \wedge T_n}; A] \quad \text{for any } A \in \mathcal{F}_s, \quad 0 \leq s \leq t \leq t_0. \quad (9.3.2)$$

To verify (9.3.2), we first note that

$$E_P[M_{t \wedge T_n}; A \cap \{T_n \leq s\}] = E_P[M_{s \wedge T_n}; A \cap \{T_n \leq s\}] \quad (9.3.3)$$

since $t \wedge T_n = T_n = s \wedge T_n$ on $\{T_n \leq s\}$. Moreover, one verifies from the definition of the σ -algebra $\mathcal{F}_{s \wedge T_n}$ that for any $A \in \mathcal{F}_s$, the event $A \cap \{T_n > s\}$ is contained in $\mathcal{F}_{s \wedge T_n}$, and hence in $\mathcal{F}_{t \wedge T_n}$. Therefore,

$$\begin{aligned} E_P[M_{t \wedge T_n}; A \cap \{T_n > s\}] &= E_Q[M_{t \wedge T_n} Z_{t \wedge T_n}; A \cap \{T_n > s\}] \\ &= E_Q[M_{s \wedge T_n} Z_{s \wedge T_n}; A \cap \{T_n > s\}] = E_P[M_{s \wedge T_n}; A \cap \{T_n > s\}] \end{aligned} \quad (9.3.4)$$

by the martingale property for $(MZ)^{T_n}$, the optional sampling theorem, and the fact that $P \ll Q$ on $\mathcal{F}_{t \wedge T_n}$ with relative density $Z_{t \wedge T_n}$. (9.3.2) follows from (9.3.3) and (9.3.4). \square

Girsanov's Theorem

We now return to our original problem of identifying the change of measure induced by a random translation of the paths of a Brownian motion. Suppose that (X_t) is a Brownian motion in \mathbb{R}^d with $X_0 = 0$ w.r.t. the probability measure Q and the filtration (\mathcal{F}_t) , and fix $t_0 \in [0, \infty)$. Let

$$L_t = \int_0^t G_s \cdot dX_s, \quad t \geq 0,$$

with $G \in \mathcal{L}_{a,loc}^2(\mathbb{R}_+, \mathbb{R}^d)$. Then $[L]_t = \int_0^t |G_s|^2 ds$, and hence

$$Z_t = \exp\left(\int_0^t G_s \cdot dX_s - \frac{1}{2} \int_0^t |G_s|^2 ds\right) \quad (9.3.5)$$

is the exponential of L . In particular, since L is a local martingale w.r.t. Q , Z is a non-negative local martingale, and hence a supermartingale w.r.t. Q . It is a Q -martingale for $t \leq t_0$ if and only if $E_Q[Z_{t_0}] = 1$:

Exercise (Martingale property for exponentials). Let $(Z_t)_{t \in [0, t_0]}$ on (Ω, \mathcal{A}, Q) be a non-negative local martingale satisfying $Z_0 = 1$.

- a) Show that Z is a supermartingale.
- b) Prove that Z is a martingale if and only if $E_Q[Z_{t_0}] = 1$.

In order to use Z_{t_0} for changing the underlying probability measure on \mathcal{F}_{t_0} we have to assume the martingale property:

Assumption. $(Z_t)_{t \leq t_0}$ is a martingale w.r.t. Q .

Theorem 9.10 below implies that the assumption is satisfied if

$$E\left[\exp\left(\frac{1}{2} \int_0^t |G_s|^2 ds\right)\right] < \infty.$$

If the assumption holds then we can consider a probability measure P on \mathcal{A} with

$$\frac{dP}{dQ}\Big|_{\mathcal{F}_{t_0}} = Z_{t_0} \quad Q\text{-a.s.} \quad (9.3.6)$$

Note that P and Q are mutually absolutely continuous on \mathcal{F}_t for any $t \leq t_0$ with

$$\frac{dP}{dQ}\Big|_{\mathcal{F}_t} = Z_t \quad \text{and} \quad \frac{dQ}{dP}\Big|_{\mathcal{F}_t} = \frac{1}{Z_t}$$

both P - and Q -almost surely. We are now ready to prove one of the most important results of stochastic analysis:

Theorem 9.9 (Maruyama 1954, Girsanov 1960). *Suppose that X is a d -dimensional Brownian motion w.r.t. Q and $(Z_t)_{t \leq t_0}$ is defined by (9.3.5) with $G \in \mathcal{L}_{a,loc}^2(\mathbb{R}_+, \mathbb{R}^d)$. If $E_Q[Z_{t_0}] = 1$ then the process*

$$B_t := X_t - \int_0^t G_s ds, \quad t \leq t_0,$$

is a Brownian motion w.r.t. any probability measure P on \mathcal{A} satisfying (9.3.6).

Proof. By the extension of Lévy's characterization of Brownian motion to the multi-dimensional case, it suffices to show that $(B_t)_{t \leq t_0}$ is an \mathbb{R}^d -valued P -martingale with $[B^i, B^j]_t = \delta_{ij}t$ P -almost surely for any $i, j \in \{1, \dots, d\}$, cf. Theorem 11.2 below. Furthermore, by Lemma 9.8, and since P and Q are mutually absolutely continuous on \mathcal{F}_{t_0} , this holds true provided $(B_t Z_t)_{t \leq t_0}$ is an \mathbb{R}^d valued local martingale under Q , and $[B^i, B^j] = \delta_{ij}t$ Q -almost surely. The identity for the covariations holds since (B_t) differs from the Q -Brownian motion (X_t) only by a continuous finite variation process. To show that $B \cdot Z$ is a local Q -martingale, we apply Itô's formula: For $1 \leq i \leq d$,

$$\begin{aligned} d(B^i Z) &= B^i dZ + Z dB^i + d[B^i, Z] \\ &= B^i ZG \cdot dX + Z dX^i - Z G^i dt + ZG^i dt, \end{aligned} \tag{9.3.7}$$

where we have used that

$$d[B^i, Z] = ZG \cdot d[B^i, X] = ZG^i dt \quad Q\text{-almost surely.}$$

The right-hand side of (9.3.7) is a stochastic integral w.r.t. the Q -Brownian motion X , and hence a local Q -martingale. \square

The theorem shows that if X is a Brownian motion w.r.t. Q , and Z defined by (9.3.5) is a Q -martingale, then X satisfies

$$dX_t = G_t dt + dB_t.$$

with a P -Brownian motion B . This can be used to construct weak solutions of stochastic differential equations by changing the underlying probability measure, see Section 11.3

below. For instance, if we choose $G_t = b(X_t)$ then the Q -Brownian motion (X_t) is a solution to the SDE

$$dX_t = b(X_t) dt + dB_t,$$

where B is a Brownian motion under the modified probability measure P .

Furthermore, Girsanov's Theorem generalizes the Cameron-Martin Theorem to non-deterministic adapted translations

$$X_t(\omega) \longrightarrow X_t(\omega) - H_t(\omega), \quad H_t = \int_0^t G_s ds,$$

of a Brownian motion X .

Remark (Assumptions in Girsanov's Theorem).

- 1) Absolute continuity and adaptedness of the "translation process" $H_t = \int_0^t G_s ds$ are essential for the assertion of Theorem 9.9.
- 2) The assumption $E_Q[Z_{t_0}] = 1$ ensuring that $(Z_t)_{t \leq t_0}$ is a Q -martingale is not always satisfied – a sufficient condition is given in Theorem 9.10 below. If (Z_t) is not a martingale w.r.t. Q it can still be used to define a positive measure P_t with density Z_t w.r.t. Q on each σ -algebra \mathcal{F}_t . However, in this case, $P_t[\Omega] < 1$. The sub-probability measures P_t correspond to a transformed process with finite life-time.

Novikov's condition

To verify the assumption in Girsanov's theorem, we now derive a sufficient condition for ensuring that the exponential

$$Z_t = \exp(L_t - 1/2 [L]_t)$$

of a continuous local (\mathcal{F}_t) martingale (L_t) is a martingale. Recall that Z is always a non-negative local martingale, and hence a supermartingale w.r.t. (\mathcal{F}_t) .

Theorem 9.10 (Novikov 1971). *Let $t_0 \in \mathbb{R}_+$. If $E[\exp([L]_{t_0}/2)] < \infty$ then $(Z_t)_{t \leq t_0}$ is an (\mathcal{F}_t) martingale.*

We only prove the theorem under the slightly more restrictive condition

$$E[\exp(p[L]_t/2)] < \infty \quad \text{for some } p > 1. \quad (9.3.8)$$

This simplifies the proof considerably, and the condition is sufficient for many applications. For a proof in the general case and under even weaker assumptions see e.g. [37].

Proof. Let $(T_n)_{n \in \mathbb{N}}$ be a localizing sequence for the martingale Z . Then $(Z_{t \wedge T_n})_{t \geq 0}$ is a martingale for any n . To carry over the martingale property to the process $(Z_t)_{t \in [0, t_0]}$, it is enough to show that the random variables $Z_{t \wedge T_n}$, $n \in \mathbb{N}$, are uniformly integrable for each fixed $t \leq t_0$. However, for $c > 0$ and $p, q \in (1, \infty)$ with $p^{-1} + q^{-1} = 1$, we have

$$\begin{aligned} & E[Z_{t \wedge T_n}; Z_{t \wedge T_n} \geq c] \\ &= E\left[\exp\left(L_{t \wedge T_n} - \frac{p}{2}[L]_{t \wedge T_n}\right) \exp\left(\frac{p-1}{2}[L]_{t \wedge T_n}\right); Z_{t \wedge T_n} \geq c\right] \quad (9.3.9) \\ &\leq E\left[\exp\left(pL_{t \wedge T_n} - \frac{p^2}{2}[L]_{t \wedge T_n}\right)\right]^{1/p} \cdot E\left[\exp\left(q \cdot \frac{p-1}{2}[L]_{t \wedge T_n}\right); Z_{t \wedge T_n} \geq c\right]^{1/q} \\ &\leq E\left[\exp\left(\frac{p}{2}[L]_t\right); Z_{t \wedge T_n} \geq c\right]^{1/q} \end{aligned}$$

for any $n \in \mathbb{N}$. Here we have used Hölder's inequality and the fact that $\exp\left(pL_{t \wedge T_n} - \frac{p^2}{2}[L]_{t \wedge T_n}\right)$ is an exponential supermartingale. If $\exp\left(\frac{p}{2}[L]_t\right)$ is integrable then the right hand side of (9.3.9) converges to 0 uniformly in n as $c \rightarrow \infty$, because

$$P[Z_{t \wedge T_n} \geq 0] \leq c^{-1} E[Z_{t \wedge T_n}] \leq c^{-1} \longrightarrow 0$$

uniformly in n as $c \rightarrow \infty$. Hence $\{Z_{t \wedge T_n} : n \in \mathbb{N}\}$ is indeed uniformly integrable, and thus $(Z_t)_{t \in [0, t_0]}$ is a martingale. \square

Example (Bounded drifts). If $L_t = \int_0^t G_s \cdot dX_s$ with a Brownian motion (X_t) and an adapted process (G_t) that is uniformly bounded on $[0, t]$ for any finite t then the quadratic variation $[L]_t = \int_0^t |G_s|^2 ds$ is also bounded for finite t . Hence $\exp(L - \frac{1}{2}[L])$ is an (\mathcal{F}_t) martingale for $t \in [0, \infty)$.

Example (Option pricing in continuous time II: Risk-neutral measure). We consider the asset price model in continuous time introduced in the beginning of Chapter 8. The stock price is modelled by an SDE

$$dS_t = \alpha_t S_t dt + \sigma_t S_t dX_t, \quad (9.3.10)$$

and the interest rate is given by (R_t) . We assume that (X_t) is a Brownian motion and $(\alpha_t), (R_t), (\sigma_t)$ and $(1/\sigma_t)$ are adapted bounded continuous processes, all defined on a filtered probability space $(\Omega, \mathcal{A}, Q, (\mathcal{F}_t))$. Then the discounted asset price

$$\tilde{S}_t := \exp\left(-\int_0^t R_s ds\right) S_t$$

satisfies

$$d\tilde{S}_t = (\alpha_t - R_t)\tilde{S}_t dt + \sigma_t\tilde{S}_t dX_t = \sigma_t\tilde{S}_t dB_t, \quad (9.3.11)$$

where

$$B_t := X_t + \int_0^t \frac{\alpha_s - R_s}{\sigma_s} ds.$$

We can apply Girsanov's Theorem and the Novikov condition to conclude that the process (B_t) is a Brownian motion under a probability measure P on (Ω, \mathcal{A}) with local densities w.r.t. Q on \mathcal{F}_t given by

$$Z_t = \exp\left(\int_0^t G_s \cdot dX_s - \frac{1}{2} \int_0^t |G_s|^2 ds\right) \quad \text{where } G_t = (R_t - \alpha_t)/\sigma_t.$$

Therefore, by (9.3.11) and by the assumptions on the coefficients, the process (\tilde{S}_t) is a martingale under Q . The measure Q can now be used to compute option prices under a no-arbitrage assumption similarly to the discrete time case considered in Section 2.3 above, see Section 9.4.

9.4 Itô's Representation Theorem and Option Pricing

We now prove two basic representation theorems for functionals and martingales that are adapted w.r.t. the filtration generated by a Brownian motion. Besides their intrinsic interest, such representation theorems are relevant e.g. for the theory of financial markets, and for stochastic filtering. Throughout this section, (B_t) denotes a Brownian motion starting at 0 on a probability space (Ω, \mathcal{A}, P) , and

$$\mathcal{F}_t = \sigma(B_s : s \in [0, t])^P, \quad t \geq 0,$$

is the completed filtration generated by (B_t) . It is crucial that the filtration does not contain additional information. By the factorization lemma, this implies that \mathcal{F}_t measurable

random variables $F : \Omega \rightarrow \mathbb{R}$ are almost surely functions of the Brownian path $(B_s)_{s \leq t}$. Indeed, we will show that such functions can be represented as stochastic integrals.

Representation theorems for functions and martingales

The first version of Itô's Representation Theorem states that random variables that are measurable w.r.t. the σ -algebra $\mathcal{F}_1 = \mathcal{F}_1^{B,P}$ can be represented as stochastic integrals:

Theorem 9.11 (Itô). *For any function $F \in \mathcal{L}^2(\Omega, \mathcal{F}_1, P)$ there exists a unique process $G \in L_a^2(0, 1)$ such that*

$$F = E[F] + \int_0^1 G_s \cdot dB_s \quad P\text{-almost surely.} \quad (9.4.1)$$

An immediate consequence of Theorem 9.11 is a corresponding representation for martingales w.r.t. the **Brownian filtration** $\mathcal{F}_t = \mathcal{F}_t^{B,P}$:

Corollary 9.12 (Itô representation for martingales). *For any right-continuous L^2 -bounded (\mathcal{F}_t) martingale $(M_t)_{t \in [0,1]}$ there exists a unique process $G \in L_a^2(0, 1)$ such that*

$$M_t = M_0 + \int_0^t G_s \cdot dB_s \quad \text{for any } t \in [0, 1], \quad P\text{-a.s.}$$

The corollary is of fundamental importance in financial mathematics where it is related to completeness of financial markets. It also proves the remarkable fact that **every martingale w.r.t. the Brownian filtration has a continuous modification!** Of course, this result can not be true w.r.t. a general filtration.

We first show that the corollary follows from Theorem 9.11, and then we prove the theorem:

Proof of Corollary 9.12. If $(M_t)_{t \in [0,1]}$ is an L^2 bounded (\mathcal{F}_t) martingale then $M_1 \in \mathcal{L}^2(\Omega, \mathcal{F}_1, P)$, and

$$M_t = E[M_1 | \mathcal{F}_t] \quad \text{a.s. for any } t \in [0, 1].$$

Hence, by Theorem 9.11, there exists a unique process $G \in L_a^2(0, 1)$ such that

$$M_1 = E[M_1] + \int_0^1 G \cdot dB = M_0 + \int_0^1 G \cdot dB \quad \text{a.s.,}$$

and thus

$$M_t = E[M_1 | \mathcal{F}_t] = M_0 + \int_0^t G \cdot dB \quad \text{a.s. for any } t \in [0, 1].$$

Since both sides in the last equation are almost surely right continuous, the identity actually holds simultaneously for all $t \in [0, 1]$ with probability 1. \square

Proof of Theorem 9.11. Uniqueness. Suppose that (9.4.1) holds for two processes $G, \tilde{G} \in L_a^2(0, 1)$. Then

$$\int_0^1 G \cdot dB = \int_0^1 \tilde{G} \cdot dB,$$

and hence, by Itô's isometry,

$$\|G - \tilde{G}\|_{L^2(P \otimes \lambda)} = \left\| \int (G - \tilde{G}) \cdot dB \right\|_{L^2(P)} = 0.$$

Hence $G_t(\omega) = \tilde{G}_t(\omega)$ for almost every (t, ω) .

Existence. We prove the existence of a representation as in (9.4.1) in several steps – starting with “simple” functions F .

1. Suppose that $F = \exp(ip \cdot (B_t - B_s))$ for some $p \in \mathbb{R}^d$ and $0 \leq s \leq t \leq 1$. By Itô's formula,

$$\exp(ip \cdot B_t + \frac{1}{2}|p|^2 t) = \exp(ip \cdot B_s + \frac{1}{2}|p|^2 s) + \int_s^t \exp(ip \cdot B_r + \frac{1}{2}|p|^2 r) ip \cdot dB_r.$$

Rearranging terms, we obtain an Itô representation for F with a bounded adapted integrand G .

2. Now suppose that $F = \prod_{k=1}^n F_k$ where $F_k = \exp(ip_k \cdot (B_{t_k} - B_{t_{k-1}}))$ for some $n \in \mathbb{N}$,

$p_1, \dots, p_n \in \mathbb{R}^d$, and $0 \leq t_0 \leq t_1 \leq \dots \leq t_n \leq 1$. Denoting by G_k the bounded adapted process in the Itô representation for F_k , we have

$$F = \prod_{k=1}^n \left(E[F_k] + \int_{t_k}^{t_{k+1}} G^k \cdot dB \right).$$

We show that the right hand side can be written as the sum of $\prod_{k=1}^n E[F_k]$ and a stochastic integral w.r.t. B . For this purpose, it suffices to verify that the product of two stochastic integrals $X_t = \int_0^t G \cdot dB$ and $Y_t = \int_0^t H \cdot dB$ with bounded adapted processes G and H is the stochastic integral of a process in $L_a^2(0, 1)$ provided $\int_0^1 G_t \cdot H_t dt = 0$. This holds true, since by the product rule,

$$X_1 Y_1 = \int_0^1 X_t H_t \cdot dB_t + \int_0^1 Y_t G_t \cdot dB_t + \int_0^1 G_t \cdot H_t dt,$$

and $XH + YG$ is square-integrable by Itô's isometry.

3. Clearly, an Itô representation also holds for any linear combination of functions as in Step 2.

4. To prove an Itô representation for arbitrary functions in $\mathcal{L}^2(\Omega, \mathcal{F}_1, P)$, we first note that the linear combinations of the functions in Step 2 form a *dense* subspace of the Hilbert space $L^2(\Omega, \mathcal{F}_1, P)$. Indeed, if ϕ is an element in $L^2(\Omega, \mathcal{F}_1, P)$ that is orthogonal to this subspace then

$$E \left[\phi \prod_{k=1}^n \exp(ip_k \cdot (B_{t_k} - B_{t_{k-1}})) \right] = 0$$

for any $n \in \mathbb{N}$, $p_1, \dots, p_n \in \mathbb{R}^d$ and $0 \leq t_0 \leq t_1 \leq \dots \leq t_n \leq 1$. By Fourier inversion, this implies

$$E[\phi \mid \sigma(B_{t_k} - B_{t_{k-1}} : 1 \leq k \leq n)] = 0 \quad \text{a.s.}$$

for any $n \in \mathbb{N}$ and $0 \leq t_0 \leq \dots \leq t_n \leq 1$, and hence $\phi = 0$ a.s. by the Martingale Convergence Theorem.

Now fix an arbitrary function $F \in L^2(\Omega, \mathcal{F}_1, P)$. Then by Step 3, there exists a sequence (F_n) of functions in $L^2(\Omega, \mathcal{F}_1, P)$ converging to F in L^2 that have a representation of the form

$$F_n - E[F_n] = \int_0^1 G^{(n)} \cdot dB \tag{9.4.2}$$

with processes $G^{(n)} \in L_a^2(0, 1)$. As $n \rightarrow \infty$,

$$F_n - E[F_n] \longrightarrow F - E[F] \quad \text{in } L^2(P).$$

Hence, by (9.4.2) and Itô's isometry, $(G^{(n)})$ is a Cauchy sequence in $L^2(P \otimes \lambda_{(0,1)})$. Denoting by G the limit process, we obtain the representation

$$F - E[F] = \int_0^1 G \cdot dB$$

by taking the L^2 limit on both sides of (9.4.2). □

Application to option pricing

We return to the asset price model considered at the end of Section 9.3. For simplicity, we now assume that the coefficients in (9.3.10) and (9.3.11) are constant:

$$\alpha_t \equiv \alpha \in \mathbb{R}, \quad \sigma_t \equiv \sigma \in (0, \infty), \quad R_t \equiv r \in \mathbb{R}.$$

Then the change of measure is given by the local densities

$$Z_t = \exp\left(\frac{r - \alpha}{\sigma} X_t - \frac{1}{2} \left(\frac{r - \alpha}{\sigma}\right)^2 t\right), \quad (9.4.3)$$

and by (9.3.11), the discounted stock price is proportional to the Itô exponential of σB where $B_t = X_t + \frac{\alpha - r}{\sigma} t$ is a Brownian motion under the risk-neutral measure Q :

$$\tilde{S}_t = S_0 \cdot \exp(\sigma B_t - \sigma^2 t/2) \quad (9.4.4)$$

Now suppose that we want to compute the no-arbitrage price of an option. For example, let us consider a European call option where the payoff at the final time t_0 is given by

$$V_{t_0} = (S_{t_0} - K)^+$$

for a positive constant K . By (9.4.4), the discounted payoff

$$\tilde{V}_{t_0} = \left(\tilde{S}_{t_0} - e^{-rt_0} K\right)^+ \quad (9.4.5)$$

is an $\mathcal{F}_{t_0}^{B,P}$ measurable random variable. Therefore, by Itô's Representation Theorem and (9.4.4), there exists a process $G \in L_a^2(0, t_0)$ such that

$$\tilde{V}_{t_0} = E_P \left[\tilde{V}_{t_0} \right] + \int_0^{t_0} G_r dB_r = E_P \left[\tilde{V}_{t_0} \right] + \int_0^{t_0} \Phi_r d\tilde{S}_r,$$

where $\Phi_r := G_r/(\sigma\tilde{S}_r)$. Hence (Φ_r) is a replicating strategy for the option, i.e., investing Φ_r units in the stock and putting the remaining money on the bank account yields exactly the payoff for the option at time t_0 provided our initial capital is given by $E_P \left[\tilde{V}_{t_0} \right]$. Since otherwise there would be an arbitrage opportunity by selling the option and investing the gain by the strategy Φ , or conversely, we can conclude that under a no-arbitrage assumption, the only possible option price at time 0 is given by

$$E_P \left[\tilde{V}_{t_0} \right] = E_P \left[\left(S_0 e^{\sigma B_{t_0} - \sigma^2 t_0 / 2} - e^{-rt_0} K \right)^+ \right]$$

Noting that $B_{t_0} \sim N(0, t_0)$ under P , we obtain the *Black-Scholes formula* for the no-arbitrage price of a European call option. Notice in particular that the price does not depend on the usually unknown model parameter α (the mean rate of return).

Application to stochastic filtering

XXX to be included

Part III

Stochastic Analysis

Appendix

Appendix A

Conditional expectations

A.1 Conditioning on discrete random variables

We first consider conditioning on the value of a random variable $Y : \Omega \rightarrow S$ where S is countable. In this case, we can define the *conditional probability measure*

$$P[A | Y = z] = \frac{P[A \cap \{Y = z\}]}{P[Y = z]}, \quad A \in \mathcal{A},$$

and the *conditional expectations*

$$E[X | Y = z] = \frac{E[X; Y = z]}{P[Y = z]}, \quad X \in \mathcal{L}^1(\Omega, \mathcal{A}, P),$$

for any $z \in S$ with $P[Y = z] > 0$ in an elementary way. Note that for $z \in S$ with $P[Y = z] = 0$, the conditional probabilities are not defined.

Conditional expectations as random variables

It will turn out to be convenient to consider the conditional probabilities and expectations not as functions of the outcome z , but as functions of the random variable Y . In this way, the conditional expectations become random variables:

Definition (Conditional expectation given a discrete random variable). Let $X : \Omega \rightarrow \mathbb{R}$ be a random variable such that $E[X^-] < \infty$, and let $Y : \Omega \rightarrow S$ be a discrete random variable. The random variable $E[X | Y]$ that is P -almost surely uniquely defined by

$$E[X | Y] := g(Y) = \sum_{z \in S} g(z) \cdot I_{\{Y=z\}}$$

with

$$g(z) := \begin{cases} E[X | Y = z] & \text{if } P[Y = z] > 0 \\ \text{arbitrary} & \text{if } P[Y = z] = 0 \end{cases}$$

is called **(a version of the) conditional expectation of X given Y** . For an event $A \in \mathcal{A}$, the random variable

$$P[A | Y] := E[I_A | Y]$$

is called **(a version of the) conditional probability of A given Y** .

The conditional expectation $E[X | Y]$ and the conditional probability $P[A | Y]$ are again random variables. They take the values $E[X | Y = z]$ and $P[A | Y = z]$, respectively, on the sets $\{Y = z\}$, $z \in S$ with $P[Y = z] > 0$. On each of the null sets $\{Y = z\}$, $z \in S$ with $P[Y = z] = 0$, an arbitrary constant value is assigned to the conditional expectation. Hence the definition is only almost surely unique.

Characteristic properties of conditional expectations

Let $X : \Omega \rightarrow \mathbb{R}$ be a non-negative or integrable random variable on a probability space (Ω, \mathcal{A}, P) . The following alternative characterisation of the conditional expectation of X given Y can be verified in an elementary way:

Theorem A.1. A real random variable $\bar{X} \geq 0$ (or $\bar{X} \in \mathcal{L}^1$) on (Ω, \mathcal{A}, P) is a version of the conditional expectation $E[X | Y]$ if and only if

(I) $\bar{X} = g(Y)$ for a function $g : S \rightarrow \mathbb{R}$, and

(II) $E[\bar{X} \cdot f(Y)] = E[X \cdot f(Y)]$ for all non-negative or bounded functions $f : S \rightarrow \mathbb{R}$, respectively.

A.2 General conditional expectations

If Y is a real-valued random variable on a probability space (Ω, \mathcal{A}, P) with continuous distribution function, then $P[Y = z] = 0$ for any $z \in \mathbb{R}$. Therefore, conditional probabilities given $Y = z$ can not be defined in the same way as above. Alternatively, one could try to define conditional probabilities given Y as limits:

$$P[A | Y = z] = \lim_{h \searrow 0} P[A | z - h \leq Y \leq z + h]. \quad (\text{A.2.1})$$

In certain cases this is possible but in general, the existence of the limit is not guaranteed.

Instead, the characterization in Theorem A.1 is used to provide a definition of conditional expectations given general random variables Y . The conditional probability of a fixed event A given Y can then be defined almost surely as a special case of a conditional expectation:

$$P[A | Y] := E[I_A | Y]. \quad (\text{A.2.2})$$

Note, however, that in general, the exceptional set will depend on the event A !

The factorization lemma

We first prove an important measure theoretic statement.

Theorem A.2 (Factorization lemma). *Suppose that (S, \mathcal{S}) is a measurable space and $Y : \Omega \rightarrow S$ is a map. Then a map $X : \Omega \rightarrow \mathbb{R}$ is measurable w.r.t. $\sigma(Y)$ if and only if*

$$X = f(Y) = f \circ Y$$

for a \mathcal{S} -measurable function $f : S \rightarrow \mathbb{R}$.

$$\begin{array}{ccc}
 & X & \\
 & \curvearrowright & \\
 (\Omega, \sigma(Y)) & \xrightarrow{Y} & (S, \mathcal{S}) \longrightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))
 \end{array}$$

Proof. (1). If $X = f \circ Y$ for a measurable function f , then

$$X^{-1}(B) = Y^{-1}(f^{-1}(B)) \in \sigma(Y) \quad \text{holds for all } B \in \mathcal{B}(\mathbb{R}),$$

as $f^{-1}(B) \in \mathcal{S}$. Therefore, X is $\sigma(Y)$ -measurable.

(2). Conversely, we have to show that $\sigma(Y)$ -measurability of X implies that X is a measurable function of Y . This is done in several steps:

(a) If $X = I_A$ is an indicator function of an set $A \in \sigma(Y)$, then $A = Y^{-1}(B)$ with $B \in \mathcal{S}$, and thus

$$X(\omega) = I_{Y^{-1}(B)}(\omega) = I_B(Y(\omega)) \quad \text{for all } \omega \in \Omega.$$

(b) For $X = \sum_{i=1}^n c_i I_{A_i}$ with $A_i \in \sigma(Y)$ and $c_i \in \mathbb{R}$ we have correspondingly

$$X = \sum_{i=1}^n c_i I_{B_i}(Y),$$

where wobei B_i are sets in \mathcal{S} such that $A_i = Y^{-1}(B_i)$.

(c) For an arbitrary non-negative, $\sigma(Y)$ -measurable map $X : \Omega \rightarrow \mathbb{R}$, there exists a sequence of $\sigma(Y)$ -measurable elementary functions such that $X_n \nearrow X$. By (b), $X_n = f_n(Y)$ with \mathcal{S} -measurable functions f_n . Hence

$$X = \sup X_n = \sup f_n(Y) = f(Y),$$

where $f = \sup f_n$ is again \mathcal{S} -measurable.

(d) For a general $\sigma(Y)$ -measurable map $X : \Omega \rightarrow \mathbb{R}$, both X^+ and X^- are measurable functions of Y , hence X is a measurable function of Y as well.

□

The factorization lemma can be used to rephrase the *characterizing properties* (I) and (II) of conditional expectations in Theorem A.1 in the following way:

\bar{X} is a version of $E[X | Y]$ if and only if

- (i) \bar{X} ist $\sigma(Y)$ -messbar;
- (ii) $E[\bar{X}; A] = E[X; A]$ für alle $A \in \sigma(Y)$.

The equivalence of (I) und (i) is a consequence of the factorization lemma, and the equivalence of (II) and (ii) follows by monotone classes, since (ii) states that

$$E[\bar{X} \cdot I_B(Y)] = E[X \cdot I_B(Y)] \quad \text{holds for all } B \in \mathcal{S}.$$

Conditional expectations given σ -algebras

A remarkable consequence of the characterization of conditional expectations by Conditions (i) and (ii) is that the *conditional expectation* $E[X | Y]$ depends on the random variable Y only via the σ -algebra $\sigma(Y)$ generated by Y ! If two random variables Y and Z are functions of each other then $\sigma(Y) = \sigma(Z)$, and hence the conditional expectations $E[X | Y]$ and $E[X | Z]$ coincide (with probability 1). Therefore it is plausible to define directly the conditional expectation given a σ -Algebra. The σ -algebra (e.g. $\sigma(Y)$, or $\sigma(Y_1, \dots, Y_n)$) then describes the available “information” on which we are conditioning.

The characterization of conditional expectations by (i) and (ii) can be extended immediately to the case of general conditional expectations given a σ -algebra or given arbitrary random variables. To this end let $X : \Omega \rightarrow \bar{\mathbb{R}}$ be a non-negative (or integrable) random variable on a probability space (Ω, \mathcal{A}, P) .

Definition (Conditional expectation, general). (I). Let $\mathcal{F} \subseteq \mathcal{A}$ be a σ -algebra. A non-negative (or integrable) random variable $\bar{X} : \Omega \rightarrow \bar{\mathbb{R}}$ is called a **version of the conditional expectation** $E[X | \mathcal{F}]$ iff:

- (a) \bar{X} is \mathcal{F} -measurable, and
 (b) $E[\bar{X} ; A] = E[X ; A]$ for any $A \in \mathcal{F}$.

(2). For arbitrary random variables Y, Y_1, Y_2, \dots, Y_n on (Ω, \mathcal{A}, P) we define

$$\begin{aligned} E[X | Y] &:= E[X | \sigma(Y)], \\ E[X | Y_1, \dots, Y_n] &:= E[X | (Y_1, \dots, Y_n)] = E[X | \sigma(Y_1, \dots, Y_n)]. \end{aligned}$$

(3). For an event $A \in \mathcal{A}$ we define

$$P[A | \mathcal{F}] := E[I_A | \mathcal{F}], \quad \text{and correspondingly} \quad P[A | Y] = E[I_A | Y].$$

Remark. By monotone classes it can be shown that Condition (b) is equivalent to:

$$(b') \quad E[\bar{X} \cdot Z] = E[X \cdot Z] \quad \text{for any non-negative (resp. bounded) } \mathcal{F}\text{-measurable } Z : \Omega \rightarrow \mathbb{R}.$$

Theorem A.3 (Existence and uniqueness of conditional expectations). *Let $X \geq 0$ or $X \in \mathcal{L}^1$, and let $\mathcal{F} \subseteq \mathcal{A}$ be a σ -algebra. Then:*

- (1). *There exists a version of the conditional expectation $E[X | \mathcal{F}]$.*
 (2). *Any two versions coincide P -almost surely.*

Proof. Existence can be shown as a consequence of the Radon-Nikodym theorem. In Theorem A.8 below, we give a different proof of existence that only uses elementary methods.

For proving uniqueness let \bar{X} and \tilde{X} be two versions of $E[X | \mathcal{F}]$. Then both \bar{X} and \tilde{X} are \mathcal{F} -measurable, and

$$E[\bar{X} ; A] = E[\tilde{X} ; A] \quad \text{for any } A \in \mathcal{F}.$$

Therefore, $\bar{X} = \tilde{X}$ P -almost surely. □

Properties of conditional expectations

Starting from the definition, we now derive several basic properties of conditional expectations that are used frequently:

Theorem A.4. *Let X, Y and X_n ($n \in \mathbb{N}$) be non-negative or integrable random variables on (Ω, \mathcal{A}, P) , and let $\mathcal{F}, \mathcal{G} \subseteq \mathcal{A}$ be σ -algebras.*

The following assertions hold:

(1). *Linearity: $E[\lambda X + \mu Y | \mathcal{F}] = \lambda E[X | \mathcal{F}] + \mu E[Y | \mathcal{F}]$ P -almost surely for any $\lambda, \mu \in \mathbb{R}$.*

(2). *Monotonicity: If $X \geq 0$ P -almost surely, then $E[X | \mathcal{F}] \geq 0$ P -almost surely.*

(3). *If $X = Y$ P -almost surely then $E[X | \mathcal{F}] = E[Y | \mathcal{F}]$ P -almost surely.*

(4). *Monotone Convergence: If (X_n) is increasing with $X_1 \geq 0$, then*

$$E[\sup X_n | \mathcal{F}] = \sup E[X_n | \mathcal{F}] \quad P\text{-almost surely.}$$

(5). *Tower Property: If $\mathcal{G} \subseteq \mathcal{F}$ then*

$$E[E[X | \mathcal{F}] | \mathcal{G}] = E[X | \mathcal{G}] \quad P\text{-almost surely.}$$

In particular,

$$E[E[X | Y, Z] | Y] = E[X | Y] \quad P\text{-almost surely.}$$

(6). *Taking out what is known: Let Y be \mathcal{F} -measurable such that $Y \cdot X \in \mathcal{L}^1$ or ≥ 0 . Then*

$$E[Y \cdot X | \mathcal{F}] = Y \cdot E[X | \mathcal{F}] \quad P\text{-almost surely.}$$

(7). *Independence: If X is independent of \mathcal{F} then $E[X | \mathcal{F}] = E[X]$ P -almost surely.*

(8). *Let (S, \mathcal{S}) and (T, \mathcal{T}) be measurable spaces. If $Y : \Omega \rightarrow S$ is \mathcal{F} -measurable and $X : \Omega \rightarrow T$ is independent of \mathcal{F} , then for any product-measurable function $f : S \times T \rightarrow [0, \infty)$ we have*

$$E[f(X, Y) | \mathcal{F}](\omega) = E[f(X, Y(\omega))] \quad \text{für } P\text{-fast alle } \omega.$$

Proof. (1). Aus der Linearität des Erwartungswertes folgt, dass $\lambda E[X | \mathcal{F}] + \mu E[Y | \mathcal{F}]$ eine Version der bedingten Erwartung $E[\lambda X + \mu Y | \mathcal{F}]$ ist.

(2). Sei \bar{X} eine Version von $E[X | \mathcal{F}]$. Aus $X \geq 0$ P -fast sicher folgt wegen $\{\bar{X} < 0\} \in \mathcal{F}$:

$$E[\bar{X}; \bar{X} < 0] = E[X; \bar{X} < 0] \geq 0,$$

und damit $\bar{X} \geq 0$ P -fast sicher.

(3). Dies folgt unmittelbar aus (1) und (2).

(4). Ist $X_n \geq 0$ und monoton wachsend, dann ist $\sup E[X_n | \mathcal{F}]$ eine nichtnegative \mathcal{F} -messbare Zufallsvariable (mit Werten in $[0, \infty]$), und nach dem "klassischen" Satz von der monotonen Konvergenz gilt:

$$E[\sup E[X_n | \mathcal{F}] \cdot Z] = \sup E[E[X_n | \mathcal{F}] \cdot Z] = \sup E[X_n \cdot Z] = E[\sup X_n \cdot Z]$$

für jede nichtnegative \mathcal{F} -messbare Zufallsvariable Z . Also ist $\sup E[X_n | \mathcal{F}]$ eine Version der bedingten Erwartung von $\sup X_n$ gegeben \mathcal{F} .

(5). Wir zeigen, dass jede Version von $E[X | \mathcal{G}]$ auch eine Version von $E[E[X | \mathcal{F}] | \mathcal{G}]$ ist, also die Eigenschaften (i) und (ii) aus der Definition der bedingten Erwartung erfüllt:

(i) $E[X | \mathcal{G}]$ ist nach Definition \mathcal{G} -messbar.

(ii) Für $A \in \mathcal{G}$ gilt auch $A \in \mathcal{F}$, und somit $E[E[X | \mathcal{G}]; A] = E[X; A] = E[E[X | \mathcal{F}]; A]$.

(6) und (7). Auf ähnliche Weise verifiziert man, dass die Zufallsvariablen, die auf der rechten Seite der Gleichungen in (6) und (7) stehen, die definierenden Eigenschaften der bedingten Erwartungen auf der linken Seite erfüllen (Übung).

(8). Dies folgt aus (6) und (7) in drei Schritten:

(a) Gilt $f(x, y) = g(x) \cdot h(y)$ mit messbaren Funktionen $g, h \geq 0$, dann folgt nach (6) und (7) P -fast sicher:

$$\begin{aligned} E[f(X, Y) | \mathcal{F}] &= E[g(X) \cdot h(Y) | \mathcal{F}] = h(Y) \cdot E[g(X) | \mathcal{F}] \\ &= h(Y) \cdot E[g(X)], \end{aligned}$$

und somit

$$E[f(X, Y) | \mathcal{F}](\omega) = E[g(X) \cdot h(Y(\omega))] = E[f(X, Y(\omega))] \quad \text{für } P\text{-fast alle } \omega.$$

- (b) Um die Behauptung für Indikatorfunktionen $f(x, y) = I_B(x, y)$ von produktmessbaren Mengen B zu zeigen, betrachten wir das Mengensystem

$$\mathcal{D} = \{B \in \mathcal{S} \otimes \mathcal{T} \mid \text{Behauptung gilt für } f = I_B\}.$$

\mathcal{D} ist ein Dynkinsystem, das nach (a) alle Produkte $B = B_1 \times B_2$ mit $B_1 \in \mathcal{S}$ und $B_2 \in \mathcal{T}$ enthält. Also gilt auch

$$\mathcal{D} \supseteq \sigma(\{B_1 \times B_2 \mid B_1 \in \mathcal{S}, B_2 \in \mathcal{T}\}) = \mathcal{S} \otimes \mathcal{T}.$$

- (c) Für beliebige produktmessbare Funktionen $f : S \times T \rightarrow \mathbb{R}_+$ folgt die Behauptung nun durch maßtheoretische Induktion.

□

Remark (Convergence theorems for conditional expectations). The Monotone Convergence Theorem (Property (4)) implies versions of Fatou's Lemma and of the Dominated Convergence Theorem for conditional expectations. The proofs are similar to the unconditioned case.

The last property in Theorem A.4 is often very useful. For independent random variables X and Y it implies

$$E[f(X, Y) | Y](\omega) = E[f(X, Y(\omega))] \quad \text{für } P\text{-fast alle } \omega, \quad (\text{A.2.3})$$

We stress that independence of X and Y ist essential for (A.2.3) to hold true. The application of (A.2.3) without independence is a common mistake in computations with conditional expectations.

A.3 Conditional expectation as best L^2 -approximation

In this section we show that the conditional expectation of a square integrable random variable X given a σ -algebra \mathcal{F} can be characterized alternatively as the best approximation of X in the subspace of \mathcal{F} -measurable, square integrable random variables, or,

equivalently, as the orthogonal projection of X onto this subspace. Besides obvious applications to non-linear predictions, this point of view is also the basis for a simple existence proof of conditional expectations

Jensen's inequality

Jensen's inequality is valid for conditional expectations as well. Let (Ω, \mathcal{A}, P) be a probability space, $X \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$ an integrable random variable, and $\mathcal{F} \subseteq \mathcal{A}$ a σ -algebra.

Theorem A.5 (Jensen). *If $u : \mathbb{R} \rightarrow \mathbb{R}$ is a convex function with $u(X) \in \mathcal{L}^1$ or $u \geq 0$, then*

$$E[u(X) | \mathcal{F}] \geq u(E[X | \mathcal{F}]) \quad P\text{-almost surely.}$$

Proof. Jede konvexe Funktion u lässt sich als Supremum von abzählbar vielen affinen Funktionen darstellen, d.h. es gibt $a_n, b_n \in \mathbb{R}$ mit

$$u(x) = \sup_{n \in \mathbb{N}} (a_n x + b_n) \quad \text{für alle } x \in \mathbb{R}.$$

Zum Beweis betrachtet man die Stützgeraden an allen Stellen einer abzählbaren dichten Teilmenge von \mathbb{R} , siehe z.B. [Williams: Probability with martingales, 6.6]. Wegen der Monotonie und Linearität der bedingten Erwartung folgt

$$E[u(X) | \mathcal{F}] \geq E[a_n X + b_n | \mathcal{F}] = a_n \cdot E[X | \mathcal{F}] + b_n$$

P -fast sicher für alle $n \in \mathbb{N}$, also auch

$$E[u(X) | \mathcal{F}] \geq \sup_{n \in \mathbb{N}} (a_n \cdot E[X | \mathcal{F}] + b_n) \quad P\text{-fast sicher.}$$

□

Corollary A.6 (L^p -contractivity). *The map $X \mapsto E[X | \mathcal{F}]$ is a contraction on $\mathcal{L}^p(\Omega, \mathcal{A}, P)$ for every $p \geq 1$, i.e.,*

$$E[|E[X | \mathcal{F}]|^p] \leq E[|X|^p] \quad \text{for any } X \in \mathcal{L}^1(\Omega, \mathcal{A}, P).$$

Proof. Nach der Jensenschen Ungleichung gilt:

$$|E[X | \mathcal{F}]|^p \leq E[|X|^p | \mathcal{F}] \quad P\text{-fast sicher.}$$

Die Behauptung folgt durch Bilden des Erwartungswertes. \square

The proof of the corollary shows in particular that for a random variable $X \in \mathcal{L}^p$, the conditional expectation $E[X | \mathcal{F}]$ is contained in \mathcal{L}^p as well. We now restrict ourselves to the case $p = 2$.

Conditional expectation as best L^2 -prediction value

The space $L^2(\Omega, \mathcal{A}, P) = \mathcal{L}^2(\Omega, \mathcal{A}, P) / \sim$ of equivalence classes of square integrable random variables is a Hilbert space with inner product $(X, Y)_{L^2} = E[XY]$. If $\mathcal{F} \subseteq \mathcal{A}$ is a sub- σ -algebra then $L^2(\Omega, \mathcal{F}, P)$ is a **closed subspace** of $L^2(\Omega, \mathcal{A}, P)$, because limits of \mathcal{F} -measurable random variables are \mathcal{F} -measurable as well. For $X \in \mathcal{L}^2(\Omega, \mathcal{A}, P)$, each version of the conditional expectation $E[X | \mathcal{F}]$ is contained in the subspace $L^2(\Omega, \mathcal{F}, P)$ by Jensen's inequality. Furthermore, the conditional expectation respects equivalence classes, see Theorem A.3. Therefore, $X \mapsto E[X | \mathcal{F}]$ induces a linear map from the Hilbert space $L^2(\Omega, \mathcal{A}, P)$ of equivalence classes onto the subspace $L^2(\Omega, \mathcal{F}, P)$.

Theorem A.7 (Characterization of the conditional expectation as best L^2 approximation and as orthogonal projection). *For $Y \in \mathcal{L}^2(\Omega, \mathcal{F}, P)$ the following statements are all equivalent:*

(1). Y is a version of the conditional expectation $E[X | \mathcal{F}]$.

(2). Y is a “**best approximation**” of X in the subspace $\mathcal{L}^2(\Omega, \mathcal{F}, P)$, i.e.,

$$E[(X - Y)^2] \leq E[(X - Z)^2] \quad \text{for any } Z \in \mathcal{L}^2(\Omega, \mathcal{F}, P).$$

(3). Y is a version of the **orthogonal projection** of X onto the subspace $L^2(\Omega, \mathcal{F}, P) \subseteq L^2(\Omega, \mathcal{A}, P)$, i.e.,

$$E[(X - Y) \cdot Z] = 0 \quad \text{for any } Z \in \mathcal{L}^2(\Omega, \mathcal{F}, P).$$




Figure A.1: $X \mapsto E[X | \mathcal{F}]$ as orthogonal projection onto the subspace $L^2(\Omega, \mathcal{F}, P)$.

Proof. **(1)** \iff **(3)**: Für $Y \in \mathcal{L}^2(\Omega, \mathcal{F}, P)$ gilt:

$$\begin{aligned} & Y \text{ ist eine Version von } E[X | \mathcal{F}] \\ \iff & E[Y \cdot I_A] = E[X \cdot I_A] \quad \text{für alle } A \in \mathcal{F} \\ \iff & E[Y \cdot Z] = E[X \cdot Z] \quad \text{für alle } Z \in \mathcal{L}^2(\Omega, \mathcal{F}, P) \\ \iff & E[(X - Y) \cdot Z] = 0 \quad \text{für alle } Z \in \mathcal{L}^2(\Omega, \mathcal{F}, P) \end{aligned}$$

Hierbei zeigt man die zweite Äquivalenz mit den üblichen Fortsetzungsverfahren (maßtheoretische Induktion).

(3) \Rightarrow (2): Sei Y eine Version der orthogonalen Projektion von X auf $L^2(\Omega, \mathcal{F}, P)$. Dann gilt für alle $Z \in \mathcal{L}^2(\Omega, \mathcal{F}, P)$:

$$\begin{aligned} E[(X - Z)^2] &= E[((X - Y) + (Y - Z))^2] \\ &= E[(X - Y)^2] + E[(Y - Z)^2] + 2E[(X - Y) \underbrace{(Y - Z)}_{\in \mathcal{L}^2(\Omega, \mathcal{F}, P)}] \\ &\geq E[(X - Y)^2] \end{aligned}$$

Hierbei haben wir im letzten Schritt verwendet, dass $Y - Z$ im Unterraum $\mathcal{L}^2(\Omega, \mathcal{F}, P)$ enthalten, also orthogonal zu $X - Y$ ist.

(2) \Rightarrow (3): Ist umgekehrt Y eine beste Approximation von X in $\mathcal{L}^2(\Omega, \mathcal{F}, P)$ und $Z \in \mathcal{L}^2(\Omega, \mathcal{F}, P)$, dann gilt

$$\begin{aligned} E[(X - Y)^2] &\leq E[(X - Y + tZ)^2] \\ &= E[(X - Y)^2] + 2tE[(X - Y)Z] + t^2E[Z^2] \end{aligned}$$

für alle $t \in \mathbb{R}$, also $E[(X - Y) \cdot Z] = 0$.

□

The equivalence of (2) and (3) is a well-known functional analytic statement: the best approximation of a vector in a closed subspace of a Hilbert space is the orthogonal projection of the vector onto this subspace. The geometric intuition behind this fact is indicated in Figure A.1.

Theorem A.7 is a justification for the interpretation of the conditional expectation as a prediction value. For example, by the factorization lemma, $E[X | Y]$ is the best L^2 -prediction for X among all functions of type $g(Y)$, $g : \mathbb{R} \rightarrow \mathbb{R}$ measurable.

Existence of conditional expectations

By the characterization of the conditional expectation as the best L^2 -approximation, the existence of conditional expectations of square integrable random variables is an

immediate consequence of the existence of the best approximation of a vector in a closed subspace of a Hilbert space. By monotone approximation, the existence of conditional expectations of general non-negative random variables then follows easily.

Theorem A.8 (Existence of conditional expectations). *For every random variable $X \geq 0$ or $X \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$, and every σ -algebra $\mathcal{F} \subseteq \mathcal{A}$, there exists a version of the conditional expectation $E[X | \mathcal{F}]$.*

Proof. (1). Wir betrachten zunächst den Fall $X \in \mathcal{L}^2(\Omega, \mathcal{A}, P)$. Wie eben bemerkt, ist der Raum $L^2(\Omega, \mathcal{F}, P)$ ein abgeschlossener Unterraum des Hilbertraums $L^2(\Omega, \mathcal{A}, P)$. Sei $d = \inf\{\|Z - X\|_{L^2} \mid Z \in \mathcal{L}^2(\Omega, \mathcal{F}, P)\}$ der Abstand von X zu diesem Unterraum. Um zu zeigen, dass eine beste Approximation von X in $L^2(\Omega, \mathcal{F}, P)$ existiert, wählen wir eine Folge (X_n) aus diesem Unterraum mit $\|X_n - X\|_{L^2} \rightarrow d$. Mithilfe der Parallelogramm-Identität folgt für $n, m \in \mathbb{N}$:

$$\begin{aligned} \|X_n - X_m\|_{L^2}^2 &= \|(X_n - X) - (X_m - X)\|_{L^2}^2 \\ &= 2 \cdot \|X_n - X\|_{L^2}^2 + 2 \cdot \|X_m - X\|_{L^2}^2 - \|(X_n - X) + (X_m - X)\|_{L^2}^2 \\ &= 2 \cdot \underbrace{\|X_n - X\|_{L^2}^2}_{\rightarrow d^2} + 2 \cdot \underbrace{\|X_m - X\|_{L^2}^2}_{\rightarrow d^2} - 4 \underbrace{\left\| \frac{X_n + X_m}{2} - X \right\|_{L^2}^2}_{\leq d^2}, \end{aligned}$$

und damit

$$\limsup_{n, m \rightarrow \infty} \|X_n - X_m\|_{L^2}^2 \leq 0.$$

Also ist die Minimalfolge (X_n) eine Cauchyfolge in dem vollständigen Raum $L^2(\Omega, \mathcal{F}, P)$, d.h. es existiert ein $Y \in \mathcal{L}^2(\Omega, \mathcal{F}, P)$ mit

$$\|X_n - Y\|_{L^2} \rightarrow 0.$$

Für Y gilt

$$\|Y - X\|_{L^2} = \left\| \lim_{n \rightarrow \infty} X_n - X \right\|_{L^2} \leq \liminf_{n \rightarrow \infty} \|X_n - X\|_{L^2} \leq d,$$

d.h. Y ist die gesuchte Bestapproximation, und damit eine Version der bedingten Erwartung $E[X | \mathcal{F}]$.

- (2). Für eine beliebige nichtnegative Zufallsvariable X auf (Ω, \mathcal{A}, P) existiert eine monoton wachsende Folge (X_n) nichtnegativer quadratintegrierbarer Zufallsvariablen mit $X = \sup_n X_n$. Man verifiziert leicht, dass $\sup_n E[X_n | \mathcal{F}]$ eine Version von $E[X | \mathcal{F}]$ ist.
- (3). Entsprechend verifiziert man, dass für allgemeine $X \in \mathcal{L}^1(\Omega, \mathcal{A}, P)$ durch $E[X | \mathcal{F}] = E[X^+ | \mathcal{F}] - E[X^- | \mathcal{F}]$ eine Version der bedingten Erwartung gegeben ist.

□

Index

- absolute continuity
 - of probability measures, 139
- absolutely continuous
 - function, 140
 - locally -, 141
- adapted, 59
- Brownian motion
 - canonical model for -, 20
 - definition, 13
 - geometric -, 246
 - Lévy process, 14
 - Markov property, 15
- canonical model for Brownian motion, 20
- colored noise, 9
- completion of a σ -algebra, 99
- Conditional Variance Process, 71
- Dirichlet problem, 88
- Doob
 - maximal inequality, 172
- Filtration, 59, 97
 - completed -, 99, 171
 - right-continuous -, 98
- first hitting time, 82
- Gaussian
 - measure, 29
- Gaussian process, 24
- Green function, 231
- harmonic, 66
 - sub-, 66
 - super-, 66
- heat kernel, 231
- i.i.d., 9
- integrable
 - uniformly -, 84
- Integral
 - Stochastic -
 - Discrete -, 77
- Itô diffusion, 256
- Itô integral
 - for predictable step functions, 165
 - backward -, 201
- Itô map, 167, 175
- Itô's formula
 - without probabilities, 198
- Lebesgue decomposition, 143

- Lebesgue density, 145
- likelihood
- ratio, 141
- local martingale, 184
- localizing sequence, 183
- lognormal distribution, 260
- Markov
- chain, 66
- Martingale, 60
- transform, 77
 - Definition, 97
 - Sub-, 60
 - Super-, 60
- Martingale Problem
- Solution of-, 74
- martingale transform, 165
- measure
- Gaussian -, 29
 - Wiener -, 20
- Mehler formula, 272
- mesh size
- of a partition, 161
- modification
- of a martingale, 172
- noise
- colored -, 9
 - white -, 9
- occupation time density, 231
- Ornstein-Uhlenbeck process, 268
- orthonormal basis, 36
- partition, 161
- predictable step functions, 165
- probability measure
- absolute continuity of -, 139
 - singular -, 139
- Process
- adapted -, 59
 - Conditional Variance -, 71
 - predictable -, 69
- process
- Gaussian -, 24
- Quadratic Variation, 196
- Radon-Nikodym
- derivative, 141
- scale function, 257
- Schauder functions, 36
- singularity
- of probability measures, 139
- Stochastic integral
- Definition, 175
 - Discrete -, 77
- stopping time, 55
- Stratonovich integral, 201
- subharmonic, 66
- superharmonic, 66
- uniformly integrable, 84
- usual conditions, 99
- Variance Process
- Conditional -, 71

white noise, 9, 25

Wiener

- measure, 20

Bibliography

- [1]
- [2]
- [3] *Sur l'équation de Kolmogoroff, par W. Doeblin.* Éditions Elsevier, Paris, 2000. C. R. Acad. Sci. Paris Sér. I Math. **331** (2000), Special Issue.
- [4] Robert J. Adler and Jonathan E. Taylor. *Random fields and geometry.* Springer Monographs in Mathematics. Springer, New York, 2007.
- [5] David Applebaum. *Lévy Processes and Stochastic Calculus.* Cambridge UP, Cambridge, 2004.
- [6] Ole E. Barndorff-Nielsen. Processes of normal inverse Gaussian type. *Finance Stoch.*, 2(1):41–68, 1998.
- [7] Richard F. Bass. *Stochastic Processes.* Cambridge University Press, Cambridge, 2011.
- [8] Jean Bertoin. *Lévy Processes.* Cambridge UP, Cambridge, 1998.
- [9] Jean-Michel Bismut. *Large deviations and the Malliavin calculus.* Birkhaeuser, Boston, 1994.
- [10] Amir Dembo and Ofur Zeitouni. *Large Deviations Techniques and Applications.* Springer, 1998, Berlin.
- [11] Frank den Hollander. *Large Deviations.* American Mathematical Society, Providence, 2008.

- [12] Andreas Eberle. Markov processes.
<http://www.uni-bonn.de/~eberle/MarkovProcesses/MPSkript.pdf>,
2009.
- [13] Andreas Eberle. Wahrscheinlichkeitstheorie und Stochastische Prozesse.
<http://www.uni-bonn.de/~eberle/StoPr/WthSkriptneu.pdf>,
2010.
- [14] Andreas Eberle. Introduction to Stochastic Analysis.
<http://www.uni-bonn.de/~eberle/StoAn1011/StoAnSkriptneu.pdf>,
2011.
- [15] David Elworthy. *On the Geometry of Diffusion Operators and Stochastic Flows*. Springer, 1999, Berlin.
- [16] Stewart N. Ethier and Thomas G. Kurtz. *Markov Processes*. Wiley-Interscience, Hoboken, NJ, 2005.
- [17] William Feller. *Introduction to Probability Theory and Its Applications*. Wiley, New York, 1957.
- [18] Peter K. Friz and Martin Hairer. *A course on rough paths*. Universitext. Springer, Cham, 2014. With an introduction to regularity structures.
- [19] Peter K. Friz and Nicolas Victoir. *Multidimensional Stochastic Processes as Rough Paths*. Cambridge UP, Cambridge, 2010.
- [20] W. Hackenbroch and A. Thalmaier. *Stochastic Analysis*. Teubner, 1991.
- [21] Martin Hairer. On malliavin's proof of hörmander's theorem. *Bull. Sci. Math.*, 135(6-7):650–666, 2011.
- [22] Elton Hsu. *Stochastic Analysis on Manifolds*. Oxford University Press, Oxford, 2002.
- [23] Nobuyuki Ikeda and Takehiko Ed. Watanabe. *Stochastic Differential Equations and Diffusion Processes*. North-Holland, Amsterdam, 1981.

-
- [24] Jean Jacod and Albert Shiryaev. *Limit Theorems for Stochastic Processes*. Springer, Berlin, 2003.
- [25] Ioannis Karatzas and Steven E. Shreve. *Brownian motion and stochastic calculus*. Springer, New York, 2010.
- [26] Peter E. Kloeden and Eckhard Platen. *Numerical solution of stochastic differential equations*, volume 23 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1992.
- [27] Hiroshi Kunita. *Stochastic Flows and Stochastic Differential Equations*. Cambridge University Press, Cambridge, 1997.
- [28] Andreas Kyprianou. *Introductory Lectures on Fluctuations of Lévy Processes with Applications*. Springer, Berlin, 2006.
- [29] Terry J. Lyons. *Differential Equations Driven by Rough Paths, Ecole d'Été de Probabilités de Saint-Flour XXXIV*. Springer, 2004.
- [30] Zhi-Ming Ma and Michael Röckner. *Introduction to the Theory of (Non-Symmetric) Dirichlet Forms*. Springer, Berlin, 1992.
- [31] Dilip B. Madan and Eugene Seneta. The variance gamma model for share market returns. *Journal of Business*, 63(4):511–524, 1990.
- [32] Paul Malliavin. *Stochastic Analysis*. Springer, Berlin, 2002.
- [33] G. N. Milstein and M. V. Tretyakov. *Stochastic numerics for mathematical physics*. Scientific Computation. Springer-Verlag, Berlin, 2004.
- [34] David Nualart. *The Malliavin Calculus and Related Topics*. Springer, Berlin, 2006.
- [35] Giuseppe Da Prato. *Introduction to Stochastic Analysis and Malliavin Calculus*. Edizioni della Normale, Pisa, 2009.
- [36] Philip E. Protter. *Stochastic Integration and Differential Equations*. Springer, Berlin, 2005.

- [37] Daniel Revuz and Marc Yor. *Continuous martingales and Brownian motion*. Springer, Berlin, 2005.
- [38] L. C. G. Rogers and David Williams. *Diffusions, Markov processes and martingales, Vol. 2: Ito calculus*. Cambridge UP, Cambridge, 2008.
- [39] Steven E. Shreve. *Stochastic calculus for finance. II*. Springer Finance. Springer-Verlag, New York, 2004. Continuous-time models.
- [40] Daniel W. Stroock and S.R. S. Varadhan. *Multidimensional Diffusion Processes*. Springer, Berlin, 2005.
- [41] Shinzo Watanabe. *Lectures on stochastic differential equations and Malliavin calculus*. Springer, Berlin, 1984.