

Diophantine properties of numbers related to Catalan's constant

Tanguy Rivoal and Wadim Zudilin

1 Introduction

This article deals with Dirichlet's beta function which is defined by

$$\beta(s) := \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^s}$$

for any complex s such that $\operatorname{Re}(s) > 0$. Euler showed that, for odd s , $\beta(s)$ is a rational multiple of π^s , namely

$$\beta(2n+1) = \frac{(-1)^n E_{2n}}{2^{2n+2} (2n)!} \pi^{2n+1} \quad \text{for } n = 0, 1, 2, \dots$$

Here the integers E_{2n} are the *Euler numbers*, whose exponential-generating function is given by the formula

$$\frac{1}{\cosh(z)} = \sum_{k=0}^{\infty} \frac{E_k}{k!} z^k.$$

Therefore, Lindemann's theorem “ π is transcendental over \mathbb{Q} ” implies that the odd beta values $\beta(2n+1)$ are transcendental over \mathbb{Q} .

For even s , no Euler-type formula is known, and no method has yet been found for determining the arithmetic nature of even beta values. Thus, the situation for the beta function appears to be the same as the situation for the Riemann zeta function

$$\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s}.$$

This is not completely true since the equivalent of Apéry's theorem “ $\zeta(3) \notin \mathbb{Q}$ ” [Ap] for Catalan's constant

$$G := \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2} = \beta(2)$$

has not yet been proved.

At the end of the last century, in [Ri1] (see also [BR]), a method was introduced for tackling the problem of determining the arithmetical nature of odd zeta values :

Theorem. *For any odd $a \geq 3$, the dimension of the \mathbb{Q} -vector space spanned by $1, \zeta(3), \zeta(5), \dots, \zeta(a)$ is greater than $\frac{1}{3} \log(a)$.*

It follows that infinitely many odd zeta values are irrational. The search of new irrational odd zeta values was started in [Ri2] “At least one of the nine numbers $\zeta(5), \zeta(7), \dots, \zeta(21)$ is irrational” and improved in [Zu2], [Zu3] :

Theorem. *At least one of the numbers $\zeta(5), \zeta(7), \zeta(9), \zeta(11)$ is irrational.*

We refer the interested reader to [Zu1] for similar results. Our aim is to translate both theorems above into the context of the beta function. We will prove the following two results.

Theorem 1. *Let a be an even integer ≥ 2 and let $\delta(a)$ denote the dimension of the \mathbb{Q} -vector space spanned by $1, \beta(2), \beta(4), \dots, \beta(a)$. Then*

$$\delta(a) \geq \frac{1 + o(1)}{2 + \log(2)} \log(a) \quad \text{as } a \rightarrow \infty. \quad (1)$$

Moreover, there exists a computable constant $c_0 > 0$ such that $\delta(a) > c_0 \log(a)$ for any even integer a .

Theorem 2. *At least one of the seven numbers*

$$\beta(2), \beta(4), \beta(6), \beta(8), \beta(10), \beta(12), \beta(14) \quad (2)$$

is irrational.

2 The hypergeometric construction

Let q be an integer ≥ 4 and let $\underline{\eta} = (\eta_0, \eta_1, \dots, \eta_q)$ be a family of $q + 1$ integral parameters satisfying the conditions

$$0 < \eta_j < \frac{1}{2}\eta_0 \quad \text{for } j = 1, \dots, q \quad \text{and} \quad \eta_1 + \eta_2 + \dots + \eta_q \leq \frac{q-1}{2}\eta_0. \quad (3)$$

For any non-negative integer n consider the tuple $\underline{h} = (h_0, h_1, \dots, h_q)$ given by the formulae

$$h_0 = \eta_0 n + 1, \quad h_j = \eta_j n + \frac{1}{2} \quad \text{for } j = 1, \dots, q-1, \quad \text{and} \quad h_q = \eta_q n + 1. \quad (4)$$

Both proofs are based on a detailed study of the following hypergeometric series

$$\begin{aligned}
S_n(z) &= S_{n,q,\underline{\eta}}(z) \\
&:= \gamma_n \cdot \frac{\Gamma(1+h_0) \prod_{j=1}^q \Gamma(h_j)}{\prod_{j=1}^q \Gamma(1+h_0-h_j)} \\
&\quad \times {}_{q+2}F_{q+1} \left(\begin{matrix} h_0, 1+\frac{1}{2}h_0, & h_1, & \dots, & h_q \\ \frac{1}{2}h_0, & 1+h_0-h_1, \dots, & 1+h_0-h_q \end{matrix} \middle| z^{-1} \right) \quad (5) \\
&= \gamma_n \cdot \sum_{t=0}^{\infty} (2t+h_0) \frac{\Gamma(t+h_0) \prod_{j=1}^q \Gamma(t+h_j)}{\Gamma(t+1) \prod_{j=1}^q \Gamma(t+1+h_0-h_j)} z^{-t}, \quad (6)
\end{aligned}$$

which, by the conditions (3) on the parameters $\underline{\eta}$, is convergent for any complex z , $|z| \geq 1$. The factor

$$\gamma_n = \gamma_{n,q,\underline{\eta}} := \frac{\prod_{j=1}^{q-1} (h_0 - 2h_j)!}{(h_q - 1)!^2} = \frac{\prod_{j=1}^{q-1} ((\eta_0 - 2\eta_j)n)!}{(\eta_q n)!^2}$$

that appears in the definition of $S_n(z)$ corresponds to an arithmetic normalization of the series.

There is an alternative ‘‘Pochhammer representation’’ of (6),

$$S_n(z) = \gamma_n \cdot \sum_{t=0}^{\infty} (2t+h_0) \frac{(t+1)_{h_q-1} (t+1+h_0-h_q)_{h_q-1}}{\prod_{j=1}^{q-1} (t+h_j)_{1+h_0-2h_j}} z^{-t}, \quad (7)$$

where $(a)_l := \Gamma(a+l)/\Gamma(a) = a(a+1)\cdots(a+l-1)$ for $l \in \mathbb{N}$ denotes Pochhammer’s symbol, which is very similar to the one introduced in [Ri1] and [BR] in the study of odd zeta values.

The key point is that the series in (5) is of a special kind. Namely it is a *well-poised hypergeometric series* in the terminology of [Bai], [Sl]: the sum of corresponding top and bottom parameters in (5) is invariant,

$$h_0 + 1 = \left(1 + \frac{1}{2}h_0\right) + \frac{1}{2}h_0 = h_1 + (1 + h_0 - h_1) = \cdots = h_q + (1 + h_0 - h_q).$$

The expression (5) gives rise to the following Euler–Beukers-type integral representation of $S_n(z)$.

Lemma 1. *The formula*

$$\begin{aligned}
S_n(z) &= \frac{\Gamma(1+h_0)}{\Gamma(h_q)^2 \Gamma(1+h_0-2h_q)} \\
&\quad \times z^{h_0} \int_{[0,1]^q} \frac{\prod_{j=1}^q t_j^{h_j-1} (1-t_j)^{h_0-2h_j}}{(z-t_1 t_2 \cdots t_q)^{h_0}} \frac{z+t_1 t_2 \cdots t_q}{z-t_1 t_2 \cdots t_q} dt_1 dt_2 \cdots dt_q \quad (8)
\end{aligned}$$

holds for any $z \in \mathbb{C}$, $|z| \geq 1$.

Proof. Using Section 4.1 of [Sl], we can derive the following integral representation for all complex z such that $|z| > 1$:

$$S_n(z) = \frac{\Gamma(1 + h_0)}{\Gamma(h_q)^2 \Gamma(1 + h_0 - 2h_q)} \times \int_{[0,1]^q} \prod_{j=1}^q t_j^{h_j-1} (1 - t_j)^{h_0-2h_j} {}_2F_1 \left(\begin{matrix} h_0, 1 + \frac{1}{2}h_0 \\ \frac{1}{2}h_0 \end{matrix} \middle| t_1 \cdots t_q z^{-1} \right) dt_1 \cdots dt_q.$$

Formula (1.5.21) in [Sl] implies that, for $|t| < 1$,

$${}_2F_1 \left(\begin{matrix} h_0, 1 + \frac{1}{2}h_0 \\ \frac{1}{2}h_0 \end{matrix} \middle| t \right) = \frac{1+t}{(1-t)^{h_0+1}}$$

and the desired integral formula follows for $|z| > 1$. In the case $|z| = 1$, *mutatis mutandis*, the proof is the same as the one of Lemma 2 in [BR]. \square

The beta values are particular cases as specialisations of the Lerch function

$$\Phi_s(z) = \Phi \left(s, \frac{1}{2}; z \right) := \sum_{k=0}^{\infty} \frac{z^k}{(k + 1/2)^s}$$

for complex z , $|z| \leq 1$, and s , $\operatorname{Re}(s) > 1$, since

$$\beta(s) = 2^s \Phi_s(-1) \quad \text{for } s \in \mathbb{C}, \operatorname{Re}(s) > 0.$$

3 The structure of $S_{n,q,\underline{\eta}}(z)$

We need to decompose $S_{n,q,\underline{\eta}}(-1)$ as a linear form in 1 and even beta values only. The following Lemma 2 is almost the same as the corresponding one in [BR] and we only include a sketch of the proof. We would like to emphasise the fact that it is essentially the well-poisedness of the series that enables us to produce this parity-dependant behaviour when $z = \pm 1$.

Consider the rational function summand in (7) or (6)

$$\begin{aligned} R_n(t) &= R_{n,q,\underline{\eta}}(t) := \gamma_n \cdot (2t + h_0) \frac{(t+1)_{h_q-1} (t+1+h_0-h_q)_{h_q-1}}{\prod_{j=1}^{q-1} (t+h_j)_{1+h_0-2h_j}} \\ &= \gamma_n \cdot (2t + h_0) \frac{\prod_{j=0}^q \Gamma(t+h_j)}{\prod_{j=0}^q \Gamma(t+1+h_0-h_j)} \end{aligned} \quad (9)$$

and define the rational numbers $c_{kj} = c_{k,j,n,q,\underline{\eta}}$, where $j = 1, \dots, q-1$ and $k = h_j + \frac{1}{2}, \dots, h_0 - h_j + \frac{1}{2}$, as the coefficients in the partial-fraction decomposition

$$R_n(t) = \sum_{j=1}^{q-1} \sum_{k=h_j+\frac{1}{2}}^{h_0-h_j+\frac{1}{2}} \frac{c_{kj}}{(t+k-\frac{1}{2})^j}. \quad (10)$$

Using this notation, we define the polynomials

$$P_{jn}(z) = P_{j,n,q,\underline{\eta}}(z) := \sum_{k=h_j+\frac{1}{2}}^{h_0-h_j+\frac{1}{2}} c_{kj} z^{k-1} \quad \text{for } j = 1, \dots, q-1, \quad (11)$$

$$P_{0n}(z) = P_{0,n,q,\underline{\eta}}(z) := \sum_{j=1}^{q-1} \sum_{k=h_j+\frac{1}{2}}^{h_0-h_j+\frac{1}{2}} c_{kj} \sum_{l=1}^{k-h_q} \frac{z^{k-l}}{(l-1/2)^j}, \quad (12)$$

which belong to $\mathbb{Q}[z]$ (for $k < h_q$ in (12), we mean $\sum_{l=1}^{k-h_q} := -\sum_{l=k-h_q}^{-1}$, and the sum is 0 if $k = h_q$).

Remark. Following the usual procedure, we can show that

$$c_{kj} = D_{q-1-j} \left(R_n(t) \left(t + k - \frac{1}{2} \right)^{q-1} \right) \Big|_{t=-k+\frac{1}{2}} \in \mathbb{Q}, \quad (13)$$

where

$$D_\lambda := \frac{1}{\lambda!} \left(\frac{d}{dt} \right)^\lambda, \quad \lambda = 0, 1, 2, \dots, \quad (14)$$

for the numbers c_{kj} , $k = h_j + \frac{1}{2}, \dots, h_0 - h_j + \frac{1}{2}$ and $j = 1, \dots, q-1$.

Lemma 2. For all complex z , $|z| \geq 1$,

$$S_n(z) = \sum_{j=1}^{q-1} P_{jn}(z) \Phi_j(z^{-1}) - P_{0n}(z). \quad (15)$$

Moreover, if q is even then for any odd $j \geq 1$, $P_{jn}(-1) = 0$ and hence

$$S_n(-1) = \sum_{l=1}^{q/2-1} P_{2l,n}(-1) \cdot 2^{-2l} \beta(2l) - P_{0n}(-1) \quad (16)$$

is a linear form in 1 and the even beta values $\beta(2), \beta(4), \dots, \beta(q-2)$.

Proof. Using the decomposition (10) and the fact that the rational function (9) has zeros at the points $t = -1, -2, \dots, 1 - h_q$, we obtain

$$\begin{aligned} S_n(z) &= \sum_{t=0}^{\infty} R_n(t) z^{-t} = \sum_{t=1-h_q}^{\infty} R_n(t) z^{-t} \\ &= \sum_{j=1}^{q-1} \sum_{k=h_j+\frac{1}{2}}^{h_0-h_j+\frac{1}{2}} c_{kj} z^{k-1} \sum_{t=1-h_q}^{\infty} \frac{z^{-(t+k-1)}}{(t+k-\frac{1}{2})^j} \\ &= \sum_{j=1}^{q-1} \sum_{k=h_j+\frac{1}{2}}^{h_0-h_j+\frac{1}{2}} c_{kj} z^{k-1} \left(\sum_{l=1}^{\infty} - \sum_{l=1}^{k-h_q} \right) \frac{z^{-l+1}}{(l-\frac{1}{2})^j} \end{aligned}$$

which gives the desired representation (15).

Further, the definition (9) gives that

$$R_n(-t - h_0) = -(-1)^{h_0(q-1)} R_n(t),$$

hence $c_{h_0-k+1,j} = (-1)^{h_0(q-1)+j-1} c_{k,j}$ for $k = h_j + \frac{1}{2}, \dots, h_0 - h_j + \frac{1}{2}$ and $j = 1, \dots, q-1$. Thus, as in Lemmas 1 of [Ri1], [Ri2], [BR] it follows that the polynomials $P_{jn}(z)$ are “ \pm -reciprocal”:

$$z^{h_0-1} P_{jn}(z^{-1}) = (-1)^{h_0(q-1)+j-1} P_{jn}(z), \quad j = 1, \dots, q-1,$$

and the substitution $z = -1$ gives us to the relations

$$P_{jn}(-1) = (-1)^{h_0q+j} P_{jn}(-1), \quad j = 1, \dots, q-1,$$

that is $P_{jn}(-1) = 0$ for odd j , $1 \leq j \leq q-1$, provided that q is even. \square

4 Asymptotic behaviour of the series

In this paragraph, we determine the asymptotics of $S_{n,q,\underline{\eta}}(-1)$.

Lemma 3. *As $n \rightarrow \infty$, the limit of $|S_n(-1)|^{1/n}$ exists and is equal to*

$$\varphi = \varphi_{q,\underline{\eta}} := \frac{\eta_0^{\eta_0} \eta_q^{-2\eta_q}}{(\eta_0 - 2\eta_q)^{\eta_0 - 2\eta_q}} \cdot \max_{\underline{t} \in]0,1]^q} \frac{\prod_{j=1}^q t_j^{\eta_j} (1 - t_j)^{\eta_0 - 2\eta_j}}{(1 + t_1 t_2 \cdots t_q)^{\eta_0}}. \quad (17)$$

In particular, $\varphi_{q,\underline{\eta}} \neq 0$.

Proof. Using the integral representation (8) for $S_n(-1)$ together with the correspondence (4) between the parameters $\underline{\eta}$ and \underline{h} , the existence and the theoretical value of the limit follow easily from the Laplace method and arguments such as those used in the first proof of Lemma 3 in [BR]. The non-vanishing of $\varphi_{q,\underline{\eta}}$ is clear from its definition. \square

Remark. In fact, the maximum in (17) is achieved at a point $\underline{t} = (t_1, \dots, t_q) \in]0, 1[^q$ that can be calculate as a function of some root $t_0 = t_0(\underline{\eta}) \in]0, 1[$ of the polynomial

$$Q(t) = Q_{q,\underline{\eta}}(t) := t \prod_{j=1}^q ((\eta_0 - \eta_j) - \eta_j t) - \prod_{j=1}^q (\eta_j - (\eta_0 - \eta_j) t). \quad (18)$$

Indeed, setting

$$F(\underline{t}) = \log \left(\frac{\prod_{j=1}^q t_j^{\eta_j} (1 - t_j)^{\eta_0 - 2\eta_j}}{(1 + t_1 t_2 \cdots t_q)^{\eta_0}} \right),$$

we see that the extrema in (17) are attained at points satisfying the conditions

$$0 = \frac{\partial F}{\partial t_j} = \frac{1}{t_j} \left(\eta_j - \frac{\eta_0 - 2\eta_j}{1 - t_j} t_j - \eta_0 \frac{t_1 t_2 \cdots t_q}{1 + t_1 t_2 \cdots t_q} \right), \quad j = 1, 2, \dots, q. \quad (19)$$

Set $t_0 := t_1 t_2 \cdots t_q$; then by (19) we obtain

$$\eta_j - \frac{\eta_0 - 2\eta_j}{1 - t_j} t_j = \frac{\eta_0 t_0}{1 + t_0}, \quad j = 1, 2, \dots, q,$$

that yields

$$t_j = \frac{\eta_j - \eta_0 t_0 / (1 + t_0)}{\eta_0 - \eta_j - \eta_0 t_0 / (1 + t_0)} = \frac{\eta_j - (\eta_0 - \eta_j) t_0}{(\eta_0 - \eta_j) - \eta_j t_0}, \quad j = 1, 2, \dots, q.$$

Substitution of these last expressions into $t_0 = t_1 t_2 \cdots t_q$ implies that t_0 is a zero of the polynomial (18).

One can ask whether t_0 is unique. The the answer is not obvious but, fortunately, we do not need it! Indeed, Lemma 3 is enough to prove Theorem 1 and the uniqueness will be numerically verified in the case of Theorem 2.

5 Denominators of linear forms

In this paragraph, we prove a very general result which gives a common denominator of the coefficients of the linear forms in even beta values which are constructed in Lemma 2. Denote the least common multiple of the numbers $1, 2, \dots, n$ by d_n and note the following consequence of the prime number theorem:

$$\lim_{n \rightarrow \infty} d_n^{1/n} = e. \quad (20)$$

To find convenient denominators for the numbers (13), we split the function $R_n(t)$ into a product of rational functions of the form

$$G(t) = G(a, b; t) := \frac{(t+b)_{a-b}}{(a-b)!} = \frac{1}{\Gamma(a-b+1)} \frac{\Gamma(t+a)}{\Gamma(t+b)} \quad \text{if } a \geq b, \quad (21)$$

and

$$H(t) = H(a, b; t) := \frac{(b-a-1)!}{(t+a)_{b-a}} = \Gamma(b-a) \frac{\Gamma(t+a)}{\Gamma(t+b)} \quad \text{if } a < b, \quad (22)$$

with an integral difference $a - b$.

For $a, b \in \mathbb{Z}$, the functions (21) are *integer-valued polynomials* and the following property is well known (see, e.g., [Bak], Lemma 1).

Lemma 4. *Let a, b be integers, $a \geq b$, and let $G(t)$ be defined in (21). Then for any non-negative integer λ , we have*

$$4^{a-b} d_{a-b}^\lambda \cdot D_\lambda G(t)|_{t=-k+\frac{1}{2}} \in \mathbb{Z} \quad \text{for } k \in \mathbb{Z},$$

where the differential operator D_λ is defined in (14).

The next assertion is proved in [BR], proof of Lemma 5.

Lemma 5. *Let a, b, a_0, b_0 be integers, $a_0 \leq a < b \leq b_0$, and let $H(t)$ be defined by (22). Then for any non-negative integer λ , we have*

$$d_{b_0-a_0-1}^\lambda \cdot D_\lambda (H(t)(t+k))|_{t=-k} \in \mathbb{Z} \quad \text{for } k = a_0, a_0 + 1, \dots, b_0 - 1.$$

The following result about the arithmetic of the polynomials (11) and (12) is a simple consequence of Lemmas 4 and 5. Suppose q is even and the parameters $\underline{\eta}$ are ordered as follows:

$$\eta_1 \leq \eta_2 \leq \dots \leq \eta_{q-1}.$$

Set

$$m_0 = \max\{\eta_0 - 2\eta_1, \eta_q\},$$

$$m_j = \max\{m_0, 2(\eta_0 - \eta_j - \eta_q), 2(\eta_q - \eta_j)\} \quad \text{for } j = 1, \dots, q-1$$

(obviously, $m_1 \geq m_2 \geq \dots \geq m_{q-1} \geq m_0$).

Lemma 6. *Under the above conditions, for $j = 0, 1, \dots, q-1$,*

$$d_{m_1 n} d_{m_2 n} \cdots d_{m_{q-1-j} n} \cdot 16^{\eta_q n} \cdot P_{jn}(z) \in \mathbb{Z}[z]. \quad (23)$$

Proof. Decomposing

$$R_{n,q,\underline{\eta}}(t) \left(t + k - \frac{1}{2}\right)^{q-1} = (2t + h_0) \cdot G(h_q, 1; t) \cdot G(h_0, 1 + h_0 - h_q; t)$$

$$\times \prod_{j=1}^{q-1} H(h_j, 1 + h_0 - h_j; t) \left(t + k - \frac{1}{2}\right),$$

and applying Lemmas 4, 5 and the Leibniz rule for differentiating a product to formulae (13), we get that

$$d_{m_0 n}^{q-1-j} \cdot 16^{\eta_q n} \cdot c_{kj} \in \mathbb{Z} \quad (24)$$

for $k = h_j + \frac{1}{2}, \dots, h_0 - h_j + \frac{1}{2}$ and $j = 1, \dots, q-1$. On the other hand, we get

$$\prod_{i=1}^j d_{\max\{2h_0-2h_i-2h_q, 2h_q-2h_i\}} \cdot \sum_{l=1}^{k-h_q} \frac{1}{(2l-1)^j} \in \mathbb{Z}. \quad (25)$$

By (24) and (25), we finally arrive at the desired result (23) for the polynomials $P_{jn}(z)$ and $P_{0n}(z)$. \square

Remark. Lemma 6 is not the end of our arithmetic investigations : we will slightly modify the relations (23) for Theorem 1 in the next paragraph and then sharpen them by using p -adic valuation of the numbers (13) for Theorem 2 in Paragraph 7.

6 Proof of Theorem 1

This paragraph is devoted to the proof of Theorem 1, which is based on the following theorem of Nesterenko [Ne].

Linear independence criterion. *Let $\theta_1, \theta_2, \dots, \theta_N$, with $N \geq 2$, be real numbers, and assume there exist N sequences of integers $(p_{ln})_{n \geq 0}$ such that:*

$$(i) \quad \alpha_1^{n+o(n)} \leq \left| \sum_{l=1}^N p_{ln} \theta_l \right| \leq \alpha_2^{n+o(n)} \text{ for certain } 0 < \alpha_1 \leq \alpha_2 < 1;$$

$$(ii) \text{ for all } l = 1, \dots, N, \quad |p_{ln}| \leq \beta^{n+o(n)} \text{ for certain } \beta > 1.$$

Then,

$$\dim_{\mathbb{Q}}(\mathbb{Q}\theta_1 + \mathbb{Q}\theta_2 + \dots + \mathbb{Q}\theta_N) \geq \frac{\log(\beta) - \log(\alpha_1)}{\log(\beta) - \log(\alpha_1) + \log(\alpha_2)}.$$

In order to complete the proof, we now take the following special values for the parameters $\underline{\eta}$:

$$\eta_0 = 2r + 1, \quad \eta_1 = \eta_2 = \dots = \eta_q = r, \quad \text{where } 1 \leq r \leq \frac{q-1}{2}. \quad (26)$$

The new parameter r is an even integer that will be chosen later as a function of q . As the criterion explicitly indicates, we also have to find the asymptotic behaviour of the coefficients of the linear form in this special case (26).

Lemma 7. *For all $j = 0, 1, \dots, q-1$,*

$$\limsup_{n \rightarrow \infty} |P_{jn}(-1)|^{1/n} \leq 2^{q-2r-1} (2r+1)^{2r+1} r^{-2r}.$$

We omit the proof of this lemma since it follows the same lines as [BR], Lemma 4: we use Cauchy's formula

$$c_{kj} = \frac{1}{2\pi i} \int_{|t+k-\frac{1}{2}|=\frac{1}{2}} R_n(t) (t+k-\frac{1}{2})^{j-1} dt$$

for the numbers (13) and then use a suitable estimation for $R_n(t)$ on the circle.

Lemma 8. For all $j = 0, 1, \dots, q-1$,

$$d_{2n}^{q-1-j} \cdot 16^{rn} \cdot \frac{(rn)!^2}{(2n)!^r} \cdot P_{jn}(z) \in \mathbb{Z}[z]. \quad (27)$$

Proof. Using the decomposition

$$\begin{aligned} & \frac{(rn)!^2}{(2n)!^r} \cdot R_n(t) \left(t + k - \frac{1}{2}\right)^{q-1} \\ &= (2t + h_0) \cdot \prod_{j=1}^{r/2} G(2jn + 1, 2(j-1)n + 1; t) \\ & \quad \times \prod_{j=1}^{r/2} G((r+2j+1)n + 1, (r+2j-1)n + 1; t) \\ & \quad \times \left(H\left(rn + \frac{1}{2}, (r+1)n + \frac{3}{2}; t\right) \left(t + k - \frac{1}{2}\right)\right)^{q-1} \end{aligned}$$

for $k = rn + 1, \dots, (r+1)n + 1$ and then proceeding in the same way as in the proof of Lemma 6, we get Lemma 8. \square

We also need a simplified version of Lemma 3, given by the following

Lemma 9. With the above choice (26) of the parameters $\underline{\eta}$, the quantity (17) satisfies

$$0 < \varphi \leq \frac{2^{2r+1}}{r^{q-1}}.$$

Proof. By Lemma 3, $\varphi_{q,\underline{\eta}} > 0$ and since $1 + t_1 t_2 \cdots t_q \geq 1$ for $t_j \in [0, 1]$, we obtain

$$\begin{aligned} \varphi_{q,\underline{\eta}} &\leq \frac{(2r+1)^{2r+1}}{r^{2r}} \cdot \max_{t \in [0,1]^q} \prod_{j=1}^q t_j^r (1-t_j) = \frac{(2r+1)^{2r+1}}{r^{2r}} \cdot \left(\max_{t \in [0,1]} t^r (1-t)\right)^q \\ &\leq \frac{(2r+1)^{2r+1}}{r^{2r}} \cdot \frac{r^{qr}}{(r+1)^{q(r+1)}} \leq \frac{2^{2r+1}}{r^{q-1}}. \end{aligned}$$

\square

In order to obtain linear forms in 1 and even beta values, we assume from now on that q is even (as in Lemma 2).

Define, for $j = 0, 1, \dots, q/2 - 1$,

$$p_{jn} := 2^{q-2} d_{2n}^{q-1} 16^{rn} (rn)!^2 (2n)!^{-r} P_{2j,n}(-1);$$

from Lemma 8, $p_{jn} \in \mathbb{Z}$ for all j .

We also define

$$\ell_n := 2^{q-2} d_{2n}^{q-1} 16^{rn} (rn)!^2 (2n)!^{-r} S_n(-1).$$

According to Lemma 2,

$$\ell_n = p_{0n} + \sum_{j=1}^{q/2-1} p_{jn} \beta(2j) \quad (28)$$

is a linear form in 1 and even beta values with integral coefficients. Stirling's formula and (20) imply that

$$\lim_{n \rightarrow \infty} \left(2^{q-2} d_{2n}^{q-1} 16^{rn} \frac{(rn)!^2}{(2n)!^r} \right)^{1/n} = e^{2(q-1)} 4^r r^{2r}.$$

From Lemma 3 we deduce that

$$\lim_{n \rightarrow \infty} |\ell_n|^{1/n} = e^{2(q-1)} 4^r r^{2r} \varphi =: \kappa$$

and from Lemma 7 that for all $j = 0, \dots, q/2 - 1$,

$$\limsup_{n \rightarrow \infty} |p_{jn}|^{1/n} \leq e^{2(q-1)} 4^r 2^{q-2r-1} (2r+1)^{2r+1} =: \tau.$$

We can now apply Nesterenko's criterion to the linear form (28) with $N = q/2$, $\alpha_1 = \alpha_2 = \kappa$ and $\beta = \tau$: we get the lower bound

$$\begin{aligned} \delta(q-2) &\geq \frac{(q-2r-1) \log(2) + (2r+1) \log(2r+1) - 2r \log(r) - \log(\varphi)}{(2 + \log(2))(q-1) + (2r+1) \log(2r+1)} \\ &\geq \frac{q \log(r) + (q-2r-1) \log(2)}{(2 + \log(2))(q-1) + (2r+1) \log(2r+1)}. \end{aligned} \quad (29)$$

In the last step, we have used Lemma 9 and the fact that $\log(2r+1) \geq \log(2r)$. Now, as $q, r \rightarrow \infty$,

$$q \log(r) + (q-2r-1) \log(2) = q \log(r) + O(q) + O(r) = q \log(r) + O(q)$$

and

$$(2 + \log(2))(q-1) + (2r+1) \log(2r+1) = (2 + \log(2))q + O(r \log(r)).$$

Thus, for $r = 2 \lfloor q / \log^2(q) \rfloor$, we obtain

$$\delta(q-2) \geq \frac{(1 + o(1))q \log(q)}{(2 + \log(2))q + o(q)} = \frac{1 + o(1)}{2 + \log(2)} \log(q),$$

which proves the estimate (1) of Theorem 1. Here and in the rest of this article, $\lfloor \cdot \rfloor$ denotes the integral part of a real number. The existence of a constant c_0 , for which the inequality $\delta(a) > c_0 \log(a)$ always holds, immediately follows from the absolute lower bound (29) for the above choice of r .

7 p -adic valuation of the coefficients

Since $\text{ord}_p d_N = 1$ for those primes p such that $\sqrt{N} < p \leq N$, Lemma 6 gives that the p -adic order of the coefficients of the polynomials $P_{jn}(z)$ and $P_{0n}(z)$ is at least $-(q-1)$ if $\sqrt{m_1 n} < p \leq m_{q-1} n$. The aim of this paragraph is to sharpen these lower bounds for p -adic orders.

The next assertion is proved in [Zu2], Lemma 18.

Lemma 10. *Let a, b, a_0, b_0 be integers, $a_0 \leq a < b \leq b_0$, and let $H(t) = H(a, b; t)$ be defined as in (22). Then for any non-negative integer j , any integer k , $a_0 \leq k < b_0$, and any prime $p > \sqrt{b_0 - a_0 - 1}$, the following holds :*

$$\text{ord}_p(H(t)(t+k))\Big|_{t=-k}^{(j)} \geq -j + \left\lfloor \frac{b-a-1}{p} \right\rfloor - \left\lfloor \frac{k-a}{p} \right\rfloor - \left\lfloor \frac{b-1-k}{p} \right\rfloor.$$

The following observation will be useful later on: $\text{ord}_p N! = \lfloor N/p \rfloor$ for $p > \sqrt{N}$. An easy consequence of this fact is the formula giving the p -adic order of the numbers

$$\frac{\Gamma(N+1/2)}{\Gamma(1/2)} = \left(\frac{1}{2}\right)_N = 4^{-N} \frac{(2N)!}{N!}, \quad (30)$$

which are sometimes written as $(2N)!! \cdot 2^{-N}$; namely,

$$\text{ord}_p(2N)!! = \left\| \frac{N}{p} \right\| \quad \text{for } p > \sqrt{2N}, p \neq 2, \quad (31)$$

where $\|x\| := \lfloor 2x \rfloor - \lfloor x \rfloor$.

Lemma 11. *Let a, b, a_0, b_0 be integers, $b_0 \leq b < a \leq a_0$, and let $G(t) = G(a, b; t)$ be defined as in (21). Then for any non-negative integer j , any integer k , $b_0 < k < a_0$, and any prime $p > \sqrt{2(a_0 - b_0 - 1)}$, $p \neq 2$, the following holds :*

$$\begin{aligned} \text{ord}_p G^{(j)}\left(-k + \frac{1}{2}\right) &\geq -j + \left\| \frac{a-k}{p} \right\| - \left\| \frac{b-k}{p} \right\| - \left\lfloor \frac{a-b}{p} \right\rfloor \\ &= -j - \left\| \frac{k-a}{p} \right\| + \left\| \frac{k-b}{p} \right\| - \left\lfloor \frac{a-b}{p} \right\rfloor. \end{aligned} \quad (32)$$

Proof. By (30) we deduce that

$$4^{a-b} G\left(-k + \frac{1}{2}\right) = \begin{cases} \frac{(2a-2k)!(b-k)!}{(a-k)!(2b-2k)!(a-b)!} & \text{if } k \leq b, \\ (-1)^{k-b} \frac{(2k-2b)!(2a-2k)!}{(k-b)!(a-k)!(a-b)!} & \text{if } b < k < a, \\ (-1)^{a-b} \frac{(2k-2b)!(k-a)!}{(k-b)!(2k-2a)!(a-b)!} & \text{if } k \geq a. \end{cases}$$

Using (31) and the formula $\lfloor -N/p \rfloor = -\lfloor N/p \rfloor$ (for integers N and primes $p > 2$), we obtain the desired estimates (32) for $j = 0$. Inductive arguments, similar to that given in the proof of Lemma 17 in [Zu2], prove (32) for all j . \square

Applying Lemmas 10, 11 and the Leibniz rule (for differentiating a product) to the decomposition

$$R_{n,q,\eta}(t) = (2t + h_0) \cdot G(h_q, 1; t) \cdot G(h_0, 1 + h_0 - h_q; t) \\ \times \prod_{j=1}^{q-1} H(h_j, 1 + h_0 - h_j; t)$$

gives rise to the estimations

$$\text{ord}_p D_\lambda (R_n(t) (t + k - \frac{1}{2})^{q-1})|_{t=-k+\frac{1}{2}} \geq -\lambda + \varpi_0 \left(\frac{n}{p}, \frac{k-1}{p} \right) \\ \geq -\lambda + \varpi \left(\frac{n}{p} \right)$$

for $p > \max\{\sqrt{2\eta_0 n}, q-1\}$, $\lambda = 0, 1, \dots, q-1$, and $k = h_1 + \frac{1}{2}, \dots, h_0 - h_1 + \frac{1}{2}$, where

$$\varpi_0(x, y) := \sum_{j=1}^{q-1} (\lfloor (\eta_0 - 2\eta_j)x \rfloor - \lfloor y - \eta_j x \rfloor - \lfloor (\eta_0 - \eta_j)x - y \rfloor) \\ + \lfloor \eta_0 x - y \rfloor - \lfloor y - \eta_q x \rfloor + \lfloor y \rfloor - \lfloor (\eta_0 - \eta_q)x - y \rfloor - 2\lfloor \eta_q x \rfloor, \\ \varpi(x) := \min_{y \in \mathbb{R}} \varpi_0(x, y) = \min_{0 \leq y < 1} \varpi_0(x, y).$$

The function $\varpi(x)$ is non-negative, 1-periodic and integer-valued. Setting

$$\Pi_n := \prod_{\sqrt{2\eta_0 n} < p \leq m_{q-1} n} p^{\varpi(n/p)} \quad (33)$$

and recalling the formula (13), we finally obtain the following sharpening of Lemma 6.

Lemma 12. *For $j = 0, 1, \dots, q-1$, the following hold*

$$d_{m_1 n} d_{m_2 n} \cdots d_{m_{q-1} n} \cdot 16^{\eta_q n} \cdot \Pi_n^{-1} \cdot P_{jn}(z) \in \mathbb{Z}[z].$$

The asymptotic behaviour of (33) as $n \rightarrow \infty$ is given by (20); we refer the reader to [Ha] and [Zu1], Lemma 4.4, for details.

Lemma 13. *In the limit when $n \rightarrow +\infty$,*

$$\lim_{n \rightarrow \infty} \frac{\log \Pi_n}{n} = \int_0^1 \varpi(x) d\psi(x) - \int_0^{1/m_{q-1}} \varpi(x) \frac{dx}{x^2},$$

where $\psi(x) = \Gamma'(x)/\Gamma(x)$.

8 Proof of Theorem 2

Unlike Theorem 1, proving Theorem 2 only requires the following

Irrationality criterion. *Let $\theta_1, \dots, \theta_N$ be real numbers. Assume there exists N sequences of integers $(p_{ln})_{n \geq 0}$ such that the linear form $p_{1n}\theta_1 + p_{2n}\theta_2 + \dots + p_{Nn}\theta_N$ is non-zero for infinitely many n and tends to 0 as $n \rightarrow \infty$. Then at least one of the numbers $\theta_1, \dots, \theta_N$ is irrational.*

This clearly follows from Nesterenko's criterion but in this case the proof is completely elementary. Note also that here we do not need an estimation of the coefficients of the linear form.

Taking $q = 16$ and

$$\underline{\eta} = (101, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 45),$$

we obtain $t_0 \approx 0.00005445875955$. In this case, it is the unique zero in $[0, 1]$ of $Q(t)$; Further, $\varphi_{q, \underline{\eta}} \approx \exp(-594.58616762)$ and

$$\lim_{n \rightarrow \infty} \frac{\log(d_{m_1 n} d_{m_2 n} \cdots d_{m_{q-1} n} \cdot 16^{n_q n} \cdot \Pi_n^{-1})}{n} \approx 593.98582857$$

since

$$\begin{aligned} m_1 = 56, \quad m_2 = 54, \quad m_3 = 52, \quad m_4 = 50, \quad m_5 = 48, \quad m_6 = 46, \\ m_7 = \cdots = m_{15} = 45, \end{aligned}$$

and the function $\varpi(x)$ takes values in $\{0, 1, \dots, 9\}$. Thus, the irrationality criterion and Lemma 2 yield that at least one of the numbers (2) is irrational.

9 Special emphasis on Catalan's constant

When $q = 4$, our general construction gives us linear forms in 1 and Catalan's constant $G = \beta(2)$, and the corresponding well-poised hypergeometric series (5) for $z = -1$ can be reduced to a simpler series using Bailey's identity [Bai], Section 4.4, formula (2) :

$$\begin{aligned} {}_3F_2 \left(\begin{matrix} 1+a-b-c, d, e \\ 1+a-b, 1+a-c \end{matrix} \middle| 1 \right) &= \frac{\Gamma(1+a) \Gamma(1+a-d-e)}{\Gamma(1+a-d) \Gamma(1+a-e)} \\ &\times {}_6F_5 \left(\begin{matrix} a, 1+\frac{1}{2}a, b, c, d, e \\ \frac{1}{2}a, 1+a-b, 1+a-c, 1+a-d, 1+a-e \end{matrix} \middle| -1 \right), \end{aligned}$$

if $\operatorname{Re}(1 + a - d - e) > 0$. We obtain

$$\begin{aligned}
s_{n,\underline{\eta}} &:= S_{n,4,\underline{\eta}}(-1) \\
&= \frac{\Gamma(h_1)\Gamma(h_2)\Gamma(h_3)\Gamma(1+h_0-2h_1)\Gamma(1+h_0-2h_2)\Gamma(1+h_0-2h_3)}{\Gamma(1+h_0-h_1)\Gamma(1+h_0-h_2)\Gamma(1+h_0-h_3-h_4)\Gamma(h_4)} \\
&\quad \times {}_3F_2\left(\begin{matrix} 1+h_0-h_1-h_2, h_3, h_4 \\ 1+h_0-h_1, 1+h_0-h_2 \end{matrix} \middle| 1\right) \\
&= \frac{\Gamma(h_1)\Gamma(h_2)\Gamma(1+h_0-2h_1)\Gamma(1+h_0-2h_2)\Gamma(1+h_0-2h_3)}{\Gamma(1+h_0-h_1-h_3)\Gamma(1+h_0-h_2-h_4)\Gamma(1+h_0-h_3-h_4)\Gamma(h_4)^2} \\
&\quad \times \int_{[0,1]^2} \frac{x^{h_3-1}(1-x)^{h_0-h_1-h_3}y^{h_4-1}(1-y)^{h_0-h_2-h_4}}{(1-xy)^{h_0-h_1-h_2}} \frac{dx dy}{1-xy}.
\end{aligned} \tag{34}$$

Since the ${}_3F_2$ -series in (34) is invariant under any permutation of its top or bottom parameters, we deduce that the permutation of its top *integral* parameters $1 + h_0 - h_1 - h_2$ and h_4 produces the transformation

$$\mathfrak{b}: \underline{\eta} \mapsto (2\eta_0 - \eta_1 - \eta_2 - \eta_4, \eta_0 - \eta_2 - \eta_4, \eta_0 - \eta_1 - \eta_4, \eta_3, \eta_0 - \eta_1 - \eta_2) \tag{35}$$

under which

$$\frac{s_{n,\underline{\eta}}}{\prod_{j=1}^3 \Gamma(\eta_j n + \frac{1}{2}) \Gamma(1 + (\eta_0 - 2\eta_j)n) \Gamma(1 + (\eta_0 - \eta_1 - \eta_2 - \eta_3 + \eta_j)n)} \tag{36}$$

is invariant. We note that (36) is also stable under permutation of the non-integral parameters h_1, h_2, h_3 of the original well-poised ${}_6F_5$ -series. They are $6 = 3!$ such permutations and we can consider the group $\mathfrak{G} = \langle \mathfrak{b}, \mathfrak{S}_3 \rangle$ of the transformations of the parameters $\underline{\eta}$ that acts trivially on (36). This group has order 24 and is a subgroup of the permutation group of order 120 discovered by Rhin and Viola [RV] for $\zeta(2)$ (where permutations of h_1, h_2, h_3 , and h_4 are allowed, cf. [Zu2], Section 6). Unfortunately, the ‘denominators’ of the linear forms $s_{n,\underline{\eta}}$ given by Lemma 6 are too large to be able to use the arithmetic arguments in [RV], [Zu2].

Finally, we want to mention an arithmetic observation for linear forms in the case $\eta_0 = 3, \eta_1 = \eta_2 = \eta_3 = \eta_4 = 1$:

$$s_{n,\underline{\eta}} = \int_{[0,1]^2} \frac{x^{n-\frac{1}{2}}(1-x)^n y^n (1-y)^{n-\frac{1}{2}}}{(1-xy)^{n+1}} dx dy \in \mathbb{Q}G + \mathbb{Q}.$$

Thanks to Zeilberger’s creative telescoping algorithm, we have found (see [Zu4]) that both the sequence $(-1)^n s_{n,\underline{\eta}} = u_n G - v_n$ and the corresponding coefficients u_n and v_n satisfy the Apéry-like difference equation

$$(2n+1)^2(2n+2)^2 p(n)u_{n+1} - q(n)u_n - (2n-1)^2(2n)^2 p(n+1)u_{n-1} = 0, \tag{37}$$

where

$$\begin{aligned} p(n) &= 20n^2 - 8n + 1, \\ q(n) &= 3520n^6 + 5632n^5 + 2064n^4 - 384n^3 - 156n^2 + 16n + 7, \end{aligned}$$

with the initial data

$$u_0 = 4, \quad u_1 = 7, \quad v_0 = 0, \quad v_1 = \frac{13}{2}.$$

Using the recursion (37), we have verified numerically (up to $n = 1000$) that $16^n \cdot u_n \in \mathbb{Z}$ and $d_{2n}^2 \cdot 16^n \cdot v_n \in \mathbb{Z}$: this is more than one can get from Lemma 6, although it is not sufficient to prove the irrationality of Catalan's constant.

Acknowledgements. We would like to express our gratitude to S. Fischler, M. Waldschmidt and the anonymous referee for their helpful remarks which have much improved this article.

References

- [Ap] R. Apéry, Irrationalité de $\zeta(2)$ et $\zeta(3)$, *Astérisque* **61**, 11–13 (1979)
- [Bai] W. N. Bailey, *Generalized hypergeometric series*, Cambridge Math. Tracts **32** (Cambridge University Press, Cambridge 1935); 2nd reprinted edition (Stechert-Hafner, New York 1964)
- [Bak] A. Baker, The theory of linear forms in logarithms, *Transcendence theory: advances and applications* (Proc. Conf., Univ. Cambridge, Cambridge, 1976), pp. 1–27 (Academic Press, London, 1977)
- [BR] K. Ball, T. Rivoal, Irrationalité d'une infinité de valeurs de la fonction zêta aux entiers impairs, *Invent. Math.* **146**, no. 1, 193–207 (2001)
- [Ha] M. Hata, Legendre type polynomials and irrationality measures, *J. Reine Angew. Math.* **407**, no. 1, 99–125 (1990)
- [Ne] Yu. V. Nesterenko, On the linear independence of numbers, *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* no. 1, 46–54 (1985); English transl., *Moscow Univ. Math. Bull.* **40**, no. 1, 69–74 (1985)
- [RV] G. Rhin, C. Viola, On a permutation group related to $\zeta(2)$, *Acta Arith.* **77**, no. 1, 23–56 (1996)

- [Ri1] T. Rivoal, La fonction zêta de Riemann prend une infinité de valeurs irrationnelles aux entiers impairs, C. R. Acad. Sci. Paris Sér. I Math. **331**, no. 4, 267–270 (2000)
- [Ri2] T. Rivoal, Irrationalité d’au moins un des neuf nombres $\zeta(5), \zeta(7), \dots, \zeta(21)$, Acta Arith. **103**, 157–167 (2002)
- [Sl] L. J. Slater, *Generalized hypergeometric functions*, 2nd edition (Cambridge University Press, Cambridge 1966)
- [Zu1] W. Zudilin, Irrationality of values of Riemann’s zeta function, Izv. Ross. Akad. Nauk Ser. Mat. **66**, no. 3, 49–102 (2002); English transl., Russian Acad. Sci. Izv. Math. **66**, no. 3 (2002)
- [Zu2] W. Zudilin, Arithmetic of linear forms involving odd zeta values, Preprint (August 2001); E-print math.NT/0206176, 42 pages (2002)
- [Zu3] V. V. Zudilin, One of the numbers $\zeta(5), \zeta(7), \zeta(9), \zeta(11)$ is irrational, Uspekhi Mat. Nauk **56**, no. 4, 149–150 (2001); English transl., Russian Math. Surveys **56**, no. 4, 774–776 (2001)
- [Zu4] W. Zudilin, Apéry-like difference equation for Catalan’s constant, E-print math.NT/0201024, 10 pages (2002)

Tanguy Rivoal
 Institut de Mathématiques de Jussieu
 CNRS UMR 7586, Théorie des Nombres, case 247
 175, rue du Chevaleret
 75013 Paris, FRANCE
 email : rivoal@math.jussieu.fr

Wadim Zudilin
 Moscow Lomonosov State University
 Department of Mechanics and Mathematics
 Vorobiovy Gory, GSP-2
 119992 Moscow, RUSSIA
 email : wadim@ips.ras.ru