

Homework 9

The homeworks are due on the Thursday of the week after the assignment was posted online¹. Please hand in your homework at the beginning of the tutorial or bring it to the lecture on Thursday morning. You can work on and submit your homework in groups of two. Please staple your pages and write your names and matriculation numbers on the first page.

Problem 25 (10 pts.)

- a) If L/K is Galois and w is an extension of v to L , then $w \circ \sigma$ is an extension of v for all $\sigma \in \text{Gal}(L/K)$.
- b) Let L/K be a finite Galois extension and let K be complete with respect to the exponential valuation v . Give a new proof (different from the one given in class) that $w(x) = \frac{1}{n}v(N_{L/K}(x))$ is the unique extension of v to L .
Hint: You can use, as seen in class, that there is at least one extension of v to L , and you should use part a) to prove b).

Problem 26 (10 pts.)

Let K be a local field with discrete valuation v . Let $f \in K[X]$ given by

$$f(X) = a_n X^n + \cdots + a_1 X + a_0$$

with $a_0 a_n \neq 0$.

First look up the definition of the Newton polygon of f from Exercise 27 below.

Let $L = K(\alpha_1, \dots, \alpha_n)$ be the splitting field of f over K . We also write v for the unique extension of the valuation from K to L . Let us suppose that $\alpha_1, \dots, \alpha_n$ are ordered, such that

$$\begin{aligned} v(\alpha_1) &= \cdots = v(\alpha_{s_1}) = m_1 \\ v(\alpha_{s_1+1}) &= \cdots = v(\alpha_{s_2}) = m_2 \\ &\quad \dots = \dots \\ v(\alpha_{s_{r-1}+1}) &= \cdots = v(\alpha_{s_r}) = m_r \end{aligned}$$

with $m_1 < m_2 < \dots < m_r$.

In the following, we suppose that $a_n = 1$ for simplicity. This is not a restriction, as the main point c) remains valid if $a_n \neq 1$ (because multiplication with a constant might only shift the Newton polygon up or down).

¹This assignment is due Thursday, 12.12.19.

a) Show that

$$\begin{aligned}
 v(a_n) &= 0 \\
 v(a_{n-1}) &\geq m_1 \\
 v(a_{n-2}) &\geq 2m_1 \\
 &\dots \\
 v(a_{n-s_1}) &= s_1 m_1 \\
 v(a_{n-s_1-1}) &\geq s_1 m_1 + m_2 \\
 &\dots \\
 v(a_{n-s_2}) &= s_1 m_1 + (s_2 - s_1)m_2 \\
 &\dots \\
 v(a_{n-s_2-1}) &\geq s_1 m_1 + (s_2 - s_1)m_2 + m_3 \\
 &\dots \\
 v(a_0) &= s_1 m_1 + (s_2 - s_1)m_2 + (s_3 - s_2)m_3 + \dots + (s_r - s_{r-1})m_r.
 \end{aligned}$$

Hint: Look up formulas for the coefficients a_i in terms of elementary symmetric polynomials in the roots α_j in an algebra book.

b) Show that the vertices of the Newton polygon are (from right to left) exactly the points

$$(n, 0), (n - s_1, s_1 m_1), (n - s_2, s_1 m_1 + (s_2 - s_1)m_2), \dots,$$

and the slopes of the lines connecting these vertices are exactly $-m_1, -m_2, \dots, -m_r$.

c) Summarize: If $(k, v(a_k)) \leftrightarrow (\ell, v(a_\ell))$ is a line ($k > \ell$) of the Newton polygon with slope $-m$, then f has exactly $\ell - k$ roots with valuation m .

Problem 27 (10 pts.)

We use the same notation as in Problem 26 and suppose that K is a local field of characteristic 0. Show that there is a decomposition

$$f(X) = a_n \prod_{j=1}^r f_j(X)$$

with

$$f_j(X) = \prod_{i=s_{j-1}+1}^{s_j} (X - \alpha_i) \in K[X]$$

and with $s_0 = 0$. In particular: If the Newton polygon has n different slopes, then f splits completely over K . If the polynomial f is irreducible, then the Newton polygon has only one slope.

Hints:

- Recall that if L/K is the splitting field of f , then L/K is a Galois extension.
- First suppose that f is irreducible.
- Then give a proof by induction on the degree n .

Bonus Problem (10 pts.)

Let K be a finite extension of \mathbb{Q}_p of degree n . Let \mathcal{O}_K be its valuation ring, \mathfrak{p} its maximal ideal and $\mathbb{k} = \mathcal{O}_K/\mathfrak{p}$ its residue field.

Let e be the *ramification index* defined in Homework 8 and let f be the *degree of inertia*, i.e., the degree of the extension of finite fields $[\mathbb{k} : \mathbb{F}_p]$.

Prove the *characteristic equation*:

$$n = e \cdot f.$$

(This shows that the degree $n = [K : \mathbb{Q}_p]$ of a finite extension K of \mathbb{Q}_p breaks up as a product $n = e \cdot f$, where e “measures” the change of the image of the p -adic valuation ν_p and $f = [\mathbb{k} : \mathbb{F}_p]$ the change in the residue field.)

The following exercises will be discussed in the tutorial and you do not need to hand in solutions for them.

Exercise 27

Newton polygons.

Let K be a local field with discrete valuation v . Let $f \in K[X]$ given by

$$f(X) = a_n X^n + \dots + a_1 X + a_0$$

with $a_0 a_n \neq 0$. Consider the set of points $S = \{(i, v(a_i)) \in \mathbb{R}^2 : i = 0, \dots, n, a_i \neq 0\}$. The Newton polygon of f is then defined to be the lower boundary of the convex hull of S .

Draw the Newton polygon of each of the following two polynomials, compute the slopes and illustrate Problem 26, c) in these cases.

1. $f(X) = 5^6 X^4 + 5^3 X^3 - 5X^2 - X + \frac{1}{7} \in \mathbb{Q}_5[X]$
2. $f(X) = 5^2 X^5 - \sqrt{5} X^3 + 7X^2 - 1 \in \mathbb{Q}_5(\sqrt{5})[X]$